

Preface

Who is This Book For?

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Why Write Yet Another Book About Linear Algebra?

The direct answer to this question is simple: I really think most resources are doing a *bad job* of explaining fundamental linear algebra to new students ¹. Not because they lack material or the complete picture - but because the common approach to teaching fundamental mathematical topics just doesn't work well for most people ².

Introduction courses to fundamental mathematical topics usually look like this: here're some basic definitions, let's derive some specific results from them - now go apply these in your field.

¹and honestly, this isn't restricted to linear algebra but applies to all fundamental mathematical topics - but it's a classic case

²the main exception is students of pure mathematics and people who can easily deal with extremely abstract concepts - a very specific group of people at the end of the day

Preface

Who is This Book For?

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Why Write Yet Another Book About Linear Algebra?

The direct answer to this question is simple: I really think most resources are doing a *bad job* of explaining fundamental linear algebra to new students ³. Not because they lack material or the complete picture - but because the common approach to teaching fundamental mathematical topics just doesn't work well for most people ⁴.

Introduction courses to fundamental mathematical topics usually look like this: here're are some basic definitions, let's derive some specific results from them - now go apply these in your field.


³and honestly, this isn't restricted to linear algebra but applies to all fundamental mathematical topics - but it's a classic case

⁴the main exception is students of pure mathematics and people who can easily deal with extremely abstract concepts - a very specific group of people at the end of the day

CHAPTER

1

MATHEMATICAL BACKGROUND



1.1 PREFACE

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Testing some equations: let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and $\alpha, \beta \in \mathbb{R}$. Then from the fact that $\mathbb{R}^n$ is a vector space, we get

$$\vec{c} = \alpha\vec{x} + \beta\vec{y} \in \mathbb{R}^n. \quad (1.1)$$

However, if $\vec{a} \in \mathbb{R}^m$, where $m \neq n$, then $\alpha\vec{x} \in \mathbb{R}^m$, and therefore $\vec{c}$ is not defined (i.e. Equation 1.1 is irrelevant).

Now, this is a test of the *aligned* environment, which allows for multiple (correctly-aligned) lines of equations and having the label only once (instead of per line, as in the *align* environment):

$$\begin{aligned} \sin(2x) &= \frac{1}{2i} \left(e^{2xi} - e^{-2xi} \right) \\ &= \frac{1}{2i} \left(\left[e^{xi} \right]^2 - \left[e^{-xi} \right]^2 \right) \\ &= \underbrace{\frac{1}{2i} \left(e^{xi} - e^{-xi} \right)}_{\sin(x)} \underbrace{\left(e^{xi} + e^{-xi} \right)}_{2 \cos(x)} \\ &= 2 \sin(x) \cos(x). \end{aligned} \quad (1.2)$$

Note 1.1 Test note

This is a test note. I will try to also refernce it later.



Now, I wish to reference [Note 1.1](#). Did it work?

Also, let us look at an example.

Example 1.1 Some test example

If $x = \begin{bmatrix} 1 & 2 & -3 \end{bmatrix}$ and $y = \begin{bmatrix} 5 & -1 \end{bmatrix}$, then $n = 3$ and $m = 2$. Clearly, $n \neq m$ - and so $\alpha \vec{x} \in \mathbb{R}^3$, while $\beta \vec{y} \in \mathbb{R}^2$. This means that $\alpha \vec{x} + \beta \vec{y}$ is not defined.



...let's check if referencing an example works correctly: can you see [Example 1.1](#)?

I believe it is now time for some figures! See [Figure 1.1](#).

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.




Figure 1.1: This is the first test figure.
Does it look ok?

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo

ger sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo

CHAPTER 1. MATHEMATICAL BACKGROUND

ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.




Figure 1.2: And this is another test figure! I hope it works.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tin-

1.1. PREFACE

cidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti




Figure 1.3

sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.3

Test text bla bla bla 0_preface.tex./chapters/1_mathsBackground0_preface.tex

$$T(\alpha \vec{x} + \beta \vec{y}) = \alpha T(\vec{x}) + \beta T(\vec{y})$$

$$\langle \vec{a} | \vec{b} \rangle = \vec{a} \cdot \vec{b} = \sum_{i=1}^n a_i b_i$$

$$\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} \perp \vec{b}$$

$$\mathbb{I}_n = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$


$$\vec{v} = \alpha_1 \hat{e}_1 + \alpha_2 \hat{e}_2 + \dots + \alpha_n \hat{e}_n$$

$$\mathbf{A} \vec{x} = \lambda \vec{x}$$

$$(\mathbf{A} \cdot \mathbf{B})^T = \mathbf{B}^T \cdot \mathbf{A}^T$$

$$\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$$

$$\mathbf{C} = \mathbf{A} \cdot \mathbf{B} \Rightarrow c_j^i = \langle a^i | b_j \rangle$$



CHAPTER

2

THE BASICS OF LINEAR ALGEBRA

2.1 INTRODUCTION AND MOTIVATION

2.2 VECTORS AND VECTOR SPACES

2.2.1 MOTIVATION

Motivating Problem 2.1 Gravity Force

Consider a simple simulation of classical (“Newtonian”) gravity, say of a planet with mass m orbiting a star with mass M (where $M \gg m$). At any given moment, the force acting on the planet due to the star’s gravity has magnitude

$$F = G \frac{Mm}{r^2}, \quad (2.1)$$

where r is the distance between the planet and the star at that moment, and G is a universal constant (specifically, $G = 6.674 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$).

However, since we live in 3-dimensional space^a, we also want this force to be pointing in the direction from the planet to the star. To do this, we need to understand how to describe orientations in 3-dimensional space, and be able to pack together both orientation and magnitude in the correct way.

^aallegedly.




Motivating Problem 2.2 Bouncing/Reflecting Off a Wall

Imagine we are writing a 3D computer game in which some object is bouncing off a wall. Alternatively, consider 3D graphics in general, where one simulates rays of light reflecting off of smooth surfaces. In both cases the same problem needs to be solved: given a flat plane of some orientation in space, and an object hitting it from some direction - what would be the direction of the object after it bounces off the plane?

continues in the next page ➡

 continues from the previous page



To solve this problem we will need to know how to represent quantities like position, velocity and orientation in 3D, and more generally points, lines and planes - and to know how to calculate the intersection between these different objects. By the end of this section you should be able to solve this problem by yourself, and even generalize it to any number of dimensions.



Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

2.2.2 VECOTRS AND THEIR GEOMTERIC PROPERTIES


The above motivating problems require objects that have both a magnitude and a direction: for example, the force of gravity between a star and a planet has magnitude given by [Equation 2.1](#) and a direction from the planet to the star. In [Motivating-problem 2.2](#), if we model some object hitting a wall, the object's

2.2. VECTORS AND VECTOR SPACES


velocity is a combination of magnitude (which we usually simply call speed) and a direction (towards the wall).

In the simplest terms, **vectors** are mathematical objects which have both a magnitude and a direction. The magnitude of a vector is called **norm**, while its direction is sometimes referred to as its **orientation**. Vectors can exist in 2-dimensions, 3-dimensions or even higher dimensions. Since higher dimensions are rather hard to conceptualize, I will first focus on 2- and 3-dimensional vectors so that we can better understand their properties, and only then generalize to higher dimensions when it's relevant.

In 2-dimensions we can represent vectors as arrows:




However, since we are usually interested in the magnitude and direction of vectors and not where they originate from, we draw them as originating in the same location which we unimagatively call the **origin**:



In fact, all vectors that have the same norm and orientation are considered identical - no matter where they originate. For example, the following vectors are actually all identical:

CHAPTER 2. THE BASICS OF LINEAR ALGEBRA



Since we will be using vectors as mathematical objects and would like to use them in an arithmetical context, we need to use some kind of notation. Usually, vectors are denoted as lower-case Latin letters, either in boldface or with a line or an arrow symbol above or below the letter. Here are few examples:

$$\begin{aligned} &\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{x}, \mathbf{y}, \mathbf{z}, \dots \\ &\bar{a}, \bar{b}, \bar{c}, \bar{x}, \bar{y}, \bar{z}, \dots \\ &\underline{a}, \underline{b}, \underline{c}, \underline{x}, \underline{y}, \underline{z}, \dots \\ &\vec{a}, \vec{b}, \vec{c}, \vec{x}, \vec{y}, \vec{z}, \dots \\ &\vec{\mathbf{a}}, \vec{\mathbf{b}}, \vec{\mathbf{c}}, \vec{\mathbf{x}}, \vec{\mathbf{y}}, \vec{\mathbf{z}}, \dots \end{aligned}$$


In this book I am using the last notation - a boldface Latin letter with an arrow above it. This ensures that it is both easy to read and is distinctive enough to quickly differentiate vectors from other objects. Sometimes, depending on context, the vectors are colored to make it easier to follow through (the process of setting up/developing a concept/equation).

There are two fundamental operations involving vectors: addition and scaling. These two operations result in creating a new vector of the same dimensions: adding together two 2-dimensional vectors results in a 2-dimensional vector, scaling a 3-dimensional vector result in a 3-dimensional vector, etc.

2.2. VECTORS AND VECTOR SPACES


Let's have a closer look at these two fundamental vector operations:

- **Adding vectors together:** any two vectors of the same dimensionality (e.g. 2-dimensional, 3-dimensional, 4-dimensional, etc.) can be added together, and the result is a third vector. Pictorially, here are two vectors $\vec{a}$ and $\vec{b}$ which we want to sum together:




The steps to produce the sum of $\vec{a}$ and $\vec{b}$ is the following:

1. move $\vec{b}$ such that its origin aligns with the head of $\vec{a}$:




2. draw a new vector from the origin to the head of the shifted $\vec{b}$:




CHAPTER 2. THE BASICS OF LINEAR ALGEBRA

3. the resulting vector $\vec{c}$ is the sum of $\vec{a}$ and $\vec{b}$. Here are all three vectors together from the same origin:




This method is called the **triangle method**, and it displays an important property of vector addition: it is commutative! This means that the order of addition doesn't matter: $\vec{a} + \vec{b} = \vec{b} + \vec{a}$. To see this visually, we can apply the triangle method twice, i.e. moving both $\vec{a}$ and $\vec{b}$:



We see that no matter which vector we add onto which (i.e. what is the order of addition) - the result is the same. The shape formed by this demonstration is called a **parallelogram**, and it is incredibly important in linear algebra as a whole as we will see in future sections.


This method of summing vectors can be applied to any number of vectors. For example, consider the following 5 vectors:




To sum them together, all we need is to start with one of the vectors (say, the red

2.2. VECTORS AND VECTOR SPACES


one) and use the triangle method to add the other vectors step by step:



Finally, we connect the origin of the first vector with the head of the last vector (in this case the orange one):



and that's it, the dashed gray vector is the sum of all five vectors. To see that the order of addition really doesn't matter we can do this again but this time starting with the green vector and changing the summation order completely:



As can be seen, the resulting dashed gray vector is exactly the same as before.

Another important property of the addition of vectors is that we can always take a vector and rotate it by 180° , giving us a vector that “cancels” the original vector by summation. For example, the following red vector is rotated to yield its opposite, colored in orange:



These vectors added together using the triangle method yield a vector with 0 norm:




(here I drew the vectors a bit shifted from each other so that we can see both)

Such a vector is called the **zero vector**, denoted as $\vec{0}$. It has no direction, and a norm of 0. Adding $\vec{0}$ to any other vector is the same as not adding any vector at all (and it doesn't matter "from which side" we add $\vec{0}$):

$$\vec{a} + \vec{0} = \vec{0} + \vec{a} = \vec{a}. \quad (2.2)$$

We say that the zero vector is *agnostic* (also: neutral) to addition.

• **Scaling vectors:** any vector can be **scaled** by any real number, which in this context we call a **scalar**. Scaling by a number simply "stretches" the vector by the number. For example, in the following figure a red vector is scaled by a scalar λ , taking on different values:




This example shows us several things:

- no matter by which positive scalar we scale a vector, it always points in the same direction.
- If we scale a vector by 0 it becomes the zero vector.
- Finally, scaling a vector by a negative scalar rotates the vector by 180° and then scales it by the absolute value of the scalar.

This is similar to how real numbers behave under multiplication: multiplying a real number by a negative number is, in a sense, "flipping" and scaling it relative to 0 (the origin). For example, multiplying 3 by -2 flips the the number to point to the left, and then scales it to be of length 6:

2.2. VECTORS AND VECTOR SPACES



Therefore, scaling a vector is denoted just like multiplication of real numbers: $\lambda \vec{a}$, where λ is the scalar. In the above example, if the red vector is $\vec{a}$, then the other five vectors can be written as $2\vec{a}$, $0\vec{a}$, $\frac{1}{2}\vec{a}$, $-1\vec{a}$ and $-\frac{3}{2}\vec{a}$.


Furthermore, we can place the vector notation in different places and even hiding the scalar depending on convenience, so for example we can write $-\vec{a}$ instead of $-1\vec{a}$ and $-\frac{3\vec{a}}{2}$ instead of $-\frac{3}{2}\vec{a}$.

Just like with real numbers, we call $-\vec{a}$ the **additive inverse** of $\vec{a}$.

In fact, the similarities to real number don't end there: subtracting a vector $\vec{b}$ from another vector $\vec{a}$ is simply adding $\vec{a}$ and the additive inverse of $\vec{b}$:

$$\vec{a} - \vec{b} = \vec{a} + (-\vec{b}). \quad (2.3)$$

Visually, if we place the origin of $\vec{a} - \vec{b}$ at the head of $\vec{b}$, its head would be at the same point as the head of $\vec{a}$:



While subtraction is not commutative, it is **antisymmetric**:

$$\vec{a} - \vec{b} = -(-\vec{a} + \vec{b}) = -(\vec{b} + (-\vec{a})) = -(\vec{b} - \vec{a}), \quad (2.4)$$

i.e. flipping the order of subtraction changes the sign of the expression.


So we can use vector subtraction as a convenient way to point from one vector to another. This can be utilized to implement the gravity force in [Motivating-problem 2.1](#): if we know the position of the planet (let's call it $\vec{r}_p$) and the position

CHAPTER 2. THE BASICS OF LINEAR ALGEBRA

of the star ($\vec{\mathbf{r}}_s$), a vector pointing from the planet's center to the star's center would be, for example,

$$\vec{\mathbf{r}}_{p \rightarrow s} = \vec{\mathbf{r}}_p - \vec{\mathbf{r}}_s. \quad (2.5)$$

(the notation $p \rightarrow s$ is just an easy way to represent that the vector is pointing from the planet to the star)



However, this vector has a norm equal to the distance between the planet and the star - and we want to use it to calculate the *force* acting on the planet, which has a different magnitude (and units). To do this, we need to be able to scale $\vec{\mathbf{r}}_{p \rightarrow s}$ to whatever norm we want. Taking inspiration from real numbers, we can employ the following procedure: say we want to scale some non-zero number x to be y . We can multiply x by $\frac{y}{x}$, canceling the x :

$$x \mapsto \cancel{x} \cdot \frac{y}{\cancel{x}} = y.$$

A different way to phrase this is that we simply multiply (or *scaling*) x by $\frac{1}{x}$, and then multiply the result by y :

$$x \mapsto \cancel{x} \cdot \frac{1}{\cancel{x}} \cdot y = y.$$

We can apply the same technique to vectors. Scaling a vector by its *reciprocal norm* yields a vector with norm 1, which we denote by replacing the arrow in the vector notation by a “hat”:

$$\hat{\mathbf{a}} = \frac{1}{\|\vec{\mathbf{a}}\|} \vec{\mathbf{a}}. \quad (2.6)$$

This operation is called **normalization**, and the resulting vector is a **unit vector** (also: normalized vector).

2.2. VECTORS AND VECTOR SPACES

Now, if we scale a unit vector $\hat{\mathbf{a}}$ by any scalar λ we get a vector of the same direction as $\hat{\mathbf{a}}$ (and also $\vec{\mathbf{a}}$), and with a norm of λ . The vectors $\vec{\mathbf{a}}$, $\hat{\mathbf{a}}$ and $\lambda\hat{\mathbf{a}}$ are all parallel to each other.

Back to our star-planet system: by normalizing the vector $\vec{\mathbf{r}}_{p \rightarrow s}$ we get a unit vector pointing from the planet to the star: $\hat{\mathbf{r}}_{p \rightarrow s}$, and scaling by the magnitude of the gravitational attraction (Equation 2.1) we finally get the correct *gravity force vector*:


$$\begin{array}{c} \boxed{\vec{\mathbf{F}}} = \boxed{G \frac{Mm}{r^2}} \boxed{\hat{\mathbf{r}}_{p \rightarrow s}} \\ \uparrow \qquad \qquad \uparrow \qquad \qquad \uparrow \\ \text{vector} \qquad \text{scalar} \qquad \text{vector} \end{array} \quad (2.7)$$

Indeed, $\vec{\mathbf{F}}$ is a vector: it has a magnitude (the scalar $G \frac{Mm}{r^2}$) and a direction (the unit vector $\hat{\mathbf{r}}_{p \rightarrow s}$). Any vector can be written as a unit vector scaled by a real number:

$$\vec{\mathbf{a}} = a\hat{\mathbf{a}}, \quad (2.8)$$

where $a \in \mathbb{R}$ is the magnitude of $\vec{\mathbf{a}}$, i.e. $a = \|\vec{\mathbf{a}}\|$. This is nothing more than a rearrangement of Equation 2.6: by dividing both sides of the equation by $a = \|\vec{\mathbf{a}}\|$ we get the definition of a unit vector.

TBW: text about how in 3D the same rules apply.



2.2.3 COMMON COORDINATE SYSTEMS IN 2- AND 3-DIMENSIONS

Everything we discussed so far is well and good - but it doesn't give us much in the way of actually doing any practical calculations. To do that, we need to "quantify" vectors somehow. There are many ways to assign values to vectors, and for now we will concentrate on the most common ones: two systems in 2-dimensions and three systems in 3-dimensions.

2.2.4 GEOMETRIC OBJECTS

2.2.5 PROJECTIONS AND REJECTIONS

2.2.6 VECTOR SPACES

2.2.7 SOLVING THE MOTIVATING PROBLEMS

2.3 LINEAR TRANSFORMATIONS

2.3.1 MOTIVATION

2.3.2 LINEAR TRANSFORMATIONS IN 2-DIMENSIONS

2.3.3 LINEAR TRANSFORMATIONS IN 3-DIMENSIONS

2.3.4 THE FORMAL DEFINITION OF LINEAR TRANSFORMATIONS

2.3.5 REPRESENTING LINEAR TRANSFORMATIONS, PART 1

2.4 MATRICES

2.4.1 REPRESENTING LINEAR TRANSFORMATIONS, PART 2

2.4.2 MATRIX REPRESENTATIONS OF 2- AND 3-DIMENSIONAL LINEAR TRANSFORMATIONS

2.4.3 TYPES OF MATRICES

2.4.4 MATRIX-MATRIX PRODUCT

2.4.5 MATRIX DETERMINANTS

2.5 LINEAR EQUATIONS

2.6 EIGENVECTORS AND EIGENVALUES

2.7 MATRIX DECOMPOSITIONS

$$T(\alpha \vec{x} + \beta \vec{y}) = \alpha T(\vec{x}) + \beta T(\vec{y})$$

$$\langle \vec{a} | \vec{b} \rangle = \vec{a} \cdot \vec{b} = \sum_{i=1}^n a_i b_i$$

$$\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} \perp \vec{b}$$

$$\mathbb{I}_n = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$


$$\vec{v} = \alpha_1 \hat{e}_1 + \alpha_2 \hat{e}_2 + \dots + \alpha_n \hat{e}_n$$

$$\mathbf{A} \vec{x} = \lambda \vec{x}$$

$$(\mathbf{A} \cdot \mathbf{B})^T = \mathbf{B}^T \cdot \mathbf{A}^T$$

$$\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$$

$$\mathbf{C} = \mathbf{A} \cdot \mathbf{B} \Rightarrow c_{ij} = a_i \cdot b_j$$



CHAPTER

3

INTRODUCING SOME FORMALISM

$$T(\alpha \vec{x} + \beta \vec{y}) = \alpha T(\vec{x}) + \beta T(\vec{y})$$

$$\langle \vec{a} | \vec{b} \rangle = \vec{a} \cdot \vec{b} = \sum_{i=1}^n a_i b_i$$

$$\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} \perp \vec{b}$$

$$\mathbb{I}_n = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$


$$\vec{v} = \alpha_1 \hat{e}_1 + \alpha_2 \hat{e}_2 + \dots + \alpha_n \hat{e}_n$$

$$\mathbf{A} \vec{x} = \lambda \vec{x}$$

$$(\mathbf{A} \cdot \mathbf{B})^T = \mathbf{B}^T \cdot \mathbf{A}^T$$

$$\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$$

$$\mathbf{C} = \mathbf{A} \cdot \mathbf{B} \Rightarrow c_{ij} = a_i \cdot b_j$$



CHAPTER

4

ADVANCED TOPICS IN LINEAR ALGEBRA

$$T(\alpha \vec{x} + \beta \vec{y}) = \alpha T(\vec{x}) + \beta T(\vec{y})$$

$$\langle \vec{a} | \vec{b} \rangle = \vec{a} \cdot \vec{b} = \sum_{i=1}^n a_i b_i$$

$$\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} \perp \vec{b}$$

$$\mathbb{I}_n = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$


$$\vec{v} = \alpha_1 \hat{e}_1 + \alpha_2 \hat{e}_2 + \dots + \alpha_n \hat{e}_n$$

$$\mathbf{A} \vec{x} = \lambda \vec{x}$$

$$(\mathbf{A} \cdot \mathbf{B})^T = \mathbf{B}^T \cdot \mathbf{A}^T$$

$$\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$$

$$\mathbf{C} = \mathbf{A} \cdot \mathbf{B} \Rightarrow c_{ij} = a_i \cdot b_j$$



CHAPTER

5

FURTHER RELATED TOPICS

CHAPTER 5. FURTHER RELATED TOPICS

Actual page layout values.

<code>\paperheight = 845.04684pt</code>	<code>\paperwidth = 597.50787pt</code>
<code>\hoffset = 0.0pt</code>	<code>\voffset = 0.0pt</code>
<code>\evensidemargin = 35.28255pt</code>	<code>\oddsidemargin = -0.56833pt</code>
<code>\topmargin = -10.73834pt</code>	<code>\headheight = 18.125pt</code>
<code>\headsep = 21.75pt</code>	<code>\textheight = 591.5302pt</code>
<code>\textwidth = 418.25368pt</code>	<code>\footskip = 50.75pt</code>
<code>\marginparsep = 12.8401pt</code>	<code>\marginparpush = 6.52495pt</code>
<code>\columnsep = 10.0pt</code>	<code>\columnseprule = 0.0pt</code>
<code>lem = 12.0pt</code>	<code>lex = 5.172pt</code>