### **Basic Maths for Non-mathematicians**

#### Peleg Bar Sapir

$$\int_{a}^{b} f(x) dx = \lim_{\Delta x \to 0} \sum_{k=1}^{N} f(x_{k}) \Delta x$$

$$(AB)^{\top} = B^{\top} A^{\top} \qquad \mathbb{R}^{n} \xrightarrow{T} \mathbb{R}^{m}$$

$$\vec{v} = \sum_{i=1}^{n} \alpha_{i} \hat{e}_{i}$$


$$\frac{df}{dx} = \lim_{\Delta x \to 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \qquad A = Q\Lambda Q^{-1}$$

$$\operatorname{Rot}(\theta) = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \int_{a}^{b} f(x) dx = F(b) - F(a)$$

$$T(\alpha \vec{u} + \beta \vec{v}) = \alpha T(\vec{u}) + \beta T(\vec{v}) \quad \langle \hat{e}_{i}, \hat{e}_{j} \rangle = \delta_{ij}$$



# **Chapter 3: Linear Transformations**



#### **Definition**

A **linear transformation** is a function  $T:A\to B$ , that obeys the following two criteria:

#### **Definition**

- A **linear transformation** is a function  $T:A\to B$ , that obeys the following two criteria:
  - 1. **Scalability**: for each  $x \in A$  and a scalar  $\alpha \in \mathbb{R}$ :

$$T(\alpha x) = \alpha T(x).$$

#### **Definition**

A **linear transformation** is a function  $T:A\to B$ , that obeys the following two criteria:

1. **Scalability**: for each  $x \in A$  and a scalar  $\alpha \in \mathbb{R}$ :

$$T(\alpha x) = \alpha T(x).$$

2. **Additivity**: For any  $x, y \in A$ :

$$T(x+y) = T(x) + T(y).$$

### **Example**

The real function f(x)=3x is linear. Proof by the above criteria:

#### **Example**

The real function f(x)=3x is linear. Proof by the above criteria:

1. **Scalability**: for any scalar  $\alpha \in \mathbb{R}$ ,

$$f(\alpha x) = 3(\alpha x) = \alpha \cdot (3x) = \alpha f(x).$$

### **Example**

The real function f(x) = 3x is linear. Proof by the above criteria:

1. **Scalability**: for any scalar  $\alpha \in \mathbb{R}$ ,

$$f(\alpha x) = 3(\alpha x) = \alpha \cdot (3x) = \alpha f(x).$$

2. **Additivity**: for any two numbers  $x, y \in \mathbb{R}$ 

$$f(x+y) = 3(x+y) = 3x + 3y = f(x) + f(y).$$

3

#### **Example**

The real function f(x) = 3x is linear. Proof by the above criteria:

1. **Scalability**: for any scalar  $\alpha \in \mathbb{R}$ ,

$$f(\alpha x) = 3(\alpha x) = \alpha \cdot (3x) = \alpha f(x).$$

2. **Additivity**: for any two numbers  $x, y \in \mathbb{R}$ 

$$f(x+y) = 3(x+y) = 3x + 3y = f(x) + f(y).$$

Therefore, f is linear.

### **Example**

Is the real function g(x) = 3x + 5 linear? Let's check:

#### **Example**

Is the real function g(x) = 3x + 5 linear? Let's check:

1. Scalability:

$$g(\alpha x) = 3\alpha x + 5, \quad \alpha g(x) = 3x\alpha + 5\alpha.$$

#### **Example**

Is the real function g(x) = 3x + 5 linear? Let's check:

1. Scalability:

$$g(\alpha x) = 3\alpha x + 5, \quad \alpha g(x) = 3x\alpha + 5\alpha.$$

If we subtitute  $\alpha=0, x=1$ , for example, we get

$$g(\alpha x) = g(0 \cdot 1) = 3 \cdot 0 + 5 = 5,$$

#### **Example**

Is the real function g(x) = 3x + 5 linear? Let's check:

#### 1. Scalability:

$$g(\alpha x) = 3\alpha x + 5, \quad \alpha g(x) = 3x\alpha + 5\alpha.$$

If we subtitute  $\alpha=0, x=1$ , for example, we get

$$g(\alpha x) = g(0 \cdot 1) = 3 \cdot 0 + 5 = 5,$$

but on the other hand

$$\alpha \cdot g(x) = 3 \cdot 1 \cdot 0 + 5 \cdot 0 = 0 \neq 5.$$

4

#### **Example**

Is the real function g(x) = 3x + 5 linear? Let's check:

### 1. Scalability:

$$g(\alpha x) = 3\alpha x + 5, \quad \alpha g(x) = 3x\alpha + 5\alpha.$$

If we subtitute  $\alpha=0, x=1$ , for example, we get

$$g(\alpha x) = g(0 \cdot 1) = 3 \cdot 0 + 5 = 5,$$

but on the other hand

$$\alpha \cdot g(x) = 3 \cdot 1 \cdot 0 + 5 \cdot 0 = 0 \neq 5.$$

Therefore, g is **NOT** linear.

4

#### Note

In order to prove that a function is linear, **both criteria** need to apply to **all** numbers  $\alpha, x$ , and y.

#### Note

In order to prove that a function is linear, **both criteria** need to apply to **all** numbers  $\alpha, x$ , and y.

In order to show that a function is **not** linear, it is enough to show that **just a single case** doesn't comply with **any of the criteria**.

5

#### Note

In order to prove that a function is linear, **both criteria** need to apply to **all** numbers  $\alpha, x$ , and y.

In order to show that a function is **not** linear, it is enough to show that **just a single case** doesn't comply with **any of the criteria**.

#### Challenge

Check whether the function g from before complies with the 2nd criterion (additivity).

#### **Example**

Is the function  $h(x) = x^2$  linear? Let's check additivity first:

$$h(x+y) = (x+y)^2 = x^2 + 2xy + y^2, \quad h(x) + h(y) = x^2 + y^2.$$

#### **Example**

Is the function  $h(x) = x^2$  linear? Let's check additivity first:

$$h(x+y) = (x+y)^2 = x^2 + 2xy + y^2, \quad h(x) + h(y) = x^2 + y^2.$$

Thus, for x = 1, y = -2:

$$h(x + y) = h(1 - 2) = h(-1) = (-1)^2 = 1.$$

#### **Example**

Is the function  $h(x) = x^2$  linear? Let's check additivity first:

$$h(x+y) = (x+y)^2 = x^2 + 2xy + y^2, \quad h(x) + h(y) = x^2 + y^2.$$

Thus, for x = 1, y = -2:

$$h(x + y) = h(1 - 2) = h(-1) = (-1)^2 = 1.$$

On the other hand,

$$h(x) + h(y) = h(1) + h(-2) = 1^2 + (-2)^2$$
  
= 1 + 4 = 5 \neq 1.

#### **Example**

Is the function  $h(x) = x^2$  linear? Let's check additivity first:

$$h(x+y) = (x+y)^2 = x^2 + 2xy + y^2, \quad h(x) + h(y) = x^2 + y^2.$$

Thus, for x = 1, y = -2:

$$h(x + y) = h(1 - 2) = h(-1) = (-1)^2 = 1.$$

On the other hand,

$$h(x) + h(y) = h(1) + h(-2) = 1^2 + (-2)^2$$
  
= 1 + 4 = 5 \neq 1.

Thus, h is also **NOT** linear.

### Challenge

Check whether h fulfills the 1st criterion (scalability).

#### Challenge

Check whether h fulfills the 1st criterion (scalability).

We can combine both criteria to a single test for linearity of a transformation  $T\colon$ 

#### **Definition**

A transformation  $T:A\to B$  is linear, if for all  $x,y\in A$  and  $\alpha,\beta\in\mathbb{R}$ 

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y).$$

7

Vectors can also be transformed, specifically by functions of the type  $T:\mathbb{R}^n\to\mathbb{R}^m$ , with  $n,m\in\mathbb{N}.$ 

Vectors can also be transformed, specifically by functions of the type  $T:\mathbb{R}^n\to\mathbb{R}^m$ , with  $n,m\in\mathbb{N}$ .

In this course we will mostly concentrate on transformations of the types

- $T: \mathbb{R}^2 \to \mathbb{R}^2$  and
- $T: \mathbb{R}^3 \to \mathbb{R}^3$ .

since they are more easy to conceptualize (and infintely easier to draw than higher dimensional transformations).

Vectors can also be transformed, specifically by functions of the type  $T:\mathbb{R}^n\to\mathbb{R}^m$ , with  $n,m\in\mathbb{N}$ .

In this course we will mostly concentrate on transformations of the types

- $T: \mathbb{R}^2 \to \mathbb{R}^2$  and
- $T: \mathbb{R}^3 \to \mathbb{R}^3$ .

since they are more easy to conceptualize (and infintely easier to draw than higher dimensional transformations).

However, everything we learn about these transformations is applicable for any linear transformation, **regardless of its dimensionality**.

#### **Example**


Applying the transformation  $T: \mathbb{R}^2 \to \mathbb{R}^2$ ,

$$T\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -x \\ 3y \end{pmatrix}$$

to the vector  $\vec{u} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ :


$$T\begin{pmatrix} 3\\1 \end{pmatrix} = \begin{pmatrix} -3\\3 \cdot 1 \end{pmatrix} = \begin{pmatrix} -3\\3 \end{pmatrix}.$$

9




### **Transforming Spaces**

We can visualize the way an entire space is transformed by a transformation  ${\cal T}$  by looking at how the axes and main gridlines of the space are transformed.



### **Transforming Spaces**

This method also allows us to see how the basis vectors  $\hat{x}$  and  $\hat{y}$  are transformed by linear transformations, and also the transformations of shapes (all this will come in handy later).



Some important properties of linear transformations are:

Some important properties of linear transformations are:

• The origin is preserved, i.e.

$$T\left(\vec{0}\right) = \vec{0}.$$

Some important properties of linear transformations are:

• The origin is preserved, i.e.

$$T\left(\vec{0}\right) = \vec{0}.$$

• Parallel lines remain parallel.

Some important properties of linear transformations are:

• The origin is preserved, i.e.

$$T\left(\vec{0}\right) = \vec{0}.$$

- Parallel lines remain parallel.
- All areas are scaled by the same number.

Some important properties of linear transformations are:

• The origin is preserved, i.e.

$$T\left(\vec{0}\right) = \vec{0}.$$

- Parallel lines remain parallel.
- All areas are scaled by the same number.


### Challenge

Show that these properties can be derived from the definition of linear transformations.

Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:


Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:

### Scaling in the x-axis




Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:

### Scaling in the y-axis




Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:

#### Rotation around the origin




Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:

#### Shear in the x-axis




Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:

#### Shear in the y-axis




Many linear transformations  $T: \mathbb{R}^2 \to \mathbb{R}^2$  can be created by composition of two or more of the following basic transformations:

Reflection by a line going through the origin




#### **Example**

The following transformation is a composition of a scaling transformation in the y-axis, followed by a rotation around the origin:



#### **Example**

The following transformation is a composition of a scaling transformation in the y-axis, followed by a rotation around the origin:



#### **Example**

The following transformation is a composition of a scaling transformation in the y-axis, followed by a rotation around the origin:

