

## Exercise 4: Determinants, Systems of Linear Equations

### Problem 1: Determinants

- Find the determinants of the following matrices:

(a)  $\begin{pmatrix} 1 & 3 \\ 2 & 0 \end{pmatrix}$

(b)  $\begin{pmatrix} 1 & 3 & -2 \\ 2 & 0 & 7 \\ 4 & 5 & 5 \end{pmatrix}$

(c)  $\begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}$

(d)  $\begin{pmatrix} 1 & 0 & 2 \\ 0 & 0 & 0 \\ 2 & 4 & 7 \end{pmatrix}$

(e)  $\begin{pmatrix} 4 & 2 & 2 \\ 2 & 1 & 1 \\ 2 & 0 & 7 \end{pmatrix}$

- What does it mean when the determinant of a  $2 \times 2$  matrix is negative?

### Problem 2: Row Operations and Rank

A matrix is said to be in its *Row Echelon Form* if:

- All nonzero rows (rows with at least one nonzero element) are above any rows of all zeroes (all zero rows, if any, belong at the bottom of the matrix)
- The leading coefficient (the first nonzero number from the left) of a nonzero row is always strictly to the right of the leading coefficient of the row above it.

If the leading coefficients are all 1, then the form is called a *Reduced Row Echelon Form*.

For example, the following matrices are all presented in reduced row echelon form:

$$\begin{pmatrix} 1 & -2 & 0 & 5 \\ 0 & 1 & 3 & 7 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 7 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 3 & 7 \\ 0 & 1 & 2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 5 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

- Use row operations to bring the following two matrices to their reduced row echelon form:

$$A = \begin{pmatrix} 3 & 1 & 0 \\ 6 & 2 & 1 \\ 3 & 6 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 1 \\ 2 & 0 & 2 \\ -3 & 0 & -3 \end{pmatrix}.$$

- What are the ranks of  $A$  and  $B$ ?

### Problem 3: Systems of Linear Equations

The following system of linear equations is given:

$$\begin{cases} -x + 3z = 20 \\ 3x + y + 3z = 15 \\ 9x + 3y = -18 \end{cases}.$$

Solve the system using the Gaussian elimination method.