

Implementing Molecular Hydrophobicity Potential Measurement for the Analysis of Dynamic Biomolecular Interactions

Peleg Bar Sapir¹

Under supervision of Prof. Maria Andrea Mroginski²

¹Freie Universität Berlin

²Technische Universität Berlin

February 15, 2018

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

- General form

- Force constants

- Distance function

Surface

- Solvent accesible surface

- Evenly distributed points

- Integration

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

- General form

- Force constants

- Distance function


Surface

- Solvent accesible surface

- Evenly distributed points

- Integration

The MHP Formula



$$\text{MHP}(\mathbf{x}') = \sum_{i=1}^k \left[f_i \cdot D(\mathbf{x} - \mathbf{x}'_i) \right]$$

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants

Distance function


Surface

Solvent accessible surface

Evenly distributed points

Integration

The MHP Formula



$$\text{MHP}(\mathbf{x}') = \sum_{i=1}^k \left[f_i \cdot D(\mathbf{x} - \mathbf{x}'_i) \right]$$

Summing over all atoms

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants

Distance function

Surface

Solvent accessible surface

Evenly distributed points

Integration

The MHP Formula

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants


Distance function

Surface

Solvent accessible surface

Evenly distributed points

Integration



$$\text{MHP}(\mathbf{x}') = \sum_{i=1}^k \left[f_i \cdot D(\mathbf{x} - \mathbf{x}'_i) \right]$$

Summing over all atoms

Force constants

The MHP Formula

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants


Distance function

Surface

Solvent accessible surface

Evenly distributed points

Integration



$$\text{MHP}(\mathbf{x}') = \sum_{i=1}^k \left[f_i \cdot D(\mathbf{x} - \mathbf{x}'_i) \right]$$

Summing over all atoms

Force constants

Distance function

Force constants

Type	Description	f_i value
<u>C in:</u>		
3	CHR_3	-0.6681
15	$=\text{CH}_2$	-0.7866
36	$\text{R}-\text{CH}-\text{X}$	-0.2405
<u>H attached to:</u>		
45	C_{sp^3} , no X attached to next carbon	0.7341
46	C_{sp^3} , C_{sp^2}	0.6301
50	Heteroatom	-0.1036
52	C_{sp^3} , 1 X attached to next carbon	0.6666
<u>O in:</u>		
56	Alcohol	-0.3567
58	Ketone	-0.0233
62	O^-	-0.7941

Source: Arup K. Ghose et al, J. Phys. Chem. A 1998, 102, 3762-3772


Distance function

Audry form

$$D(x) = \frac{1}{1+x}$$

Exponential decay form

$$D(x) = e^{-\alpha x}$$



Solvent accessible surface

Molecular
Hydrophobicity
Potential

Pelg Bar Sapir

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants

Distance function

Surface


Solvent accessible surface

Evenly distributed points

Integration


Evenly distributed points

How to distribute N points on a surface of a sphere?



Evenly distributed points


How to distribute N points on a surface of a sphere?



$$\varphi_i = i \cdot \frac{2\pi}{N}$$
$$\theta_j = j \cdot \frac{\pi}{N}$$


Evenly distributed points

How to distribute N points on a surface of a sphere?



$$\varphi_i = i \cdot \frac{2\pi}{N}$$
$$\theta_j = j \cdot \frac{\pi}{N}$$

Molecular Hydrophobicity Potential





- ▶ Points are not evenly distributed

$$\begin{aligned}\varphi_i &= i \cdot \frac{2\pi}{N} \\ \theta_j &= j \cdot \frac{\pi}{N}\end{aligned}$$

$$\theta_j = j \cdot \frac{\pi}{N}$$

Evenly distributed points

How to distribute N points on a surface of a sphere?



$$\varphi_i = i \cdot \frac{2\pi}{N}$$
$$\theta_j = j \cdot \frac{\pi}{N}$$

- Points are not evenly distributed
- Several points overlap at poles

Evenly distributed points

Solution: **Vogel's method**

In 2 dimensions:

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants

Distance function

Surface

Solvent accessible surface

Evenly distributed points

Integration

Evenly distributed points

Solution: **Vogel's method**

In 2 dimensions:


- ▶ Distances: $r_i = \sqrt{\frac{i}{N}}$
- ▶ Angle: $\theta_i = \varphi i$
(φ is the golden ratio!)

Evenly distributed points

Solution: **Vogel's method**

In 2 dimensions:

- ▶ Distances: $r_i = \sqrt{\frac{i}{N}}$
- ▶ Angle: $\theta_i = \varphi i$
(φ is the golden ratio!)




In 3 dimensions (cylindrical coordinates):

Evenly distributed points

Solution: **Vogel's method**

In 2 dimensions:

- ▶ Distances: $r_i = \sqrt{\frac{i}{N}}$
- ▶ Angle: $\theta_i = \varphi i$
(φ is the golden ratio!)



In 3 dimensions (cylindrical coordinates):


- ▶ Distances: $z_i = \left(1 - \frac{1}{N}\right) \left(1 - \frac{2i}{N-1}\right)$
- ▶ Angles:
 $\theta_i = \varphi i, \rho_i = \sqrt{1 - z_i^2}$

Evenly distributed points

Solution: **Vogel's method**

In 2 dimensions:

- ▶ Distances: $r_i = \sqrt{\frac{i}{N}}$
- ▶ Angle: $\theta_i = \varphi i$
(φ is the golden ratio!)



In 3 dimensions (cylindrical coordinates):

- ▶ Distances: $z_i = \left(1 - \frac{1}{N}\right) \left(1 - \frac{2i}{N-1}\right)$
- ▶ Angles:
 $\theta_i = \varphi i$, $\rho_i = \sqrt{1 - z_i^2}$




Image source: Marmakoide's Blog

Integration

Molecular Hydrophobicity Potential

Pelg Bar Sapir

Introduction

Hydrophobicity and log P

Molecular Hydrophobicity Potential

Potential

General form

Force constants

Distance function

Surface

Solvent accessible surface

Evenly distributed points

Integration

