Kinases as Drug Targets: A History of Failures and Successes

Supervisor: Dr Gian De Nicola

Word count: 2,999

Contents

Abstract 4

1. Introduction 4

2. Discussion 5

2.1. Imatinib success in Chronic Myeloid Leukaemia 5

2.2. The Initial Failure and Ultimate Success of Gefitinib in Non-Small Cell Lung Cancer 7

2.3 The Systematic Failure of p38 MAPK Inhibitors in Rheumatoid Arthritis 8

2.4 The Universal Challenge of Acquired Resistance and Emerging Strategies 12

3. Conclusion 14

References 15

List of Figures

Figure 1. JAK vs p38 MAPK Pathway Positioning 10

Figure 2. Structural basis of kinase inhibitor resistance and strategies to overcome them 13

List of Tables

Table 1. Comparative summary of key first-generation kinase inhibitors 11

Abstract

Protein kinases are central regulators of cellular signalling and represent a major class of therapeutic drug targets. In the history of kinase inhibitor development, most of the field's significant advances have been driven by an understanding of clinical failures. The initial success of Imatinib in chronic myeloid leukaemia (CML) established a powerful but simple paradigm of targeting a single, dominant oncogenic driver. This "best-case scenario" contrasts with two subsequent, more complex case studies. The first is the initial failure and ultimate success of the Epidermal Growth Factor Receptor (EGFR) inhibitor Gefitinib in non-small cell lung cancer (NSCLC), which emphasised the importance of biomarker-driven patient stratification. The second is the systematic failure of at least 14 p38 mitogen-activated protein kinase (MAPK) inhibitors in rheumatoid arthritis (RA), which revealed the profound challenge of targeting a single downstream node in a redundant signalling network. As such, the aim of this dissertation is to argue that a deep understanding of disease heterogeneity, genetic context, and signalling network architecture are the key determinants of success or failure in kinase-targeted therapy.

1. Introduction

Protein kinases are a family of enzymes that catalyse the transfer of a phosphate group from ATP to a substrate protein, a process known as phosphorylation. Within the cell, kinases operate as nodes in signalling networks where they receive signals from cell surface receptors and transmit these signals downstream through phosphorylation cascades. This post-translational modification acts as a molecular switch, regulating a vast array of cellular processes including proliferation, differentiation, metabolism, and apoptosis (1). The human genome encodes over 500 protein kinases, collectively referred to as the human kinome, which form the backbone of the cell's signalling architecture (1). Given their central role in cellular regulation, it is unsurprising that abnormal kinase activity is a hallmark of many human diseases, especially cancer. As such, the discovery that specific kinases can act as oncogenic drivers has made them highly attractive therapeutic targets (1).

The appeal of kinase inhibitors as therapeutic agents lies in their potential for high selectivity, oral bioavailability, and a more favourable toxicity profile compared to conventional cytotoxic

chemotherapy (1). Importantly, these principles are not limited to oncology; they are also a driving force for kinase inhibitor development in chronic inflammatory and autoimmune diseases like rheumatoid arthritis, where the need for long-term, well-tolerated oral therapies is particularly important (2). Most clinically approved kinase inhibitors function by competing with ATP for binding at the kinase active site, and subtle structural differences between kinases can be exploited to achieve target selectivity (1). Since the approval of Imatinib in 2001, over 70 kinase-targeted agents have now reached clinical use across oncology, immunology, and other therapeutic areas (1), reflecting both the therapeutic promise and the broader pharmaceutical confidence in molecularly targeted therapy.

However, the history of kinase drug discovery is not a simple story of linear progress. For every clinical success, there have been equally significant failures, and the most important advances have often been born from analysis of clinical setbacks. Despite the large number of approved inhibitors, fundamental questions remain about why some kinase-targeted strategies succeed while others fail, even when the preclinical rationale appears equally strong. Understanding these principles is essential for guiding future drug design and avoiding the repetition of costly clinical failures. Therefore, this dissertation examines three case studies that together illustrate how disease biology, patient selection, and network complexity determine the success or failure of a kinase-targeted strategy, arguing that drug potency alone is insufficient, and that disease heterogeneity, genetic context, and signalling network architecture are the key determinants of therapeutic outcome.

2. Discussion

2.1. Imatinib success in Chronic Myeloid Leukaemia

The story of Imatinib is the foundational success of rational drug design in oncology. Chronic Myeloid Leukaemia is a myeloproliferative neoplasm characterised by the overproduction of granulocytes. The disease is driven by a specific genetic abnormality: the Philadelphia chromosome, a shortened chromosome 22 that results from a reciprocal translocation between chromosomes 9 and 22, t(9;22)(q34;q11) (3). This translocation creates the BCR-ABL fusion gene, which encodes a constitutively active, non-receptor tyrosine kinase. The BCR-ABL protein

drives uncontrolled proliferation and survival of myeloid cells through downstream signalling cascades including RAS/MAPK and PI3K/AKT (1). This direct, causal link between a single genetic mutation and the disease phenotype makes BCR-ABL an ideal therapeutic target.

Mechanistically, Imatinib functions as a potent and selective ATP-competitive inhibitor of the ABL kinase domain. Crucially, Imatinib binds to and stabilises the inactive (closed) conformation of the ABL kinase. In this conformation, the activation loop is folded in a way that blocks substrate access and prevents autophosphorylation (3). By locking the kinase in this inactive state, Imatinib prevents ATP from binding, thereby shutting down the downstream signalling responsible for the cancerous phenotype (1).

The clinical outcomes were revolutionary. Before Imatinib, the prognosis for CML was grim, with the disease historically transforming to fatal blast crisis within an average of 4–6 years (3). Its introduction in 2001 transformed CML into a manageable chronic condition. The landmark International Randomized Study of Interferon and STI571 (IRIS) trial demonstrated that in newly diagnosed chronic phase CML patients, Imatinib induced a complete haematologic response (CHR) in 98% and a complete cytogenetic response (CCyR) in 87% at five years, with an estimated overall survival rate of 89% (4). A single-institution historical comparison found that the 8-year survival rate jumped from 42–65% in the pre-Imatinib era (1983–2000) to 87% after 2001 (5).

The profound success of Imatinib can be attributed to the near-perfect alignment of disease biology and drug mechanism. CML is a classic example of oncogene addiction, a state where a cancer cell is mostly dependent on a single activated oncogene for its survival (1). The entire pathophysiology is driven by BCR-ABL, making it a highly vulnerable, single therapeutic target. Another important part of the Imatinib story is that acquired resistance emerges in a significant proportion of patients, particularly those in advanced disease, establishing a cycle of resistance and next-generation drug development that is explored further in Section 2.4. As such, Imatinib's success established a powerful but deceptively simple paradigm: identify a single driving kinase and inhibit it. As the later case studies demonstrate, this paradigm does not generalise to most solid tumours or complex inflammatory diseases, where pathway redundancy, heterogeneous drivers, and adaptive evolution limit durable responses.

2.2. The Initial Failure and Ultimate Success of Gefitinib in Non-Small Cell Lung Cancer

In contrast to CML, the story of Gefitinib in Non-Small Cell Lung Cancer (NSCLC) is a crucial lesson in patient stratification and biomarker discovery. NSCLC is a highly heterogeneous disease comprising multiple distinct molecular subtypes, which was not fully appreciated in the early 2000s (6). The Epidermal Growth Factor Receptor (EGFR), a receptor tyrosine kinase, was known to be overexpressed in many NSCLC tumours. Following the Imatinib paradigm, the initial assumption was that targeting this overexpressed kinase would be broadly effective, treating EGFR overexpression as equivalent to BCR-ABL in CML. Two years after Imatinib's approval, in May 2003, Gefitinib (Iressa) received accelerated approval from the US FDA for advanced NSCLC after failure of chemotherapy (6). However, the pivotal discovery of activating EGFR mutations that explained drug sensitivity did not occur until 2004 (7, 8). Critically, Gefitinib was not designed to target a specific EGFR mutation; it was developed as a general EGFR inhibitor before mutation-based stratification existed. This timeline explains why early trials were conducted in unselected populations: the predictive biomarker for sensitivity was not yet known.

The initial trial outcomes were considerably disappointing. In unselected populations, Gefitinib demonstrated only a modest ~10% response rate (6). The Phase III ISEL trial failed to show a statistically significant survival benefit for Gefitinib over placebo (9). This led to restriction of Gefitinib to existing patients in 2005 and its eventual withdrawal from the US market (6, 9).

The turning point came in 2004 when two independent research groups discovered that the tumours of the small subset of responding patients contained specific activating mutations within the EGFR kinase domain, primarily small in-frame deletions in exon 19 and a point mutation in exon 21 (L858R) (7, 8). These responders were disproportionately of East Asian ethnicity, female, non-smokers, and had adenocarcinoma histology. Adenocarcinoma is the most common histological subtype of NSCLC, arising from the glandular epithelial cells of the distal airways, and is distinct from squamous cell carcinoma. EGFR mutations are strongly enriched in adenocarcinoma, particularly in never-smokers and East Asian patients (6).

This discovery led to a re-evaluation of Gefitinib's mechanism. The activating mutations destabilise the inactive conformation of the kinase, shifting the equilibrium towards the active state. This simultaneously increases kinase sensitivity to Gefitinib and reduces the effective affinity for ATP, creating a therapeutic window that does not exist in wild-type EGFR (7, 8). For the non-responders, the issue was twofold. Firstly, in tumours with wild-type EGFR, the drug cannot inhibit the kinase sufficiently at tolerable doses because the inhibitor must compete against normal ATP concentrations without an altered binding equilibrium. Crucially, this does not reflect poor inhibitor design; Gefitinib potently inhibits EGFR. The problem is that the therapeutic window only exists where the activating mutation has altered the kinase's biochemistry, making the distinction between an ineffective drug and an incorrectly applied one central to this case study. Secondly, and more fundamentally, many of these tumours are not dependent on EGFR signalling for survival; their growth is driven by other oncogenic pathways (such as KRAS mutations), making EGFR a bystander rather than a driver.

The core lesson of the Gefitinib story was not a failure of the drug, but a failure of patient selection and disease understanding. It proved that overexpression of a kinase is not a reliable proxy for dependency; a specific, activating genetic driver is the key. Gefitinib was eventually re-approved specifically for patients with EGFR-mutation-positive NSCLC, completing the transformation from failure to precision success (6).

2.3 The Systematic Failure of p38 MAPK Inhibitors in Rheumatoid Arthritis

If the Gefitinib story was a lesson in patient selection, p38 MAPK inhibitors in rheumatoid arthritis (RA) are a lesson in the complexities of signalling networks. RA is a chronic autoimmune inflammatory disease where immune cells attack the joints, producing pro-inflammatory cytokines like tumour necrosis factor-α (TNF-α) and interleukin-1 (IL-1), which drive joint destruction (10). The p38 MAPK pathway is a key intracellular cascade that regulates production of these cytokines, making it a logical therapeutic target. The strong preclinical rationale rested on clear evidence: p38 directly regulates TNF-α mRNA stability and translation, and TNF-α itself was already a clinically validated target in RA through biological therapies. Specifically, infliximab, a chimeric monoclonal antibody that directly neutralises circulating TNF-

α protein, had demonstrated significant clinical efficacy, establishing that reducing TNF-α activity could control disease progression (10). Because TNF-α blockade at the protein level was proven effective, the logic of inhibiting the intracellular kinase that controls TNF-α production appeared compelling.

Following this rationale, the pharmaceutical industry invested heavily in ATP-competitive inhibitors of p38α. However, despite extensive clinical testing, no compound delivered durable benefit. In multiple trials, an initial drop in C-reactive protein (CRP) was transient, rebounding within weeks despite continued dosing. CRP is synthesised by hepatocytes primarily in response to interleukin-6 (IL-6), whose production itself is regulated by p38 MAPK; CRP therefore serves as an indirect pharmacodynamic biomarker of p38 pathway activity (11). For example, VX-702 produced a temporary CRP reduction but no significant improvement in the ACR20 (American College of Rheumatology 20% response criteria) clinical endpoint compared to placebo (11). This divergence between a transient biomarker response and lack of clinical efficacy pointed towards rapid compensatory mechanisms within the signalling network.

The mechanistic explanation is multi-faceted. Firstly, unlike CML, RA is a complex, polygenic disease with no single genetic driver. Secondly, the preclinical animal models did not accurately predict drug effects in humans. The collagen-induced arthritis (CIA) model in mice is an induced, relatively acute inflammatory syndrome that only partially captures the chronicity and clinical heterogeneity of established human RA. Furthermore, many rodent studies dosed prophylactically or at exposures difficult to sustain in humans because of dose-limiting hepatic and central nervous system toxicities described across the class, so efficacy in rodents overestimated what was clinically achievable (10). Despite these limitations, the compelling biology of p38-mediated TNF-α regulation led the field to proceed with clinical development.

Most critically, the failure resulted from the inherent redundancy of the inflammatory signalling network. The p38 pathway is just one of several parallel pathways, including JNK, ERK, and NF-κB, that drive cytokine production. When p38 was blocked, the network re-routed through these alternatives, compensating for the blockade and restoring cytokine production (10). Furthermore, p38 is a downstream node in the cascade, contrasting sharply with the success of targeting upstream drivers like BCR-ABL and mutant EGFR.




Figure 1. JAK vs p38 MAPK Pathway Positioning. Created using BioRender. (A) Upstream targeting: Tofacitinib blocks JAK1/JAK3 at the receptor-proximal level, suppressing STAT-dependent inflammatory gene transcription. (B) Downstream targeting: VX-702 blocks p38α, but parallel ERK, NF-κB, and JNK cascades compensate, maintaining cytokine output. The terms 'upstream' and 'downstream' refer to proximity to the initiating cytokine signal, not a hierarchical relationship between these parallel cascades.

This systematic failure led to abandonment of the p38 class for RA and a strategic shift to Janus kinase (JAK) inhibitors. JAK kinases are positioned upstream, directly associating with cytokine receptors at the cell surface. By blocking signalling at this receptor-proximal level, JAK inhibitors suppress the entire downstream cascade rather than a single node within it, making compensatory rerouting far more difficult. Tofacitinib demonstrated a ~60% ACR20 response rate compared to ~27% for placebo, supporting this upstream approach (2). The p38 saga demonstrates that a strong preclinical rationale and a potent drug are insufficient for clinical success when faced with a redundant signalling network. The original logic of targeting p38 as a downstream effector was that it would be more selective and better tolerated than upstream

blockade, but the biology of RA proved this reasoning wrong. The failure was not one of chemistry, but of biological strategy, teaching the field that understanding signalling network topology is as important as understanding the target itself.

Table 1. Comparative summary of key first-generation kinase inhibitors discussed in this dissertation. *CHR is Complete Haematologic Response (normalisation of peripheral blood counts and leukocyte differential), CCyR is Complete Cytogenetic Response (0% Philadelphia chromosome-positive metaphases), OS is Overall Survival, ACR20 is American College of Rheumatology 20% improvement criteria (≥20% improvement in tender and swollen joint counts and ≥20% improvement in at least three of five measures: patient pain assessment, patient and physician global assessments, disability index, and acute phase reactant) and CRP is C-Reactive Protein.*

Drug	Molecular Formula	Target Kinase	Inhibitor Type	Key Clinical Outcome
Imatinib	$C_{29}H_{31}N_7O$	BCR-ABL	ATP-competitive (Type II)	98% CHR; 87% CCyR; 89% 5-yr OS (4)
Gefitinib	$C_{22}H_{24}ClFN_4O_3$	EGFR (mutant)	ATP-competitive (reversible)	~10% unselected; ~70% in EGFR-mutant (6,7)
VX-702	$C_{19}H_{12}F_4N_4O_2$	p38α MAPK	ATP-competitive	CRP decrease then rebound; ACR20 not significantly improved vs placebo (11)
Tofacitinib	$C_{16}H_{20}N_6O$	JAK1/JAK3	ATP-competitive	~60% ACR20 vs ~27% placebo (2)

2.4 The Universal Challenge of Acquired Resistance and Emerging Strategies

Even when initial therapy succeeds, the long-term efficacy of kinase inhibitors is almost universally challenged by the emergence of acquired resistance. The potent selective pressure exerted by a targeted drug eliminates sensitive cells, allowing the outgrowth of pre-existing or newly mutated resistant clones (12). A primary mechanism of this resistance is the development of secondary mutations at the “gatekeeper” residue, which sits within the ATP-binding pocket and controls access to a hydrophobic sub-pocket (13). This residue is named because it acts as a molecular gatekeeper regulating access to the deeper regions of the binding cleft, and mutations at this position can fundamentally alter the drug-kinase interaction. Importantly, different gatekeeper mutations confer resistance through distinct mechanisms. The T790M mutation in EGFR, first identified clinically in patients who had relapsed on Gefitinib (12), confers resistance primarily by increasing the kinase’s affinity for ATP by nearly 18-fold, thereby outcompeting reversible inhibitors rather than simply blocking drug binding sterically (13). By contrast, the T315I mutation in BCR-ABL, first reported by Gorre et al. (14), confers resistance through both steric clash with Imatinib and loss of a critical hydrogen bond at the gatekeeper position (3, 14). It is important to note that gatekeeper mutations, while clinically prominent, are not the only resistance mechanism: gene amplification, activation of bypass signalling pathways (such as MET amplification in EGFR-mutant NSCLC), and histological transformation also contribute to treatment failure (1).

The discovery of these resistance mutations has driven an “arms race” of next-generation inhibitor design. In CML, the T315I mutation conferred resistance to both first-generation (Imatinib) and second-generation (Dasatinib, Nilotinib) inhibitors because all share an ATP-competitive mechanism vulnerable to the same steric obstruction at the gatekeeper position. Ponatinib was specifically engineered to overcome this, using a carbon–carbon triple bond linker that accommodates the bulky isoleucine residue without steric clash (15). In NSCLC, Osimertinib, a third-generation covalent EGFR inhibitor, forms an irreversible bond with cysteine 797 (C797) near the ATP-binding pocket, sustaining kinase inhibition despite the T790M-mediated increase in ATP affinity. The AURA3 trial demonstrated its efficacy specifically in T790M-positive patients who had progressed on earlier EGFR inhibitors (16), and the FLAURA

trial subsequently established Osimertinib as first-line treatment for EGFR-mutant NSCLC due to its superior progression-free survival and central nervous system penetration (17).

Beyond sequential ATP-competitive inhibitors, an alternative strategy is allosteric inhibition. GNF-2 was identified as an allosteric inhibitor proposed to bind the myristoyl pocket on the C-lobe of ABL (18). Structural studies subsequently confirmed this binding site using NMR and X-ray crystallography and demonstrated that combining allosteric and ATP-site inhibitors can suppress even the T315I-mutant kinase in preclinical models (19). This concept was translated clinically as Asciminib, an allosteric BCR-ABL1 inhibitor targeting the same regulatory pocket, which showed activity in heavily pre-treated CML patients, including those with the T315I mutation (20). Its distinct mechanism of action also opens the possibility of combining allosteric and ATP-site inhibitors to raise the genetic barrier to resistance (19, 20).




Figure 2. Structural basis of kinase inhibitor resistance and strategies to overcome them. Created using BioRender. (A) Normal drug binding at the ATP-binding pocket, showing the gatekeeper threonine residue forming a hydrogen bond with the drug. (B) Gatekeeper mutations disrupt drug binding; the boxes distinguish two principal mechanisms: ABL T315I (steric clash and lost hydrogen bond) versus EGFR T790M (increased ATP affinity reducing

inhibitor potency) (13). (C) Strategies to overcome resistance: allosteric inhibition (GNF-2/Asciminib binding the myristoyl pocket on the C-lobe, stabilising the autoinhibited conformation) and covalent inhibition (Osimertinib forming an irreversible bond with Cys797).

3. Conclusion

The history of kinase inhibitor development illustrates how progress in drug discovery is driven as much by failure as by success. Imatinib provided the foundational proof-of-concept for targeted therapy, but its simplicity was misleading, creating the expectation of a "magic bullet" that was not broadly applicable. Gefitinib demonstrated that understanding the specific genetic context of a tumour is paramount, establishing biomarker-driven patient selection as the standard of care. The p38 story highlighted that targeting a single, downstream node in a redundant signalling network is a fundamentally flawed strategy, regardless of preclinical rationale or drug potency.

These case studies reveal that the success of a kinase inhibitor depends not on potency alone, but on the interaction between the drug, the genetic makeup of the disease, and the signalling network in which the target is embedded. Effective kinase-targeted therapy requires a systems-level understanding integrating the genomic context of the patient (the Gefitinib lesson), the topology of the relevant signalling network (the p38 lesson), and the evolutionary dynamics of resistance (the Imatinib lesson). However, the emergence of tertiary resistance mutations such as C797S in Osimertinib-treated patients suggests that sequential monotherapy with ATP-site inhibitors may be inherently self-limiting. This raises the possibility that the future of the field may lie less in ever-more-selective single agents and more in rational upfront combinations of mechanistically orthogonal inhibitors, such as allosteric and ATP-competitive agents, that suppress resistance before it emerges. Whether such strategies can achieve durable responses without unacceptable toxicity remains a critical open question and addressing it will require the convergence of precision biology, structural pharmacology, and network-level thinking that represents the future of kinase drug development.

References

[1] Bhullar KS, Lagarón NO, McGowan EM, Parmar I, Jha A, Hubbard BP, et al. Kinase-targeted cancer therapies: progress, challenges and future directions. Molecular Cancer. 2018 Feb 19;17(1):48.

[2] Fleischmann R, Kremer J, Cush J, Schulze-Koops H, Connell CA, Bradley JD, et al. Placebo-controlled trial of tofacitinib monotherapy in rheumatoid arthritis. The New England Journal of Medicine. 2012 Aug 9;367(6):495-507.

[3] Druker BJ. Translation of the Philadelphia chromosome into therapy for CML. Blood. 2008 Dec 15;112(13):4808-4817.

[4] Druker BJ, Guilhot F, O'Brien SG, Gathmann I, Kantarjian H, Gattermann N, et al. Five-year follow-up of patients receiving imatinib for chronic myeloid leukemia. The New England Journal of Medicine. 2006 Dec 7;355(23):2408-2417.

[5] Kantarjian H, O'Brien S, Jabbour E, Garcia-Manero G, Quintas-Cardama A, Shan J, et al. Improved survival in chronic myeloid leukemia since the introduction of imatinib therapy: a single-institution historical experience. Blood. 2012 Jan 6;119(9):1981-1987.

[6] Sharma SV, Bell DW, Settleman J, Haber DA. Epidermal growth factor receptor mutations in lung cancer. Nature Reviews Cancer. 2007 Mar;7(3):169-181.

[7] Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. The New England Journal of Medicine. 2004 May 20;350(21):2129-2139.

[8] Paez JG, Jänne PA, Lee JC, Tracy S, Greulich H, Gabriel S, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. Science. 2004 Jun 4;304(5676):1497-1500.

[9] Thatcher N, Chang A, Parikh P, Rodrigues Pereira J, Ciuleanu T, von Pawel J, et al. Gefitinib plus best supportive care in previously treated patients with refractory advanced non-small-cell

lung cancer: results from a randomised, placebo-controlled, multicentre study (Iressa Survival Evaluation in Lung Cancer). The Lancet. 2005 Oct;366(9496):1527-1537.

[10] Hammaker D, Firestein GS. “Go upstream, young man”: lessons learned from the p38 saga. Annals of the Rheumatic Diseases. 2009 Dec 7;69(Suppl 1):i77-i82.

[11] Damjanov N, Kauffman RS, Spencer-Green GT. Efficacy, pharmacodynamics, and safety of VX-702, a novel p38 MAPK inhibitor, in rheumatoid arthritis: results of two randomized, double-blind, placebo-controlled clinical studies. Arthritis & Rheumatism. 2009 May;60(5):1232-1241.

[12] Pao W, Miller VA, Politi KA, Riely GJ, Somwar R, Zakowski MF, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the EGFR kinase domain. PLoS Medicine. 2005 Feb 22;2(3):e73.

[13] Yun CH, Mengwasser KE, Toms AV, Woo MS, Greulich H, Wong KK, et al. The T790M mutation in EGFR kinase causes drug resistance by increasing the affinity for ATP. Proceedings of the National Academy of Sciences of the United States of America. 2008 Jan 28;105(6):2070-2075.

[14] Gorre ME, Mohammed M, Ellwood K, Hsu N, Paquette R, Rao PN, et al. Clinical resistance to STI-571 cancer therapy caused by BCR-ABL gene mutation or amplification. Science. 2001 Jun 21;293(5531):876-880.

[15] O'Hare T, Shakespeare WC, Zhu X, Eide CA, Rivera VM, Wang F, et al. AP24534, a pan-BCR-ABL inhibitor for chronic myeloid leukemia, potently inhibits the T315I mutant and overcomes mutation-based resistance. Cancer Cell. 2009 Nov;16(5):401-412.

[16] Mok TS, Wu YL, Ahn MJ, Garassino MC, Kim HR, Ramalingam SS, et al. Osimertinib or platinum-pemetrexed in EGFR T790M-positive lung cancer. The New England Journal of Medicine. 2017 Feb 16;376(7):629-640.

[17] Soria JC, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. The New England Journal of Medicine. 2018 Jan 11;378(2):113-125.

[18] Adrián FJ, Ding Q, Sim T, Velentza A, Sloan C, Liu Y, et al. Allosteric inhibitors of Bcr-Abl-dependent cell proliferation. Nature Chemical Biology. 2006 Jan 15;2(2):95-102.

[19] Zhang J, Adrián FJ, Jahnke W, Cowan-Jacob SW, Li AG, Iacob RE, et al. Targeting Bcr-Abl by combining allosteric with ATP-binding-site inhibitors. Nature. 2010 Jan;463(7280):501-506.

[20] Hughes TP, Mauro MJ, Cortes JE, Minami H, Rea D, DeAngelo DJ, et al. Asciminib in chronic myeloid leukemia after ABL kinase inhibitor failure. The New England Journal of Medicine. 2019 Dec 12;381(24):2315-2326.