


LECTURE 11: Derived distributions

- Given the distribution of X ,
find the distribution of $Y = g(X)$
 - the discrete case
 - the continuous case
 - general approach, using CDFs
 - the linear case: $Y = aX + b$
 - general formula when g is monotonic
- Given the (joint) distribution of X and Y ,
find the distribution of $Z = g(X, Y)$

Derived distributions — the discrete case

$$Y = g(X)$$





$$p_Y(4) = \mathbb{P}(Y = 4)$$

$$= \mathbb{P}(X = 4) + \mathbb{P}(X = 5)$$


$$= p_X(4) + p_X(5) = 0.3 + 0.4$$

$$\begin{aligned} p_Y(y) &= \mathbb{P}(g(X) = y) \\ &= \sum_{x: g(x) = y} p_X(x) \end{aligned}$$

A linear function of a discrete r.v.






$$Y = 2X + 3 = Z + 3$$



$$\begin{aligned} P_Y(y) &= P(Y=y) = P(2X+3=y) \\ &= P\left(X = \frac{y-3}{2}\right) = p_X\left(\frac{y-3}{2}\right) \end{aligned}$$

$$Y = aX + b : \quad p_Y(y) = p_X\left(\frac{y-b}{a}\right)$$

A linear function of a continuous r.v.



A linear function of a continuous r.v.

$$Y = aX + b$$

$$a > 0$$

$$P(Y = \gamma) = P(aX + b = \gamma) = P\left(X = \frac{\gamma - b}{a}\right)$$

$$\begin{aligned} F_Y(\gamma) &= P(Y \leq \gamma) = P(aX + b \leq \gamma) \\ &= P\left(X \leq \frac{\gamma - b}{a}\right) = F_X\left(\frac{\gamma - b}{a}\right) \end{aligned}$$

$$f_Y(\gamma) = f_X\left(\frac{\gamma - b}{a}\right) \cdot \frac{1}{a}$$

$$f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y - b}{a}\right)$$

$$a < 0$$

$$\begin{aligned} &= P\left(X \geq \frac{\gamma - b}{a}\right) \\ &= 1 - P\left(X \leq \frac{\gamma - b}{a}\right) \\ &= 1 - F_X\left(\frac{\gamma - b}{a}\right) \end{aligned}$$

$$f_Y(\gamma) = -f_X\left(\frac{\gamma - b}{a}\right) \cdot \frac{1}{a}$$

$$p_Y(y) = p_X\left(\frac{y - b}{a}\right) \cdot \frac{1}{|a|}$$

A linear function of a normal r.v. is normal

$$X \sim \mathcal{N}(\mu, \sigma^2)$$

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$$

$$Y = aX + b, \quad a \neq 0$$

$$f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$$

$$\begin{aligned} f_Y(y) &= \frac{1}{|a|} \frac{1}{\sqrt{2\pi}\sigma} e^{-\left(\frac{y-b}{a} - \mu\right)^2/2\sigma^2} \\ &= \frac{1}{\sqrt{2\pi}\sigma|a|} e^{-\frac{(y-b-a\mu)^2}{2\sigma^2 a^2}} \end{aligned}$$



If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $aX + b \sim \mathcal{N}(a\mu + b, a^2\sigma^2)$

A general function $g(X)$ of a continuous r.v.

- Two-step procedure:

- Find the CDF of Y : $F_Y(y) = \mathbf{P}(Y \leq y) = \mathbf{P}(g(x) \leq y)$
- Differentiate: $f_Y(y) = \frac{dF_Y}{dy}(y)$

Example: $Y = X^3$; X uniform on $[0, 2]$





$$0 \leq y \leq 8$$

$$F_Y(y) = P(Y \leq y) = P(X^3 \leq y) = P(X \leq y^{1/3}) = \frac{1}{2} y^{1/3}$$

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{1}{2} \cdot \frac{1}{3} y^{-2/3} = \frac{1}{6} \cdot \frac{1}{y^{2/3}}$$

Example: $Y = a/X$

- You go to the gym and set the speed X of the treadmill to a number between 5 and 10 km/hr (with a uniform distribution). Find the PDF of the time it takes to run 10km.



$$\text{time} = Y = \frac{10}{X} \quad 1 \leq Y \leq 2$$

$$F_Y(Y) = P(Y \leq Y) = P\left(\frac{10}{X} \leq Y\right)$$

$$= P\left(X \geq \frac{10}{Y}\right) = \frac{1}{5} \left(10 - \frac{10}{Y}\right)$$

$$f_Y(Y) = \frac{1}{5} \frac{(-10)}{-Y^2} = \frac{2}{Y^2} \quad , 1 \leq Y \leq 2$$
$$= 0 \quad , \text{otherwise}$$

A general formula for the PDF of $Y = g(X)$ when g is monotonic $x^3 \frac{a}{x}$

decreasing $x < x' \Rightarrow g(x) > g(x')$

Assume g strictly increasing


and differentiable

$$F_Y(y) = P(Y \leq y) = P(X \leq h(y)) = F_X(h(y))$$

$$f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$$

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(X \geq h(y)) \\ &= 1 - P(X \leq h(y)) = 1 - F_X(h(y)) \end{aligned}$$

$$f_Y(y) = \textcolor{green}{1} f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$$




inverse function $h \rightarrow$ decreasing

$$f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$$

Example: $Y = X^2$; X uniform on $[0, 1]$

$$f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$$




$$y = x^2 \Leftrightarrow x = \sqrt{y} \quad h(y) = \sqrt{y}$$

$$f_Y(y) = \frac{1}{2\sqrt{y}}$$

$$0 \leq y \leq 1$$

An intuitive explanation for the monotonic case



$$y = g(x) \quad \delta_2 \approx \delta_1 \frac{dg}{dx}(x)$$

$$x = h(y) \quad \delta_1 \approx \delta_2 \cdot \frac{dh}{dy}(y) \quad \textcircled{*}$$


$$f_Y(y) \delta_2 \approx \mathbb{P}(y \leq Y \leq y + \delta_2) = \mathbb{P}(x \leq X \leq x + \delta_1)$$

$$\approx f_X(x) \delta_1 \approx f_X(x) \delta_2 \frac{dh}{dy}(y)$$

$$f_Y(y) = f_X(x) \frac{dh}{dy}(y)$$

$$= f_X(h(y)) \frac{dh}{dy}(y)$$

A nonmonotonic example: $Y = X^2$



- The discrete case:

$$p_Y(9) = P(X=3) + P(X=-3)$$

$$p_Y(y) = P_X(\sqrt{y}) + P_X(-\sqrt{y})$$

- The continuous case: $y \geq 0$

$$F_Y(y) = P(Y \leq y) = P(X^2 \leq y) = P(|X| \leq \sqrt{y}) = P(-\sqrt{y} \leq X \leq \sqrt{y})$$



$$= F_X(\sqrt{y}) - F_X(-\sqrt{y})$$

$$f_Y(y) = f_X(\sqrt{y}) \frac{1}{2\sqrt{y}} + f_X(-\sqrt{y}) \frac{\cancel{(-1)}}{2\sqrt{y}}$$

A function of multiple r.v.'s: $Z = g(X, Y)$

$$Y = g(x)$$

- Same methodology: find CDF of Z
- Let $Z = Y/X$; X, Y independent, uniform on $[0, 1]$



$$\begin{aligned} F_Z(z) &= P\left(\frac{Y}{X} \leq z\right) = 0, \quad z < 0 \\ &= \frac{1}{2} \cdot z, \quad 0 \leq z \leq 1 \\ &= 1 - \frac{1}{2z}, \quad z > 1 \end{aligned}$$

