

## Week 13 - Quiz

### Problem 1.

The correlation coefficient is used to determine:

- A. A specific value of the y-variable given a specific value of the x-variable
- B. A specific value of the x-variable given a specific value of the y-variable
- C. The strength of the linear relationship between the x and y variables
- D. The relationship between the x and y variables

**Answer – C.** It measures the linear relationship.

### Problem 2.

If there is a very strong correlation between two variables, then the correlation coefficient must be

- A. any value larger than 1
- B. much smaller than 0, if the correlation is negative
- C. much larger than 0, regardless of whether the correlation is negative or positive
- D. either close to 1 or -1.

**Answer - D:** It is close to 1 if the two variables have a very strong positive correlation; it is close to -1 if the two variables have a very strong negative correlation.

### Problem 3.

In regression, the equation that describes how the response variable (y) is related to the explanatory variable (x) is:

- A. the correlation model
- B. the regression model
- C. used to compute the correlation coefficient
- D. a two-sample test

**Answer - B:**

### Problem 4.

In regression analysis, the variable that is being predicted is the

- A. response, or dependent, variable
- B. independent variable
- C. intervening variable
- D. is usually x

**Answer – A.**

**Problem 5.**

The coefficient of correlation

- A. is the square of the coefficient of determination
- B. is the square root of the coefficient of determination
- C. is the same as r-square
- D. can never be negative.

**Answer- B.**

**Problem 6.**

If the coefficient of determination is a positive value, then the regression equation

- A. must have a positive slope
- B. must have a negative slope
- C. could have either a positive or a negative slope
- D. must have a positive y-intercept

**Answer-C:** because we don't know the sign of the slope.

**Problem 7.**

If the coefficient of determination is equal to 1, then the correlation coefficient

- A. must also be equal to 1
- B. can be either -1 or +1
- C. can be any value between -1 to +1
- D. must be -1

**Answer-B.** The square of the correlation coefficient is equal to the coefficient of determination.

**Problem 8**

If the correlation coefficient is 0.8, the percentage of variation in the response variable explained by the variation in the explanatory variable, i.e., the coefficient of determination, is

- A. 0.80%
- B. 80%
- C. 0.64%
- D. 64%

**Answer- D.** The coefficient of determination represents the percentage of variation explained by the explanatory variable.

**Problem 9**

If the correlation coefficient is a positive value, then the slope of the regression line

- A. must also be positive
- B. can be either negative or positive
- C. can be zero
- D. cannot be zero

**Answer- A.**

**Problem 10**

If the coefficient of determination is 0.81, the correlation coefficient

- A. is 0.6561
- B. could be either + 0.9 or - 0.9
- C. must be positive
- D. must be negative

**Answer-B.** The square of the correlation coefficient is equal to the coefficient of determination.

**Problem 11**

A variety of summary statistics were collected for a small sample (10) of bivariate data, where the dependent variable was y and the independent variable was x

$$\begin{aligned}\Sigma(Y - \bar{Y})(X - \bar{X}) &= 466 \\ \Sigma(X - \bar{X})^2 &= 234 \\ \Sigma(Y - \bar{Y})^2 &= 1434\end{aligned}$$

Using the following coefficient of correlation,

$$r = \frac{\sum_{i=1}^n ((x_i - \bar{x})(y_i - \bar{y}))}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2 \sum_{i=1}^n (y_i - \bar{y})^2}}$$

we have r =

- A. 0.8045
- B. -0.8045
- C. 0
- D. 1

**Answer-A.**

**Problem 12**

The data above is from a research study. Ice cream consumption was measured over 30 four-week periods. One purpose of the study was to determine whether ice cream consumption depended on the price of ice cream. The researcher uses least-squares linear regression to determine this prediction equation:

$$y = -2.05x + 0.9230.$$

Which statement best interprets the meaning of the **slope** of the prediction equation?

- A. For a \$1 increase in the price of a pint of ice cream, we can estimate a per capita ice cream consumption increase of approximately \$2.05.
- B. For a \$1 increase in the price of a pint of ice cream, we can estimate a per capita ice cream consumption decrease of approximately 2.05 pints.
- C. For ice cream priced at \$1 per pint, we can estimate that the per capita ice cream consumption will decrease by 0.9230 pints.
- D. For 1 pint of ice cream consumed per capita, we can estimate its price will be approximately \$2.05.

**Answer- B.**

## Summary of Weekly Assignment #13

### 1. Five Number Summary :

The five-number summary is used to describe the shape of the distribution of a given numerical data. It consists of five numbers: minimum data value, first quartile, median, the third quartile, and the maximum data value.

The five-number summary of this given data set is:

stats	value
Min.	35.00
1st Qu.	90.00
Median	95.00
3rd Qu.	100.00
Max.	100.00

### 2. Boxplot :

The boxplot is a geometric representation of the five-number summary. The boxplot of the given data set is given below.

