

Answers to Chapter One HW

STA 321 Topics in Advanced Statistics

2023-02-12

Contents

Chapter 1: Section 1.1	1
Problem 1. Sec 1.1 1(a)	1
Problem 2. Sec 1.1 4(b)	2
Problem 3 Sec 1.1 11(a)	3
Problem 4 Sec 1.1 19	3
Problem 5 Sec 1.1 21	4
Chapter 1: Section 1.2	4
Problem 1 Sec 1.2: 1(a)	4
Problem 2 Sec 1.2: 19	5
Problem 3 Sec 1.2: 21	6
Chapter 1: Section 1.3	6
Problem 1 Sec 1.3: 6(a)	6
Problem 2 Sec 1.3: 7(a)	7
Problem 3 Sec 1.3: 11	7
Problem 3 Sec 1.3: 13(a)	8

Chapter 1: Section 1.1

Problem 1. Sec 1.1 1(a)

Textbook Section 1.1.

1(a). Show that the following equations have at least one solution in the given intervals.

$$x \cos x - 2x^2 + 3x - 1 = 0, [0.2, 0.3] \text{ and } [1.2, 1.3]$$

Solution: We use the mean value theorem to show the existence of the roots on the given interval. Let $f(x) = x \cos x - 2x^2 + 3x - 1$.

define the following function to evaluate the end points of the intervals.

```
fun = function(x) x*cos(x) - 2*x^2 + 3*x - 1
```

calling the above function

```
fun(0.2)*fun(0.3)
```

```
## [1] -0.001874581
```

```
fun(1.2)*fun(1.3)
```

```
## [1] -0.02047641
```

Since $f(0.2) \times f(0.3) < 0$ and $f(1.2) \times f(1.3) < 0$, equation $f(x) = 0$ has at least one root in each of the two intervals.

Problem 2. Sec 1.1 4(b)

Textbook Section 1.1.

4(b). Find $\max_{a \leq x \leq b} |f(x)|$ for the following functions and intervals.

b. $f(x) = (4x - 3)/(x^2 - 2x)$ on $[0.5, 1]$

Solution: We first find the extreme values in the following.

$$f'(x) = \frac{4(x^2 - 2x) - (4x - 3)(2x - 2)}{(x^2 - 2x)^2} = \frac{-4x^2 + 6x - 6}{x^4 - 4x^3 + 4x^2}.$$


Since $-4x^2 + 6x - 6 = 0$ has no real root. We only need to evaluate the function on the end points of the given interval to determine the maximum of $|f(x)|$. We do in the in the following R code

```
# define the R function
fun1.1.4b = function(x) (4*x - 3)/(x^2 - 2*x)
# Evaluating the function
c(fun1.1.4b(0.5), fun1.1.4b(1))
```

```
## [1] 1.333333 -1.000000
```

Next we plot the function on interval $[0.5, 1]$.

```
xx = seq(0.5, 1, length = 200)
yy = fun1.1.4b(xx)
plot(xx, yy, xlab = "", ylab = "", type = "l", lwd = 2, col = "blue", main = "", bty = "n")
```



Therefore $\max_{a \leq x \leq b} |f(x)| = 4/3$.

Problem 3 Sec 1.1 11(a)

Textbook Section 1.1.

11(a): Find the third Taylor polynomial $P_3(x)$ for the function $f(x) = (x - 1) \ln x$ about $x_0 = 1$.

- a. Use $P_3(0.5)$ to approximate $f(0.5)$. Find an upper bound for error $|f(0.5) - P_3(0.5)|$ using the error formula, and compare it to the actual error.

Solution:

$$f(x) = (x - 1)^2 - \frac{1}{2}(x - 1)^3 + \frac{\frac{2}{\xi^3} + \frac{6}{\xi^4}}{3!}(x - 1)^4$$

Where

$$P_3(x) = (x - 1)^2 - \frac{1}{2}(x - 1)^3 \quad \text{and} \quad R_3(x) = \frac{\frac{2}{\xi^3} + \frac{6}{\xi^4}}{4!}(x - 1)^4.$$

Since $\xi \in [0.5, 1]$,

$$|R_3(x)| = \frac{\frac{2}{\xi^3} + \frac{6}{\xi^4}}{4!}(x - 1)^4 \leq \frac{\frac{2}{0.5^3} + \frac{6}{0.5^4}}{4!}(0.5 - 1)^4 = 0.2916667.$$

The approximated value and the true values are

```
# Define f(x) and Pn(x)
true.fun = function(x) (x-1)*log(x)
Pn = function(x) (x-1)^2 - 0.5*(x-1)^3
##
true.val = true.fun(0.5)
est.val = Pn(0.5)
est.error = abs(true.val - est.val)
##
kable(c(true.val = true.val, est.val = est.val, est.error = est.error))
```

	x
true.val	0.3465736
est.val	0.3125000
est.error	0.0340736

The true, estimated values and the estimated errors are given in the above table.

Problem 4 Sec 1.1 19

Let $f(x) = e^x$ and $x_0 = 0$. Find the nth Taylor polynomial $P_n(x)$ for $f(x)$ about x_0 . Find a value of n necessary for $P_n(x)$ to approximate $f(x)$ to within 10^{-6} on $[0, 0.5]$.

Solution. The Taylor expansion of $f(x) = e^x$ has the following form

$$e^x = \sum_{k=0}^n \frac{x^k}{k!} + \frac{e^\xi}{(n+1)!}x^{n+1}$$

where $\xi \in (0, x)$. The last term is the remainder which indicates the error when we approximate e^x by $P_n(x) = \sum_{k=0}^n \frac{x^k}{k!}$. However, the remainder error on interval $[0, 0.5]$ can be approximated by

$$R_n(x) = \frac{e^\xi}{(n+1)!}x^{n+1} \leq \frac{e^{0.5}}{(n+1)!}0.5^{n+1}$$

To find n , we solve the following inequality

$$\frac{e^{0.5}}{(n+1)!} 0.5^{n+1} < 10^{-10}$$

The algebraic solution to the above inequality is infeasible. We use numerical method to find the solution.

```
# define the remainder function
Rn = function(x){exp(0.5)*0.5^(x+1) / factorial(x+1)}
# choose a sequence of x to find the smallest value that
# meets the error condition.
x = 1:10
Rn(x)

## [1] 2.060902e-01 3.434836e-02 4.293545e-03 4.293545e-04 3.577954e-05
## [6] 2.555682e-06 1.597301e-07 8.873894e-09 4.436947e-10 2.016794e-11
```

From the above output, the minimum degree of the estimated polynomial is $n = 7$.

Problem 5 Sec 1.1 21

21. The polynomial $P_2(x) = 1 - x^2/2$ is to be used to approximate $f(x) = \cos x$ in $[-1/2, 1/2]$. Find a bound for the maximum error.

Solution: Note that the Taylor expansion of $f(x) = \cos(x)$ has the following general form.

$$\cos(x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!}$$

The remainder error term can be written as

$$R_n(x) = \frac{(-1)^k \cos(\xi) x^{2k}}{(2k)!} \quad \text{where } \xi \in [0, x]$$

Since $\cos(x) \leq 1$. Therefore,

$$|R_n(x)| = \left| \frac{(-1)^k \cos(\xi) x^{2k}}{(2k)!} \right| \leq \frac{x^{2k}}{(2k)!}.$$

When $P_n(x) = 1 - x^2/2$ is used to approximate $f(x) = \cos(x)$, the degree of $P_n(x)$ is $n = 2k = 2$. Therefore, for $x \in [-0.5, 0.5]$, the error bound is given by

$$R_2(x) \leq \frac{0.5^{2 \times 2}}{(2 \times 2)!} = 0.002604167.$$

Chapter 1: Section 1.2

Problem 1 Sec 1.2: 1(a)

1(a): Compute the absolute error and relative error in approximations of p by p^* .

a. $p = \pi, p^* = 22/7$

Solution: We use the following R code to do the calculation.

```
abs.error = abs(pi - 22/7)
rel.error = abs(pi - 22/7)/pi
kable(c(abs.error = abs.error, rel.error = rel.error))
```

	x
abs.error	0.0012645
rel.error	0.0004025

Problem 2 Sec 1.2: 19

19: The two-by-two linear system

$$ax + by = e,$$

$$cx + dy = f,$$

where a, b, c, d, e, f are given, can be solved for x and y as follows:

set

$$m = c/a, \text{ provided } a \neq 0;$$

$$d_1 = d - mb;$$

$$f_1 = f - me;$$

$$y = f_1/d_1;$$

$$x = (e - by)/a.$$

Solve the following linear systems using four-digit rounding arithmetic.

(a). The system is given by

$$1.130x - 6.990y = 14.20$$

$$1.013x - 6.099y = 14.22$$

Solution: The expression of the exact solution to the linear system is given. The error in the solution is due to the rounding error. We use the following R code to do the calculation.

```
# coefficients
a = 1.130
b = -6.990
c = 1.013
d = -6.099
e = 14.20
f = 14.22
####
m = c/a
d1 = d - m*b
f1 = f - m*e
```

```

y = round(f1/d1,4)           # 4-digit arithmetic requirement
x = round((e - b*y)/a,4)
c(x = x, y = y)

##           x           y
## 67.6828  8.9101

```

Problem 3 Sec 1.2: 21

21. (a) Show that the polynomial nesting technique can also be applied to the evaluation of

$$f(x) = 1.01e^{4x} - 4.62e^{3x} - 3.11e^{2x} + 12.2e^x - 1.99.$$

(b). Use three-digit rounding arithmetic, the assumption that $e^{1.53} = 4.62$, and the fact that $e^{nx} = (e^x)^n$ to evaluate $f(1.53)$ as given in part (a).

(c). Redo the calculation in part (b) by first nesting the calculations.

(d). Compare the approximations in parts (b) and (c) to the true three-digit result $f(1.53) = -7.61$.

Solution: (a). Note that the given $f(x)$ can be re-expressed in the following nested structure.

$$f(x) = (((1.01 \times e^x - 4.62)e^x - 3.11)e^x + 12.2)e^x - 1.99$$

Therefore, we can use nested arithmetic to evaluate the above polynomial.

(b)-(d). This means the direct evaluation by term using 3-digit arithmetic (rounding each term by keeping 3 decimal places). We do the calculation using R in the following. The results are summarized in the output.

```

x = 1.53
E = 4.62
True.val = 1.01*exp(1.53)^4 - 4.62*exp(1.53)^3 - 3.11*exp(1.53)^2 + 12.2*exp(1.53) - 1.99
PartB.val = 1.01*E^4 - 4.62*E^3 - 3.11*E^2 + 12.2*E - 1.99
PartC.val = round((round((round((round(1.01*E,3) - 4.62)*E,3) - 3.11)*E,3) + 12.2)*E,3) - 1.99
##
PartD.Rel.ErrB = abs((True.val - PartB.val)/True.val)
PartD.Rel.ErrC = abs((True.val - PartC.val)/True.val)
c(True.val = True.val, PartB.val = PartB.val, PartC.val = PartC.val, PartD.Rel.ErrB = PartD.Rel.ErrB, PartD.Rel.ErrC = PartD.Rel.ErrC)

##           True.val           PartB.val           PartC.val PartD.Rel.ErrB PartD.Rel.ErrC
##      -7.60787145      -7.45124989      -7.46000000      0.02058678      0.01943664

```

The absolute relative error of the nest algorithm is smaller than that of the regular approximation in (b).

Chapter 1: Section 1.3

Problem 1 Sec 1.3: 6(a)

6(a): Find the rates of convergence of the following sequences as $n \rightarrow \infty$.

$$\lim_{n \rightarrow \infty} \sin(1/n) = 0$$

Solution: We need to use an important result in Calculus: $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$. Using this result, we have

$$\lim_{n \rightarrow \infty} \frac{\sin(1/n)}{1/n} = 1.$$

By definition, $\sin(1/n) = O(1/n)$. That is, the convergence rate of $\sin(1/n)$ as $n \rightarrow \infty$ is the same as that of $\{1/n\}$.

Problem 2 Sec 1.3: 7(a)

7(a). Find the rates of convergence of the following functions as $h \rightarrow 0$. **a.**

$$\lim_{h \rightarrow 0} \frac{\sin(h)}{h} = 1$$

Solution: We first use the Taylor expansion

$$\frac{\sin(h)}{h} = 1 - h^2/6 + h^4/120 - h^6/5040 + h^8/362880 - \dots = 1 + O(h^2)$$

That is the convergence rate of given limit is the same as h^2 , $O(h^2)$.

Problem 3 Sec 1.3: 11

11: Construct an algorithm that has as input an integer $n \geq 1$, numbers x_0, x_1, \dots, x_n , and a number x and that produces as output the product $(x - x_0)(x - x_1) \dots (x - x_n)$.

Solution: We first write the pseudo code for the algorithm.

```

INPUT: vector[x1, x2, ..., xn]
      n
      x
PUTPUT: cumulative product

STEP 1: P = 1      (initial cumulative product)

STEP 2: FOR i < n DO:
      Pi = x - xi
      P = P*Pi
      OUTPUT (P = result)

```

The next example of implementing the algorithm is optional.

Example: Assume that the given vector $V = c(1, 3, 2, 5, 7, 5, 9, 4, 2, 0, 7)$. We use n and x as input variables (arguments). The following R function implements the above algorithm.

```

cumProduct = function(n, x, V){
  if(n > length(V)) stop("n should be smaller than the dimension of the vector!")
  P = 1
  for(i in 1:n){
    P = P*(x-V[i])
  }
  return(P)
}

```

```
V = c(1,3,2,5,7,5,9,4,2,0,7)
##
cumProduct( n=8,
            x = 5.5,
            V = V)
```

```
## [1] 77.51953
```

Problem 3 Sec 1.3: 13(a)

a. Suppose that $0 < q < p$ and that $\alpha_n = \alpha + O(n^{-p})$. Show that $\alpha_n = \alpha + O(n^{-q})$.

Solution: Using the definition of **big O** notation and the given condition $\alpha_n = \alpha + O(n^{-p})$, we have

$$|\alpha_n - \alpha| \leq Kn^{-p} < Kn^{-q} \quad \text{since } 0 < q < p.$$

Therefore,

$$\frac{\alpha_n - \alpha}{n^{-q}} \leq K \quad \text{as } n \rightarrow \infty.$$

This implies that the rate of convergence is $O(n^{-q})$.