

## STA 504 Homework #8

Due: Tuesday, November 7

Answer the following questions based on the given joint density function in problems 1 - 4.

- (i) Find the marginal densities  $f(x)$  and  $f(y)$ .
- (ii) Check if the variables  $X$  and  $Y$  are independent.
- (iii) find the conditional probability distribution function of  $X$ , given  $Y = y$


### Problem 1.

Let the continuous random vector  $(X, Y)$  have joint probability density function

$$f(x, y) = e^{-y}, 0 < x < y < \infty.$$

In addition to the (i)-(iii), compute the  $P(X \geq 2; Y \geq 4)$ .

**Solution:** We first sketch the domain of the joint density function in the left panel (a) of the following figure. It is both type I and II region.



(i). Marginal density functions.

$$f_X(x) = \int_x^{\infty} e^{-y} dy = -e^{-y} \Big|_x^{\infty} = e^{-x}$$

$$f_Y(y) = \int_0^y e^{-y} dx = e^{-y} x \Big|_0^y = ye^{-y}$$

The above two densities represent exponential and gamma distributions respectively.

(ii) Since  $f_X(x)f_Y(y) = ye^{-x-y} \neq e^{-y} = f(x, y)$ , random variable X and Y are dependent on each other.

$$(iii). f(x|y) = \frac{f(x,y)}{f(y)} = \frac{e^{-y}}{ye^{-y}} = \frac{1}{y}, \quad x < y < \infty$$

(iv).  $P(X > 2, Y > 4)$  is the volume between the sky-blue region and the surface of the density function. However, the sky-blue region is irregular. We can partition it into two regular regions: one type I and one type II (see (b) of the above figure). Therefore,

$$\begin{aligned} P(X > 2, Y > 4) &= \iint_{R_1} f(x, y) dA + \iint_{R_2} f(x, y) dA \\ &= \int_2^4 \int_4^\infty e^{-y} dy dx + \int_4^\infty \int_4^y e^{-y} dx dy = \int_2^4 e^{-4} dx + \int_4^\infty (y-4)e^{-y} dy \\ &= (4-2)e^{-4} + \int_4^\infty (y-4)e^{-y} dy = 2e^{-4} + \int_4^\infty (y-4)e^{-y} dy \end{aligned}$$

Note that

$$\begin{aligned} \int_4^\infty (y-4)e^{-y} dy &= -\int_4^\infty (y-4)e^{-y} d(-y) = -\int_4^\infty (y-4)de^{-y} \\ &= -\{(y-4)e^{-y}|_4^\infty - \int_4^\infty e^{-y} d(y-4)\} \\ &= -\left\{0 + \int_4^\infty e^{-y} d(-y)\right\} = -e^{-y}|_4^\infty = -(0 - e^{-4}) = e^{-4}. \end{aligned}$$

$$\text{Therefore, } P(X > 2, Y > 4) = 2e^{-4} + \int_4^\infty (y-4)e^{-y} dy = 2e^{-4} + e^{-4} = 3e^{-4}$$


## Problem 2.

Let X and Y be a random vector with joint probability density function

$$f(x, y) = \begin{cases} 6xy(2 - x - y), & \text{if } 0 < x < 1, 0 < y < 1 \\ 0, & \text{otherwise} \end{cases}$$

Then find the conditional probability distribution function of X, given  $Y = y$ , where  $0 < y < 1$ .

**Solution.** We first sketch the domain in the following.



(i). Marginal density functions.

$$\begin{aligned} f_X(x) &= \int_0^1 6xy(2 - x - y) dy = \int_0^1 (12xy - 6x^2y - 6xy^2) dy \\ &= (6xy^2 - 3x^2y^2 - 2xy^3) \Big|_{y=0}^{y=1} = 6x - 3x^2 - 2x = 4x - 3x^2 \end{aligned}$$

$$\begin{aligned} f_Y(y) &= \int_0^1 6xy(2 - x - y) dx = \int_0^1 (12xy - 6x^2y - 6xy^2) dx \\ &= (6x^2y - 2x^3y - 3x^2y^2) \Big|_{x=0}^{x=1} = 6y - 3y^2 - 2y = 4y - 3y^2 \end{aligned}$$

(ii) Since  $f_X(x)f_Y(y) = (6x - 3x^2 - 2x)(6y - 3y^2 - 2y) \neq 6xy(2 - x - y) = f(x, y)$ , random variable X and Y are dependent on each other.


$$(iii). \quad f(x|y) = \frac{f(x,y)}{f(y)} = \frac{6xy(2-x-y)}{4y-3y^2} = \frac{6x(2-x-y)}{4-3y}, \quad x < y < \infty$$

### Problem 3.

Let X and Y be a random vector with joint probability density function.

$$f(x, y) = \begin{cases} 1/x, & \text{if } 0 < y < x < 1 \\ 0, & \text{otherwise} \end{cases}$$

**Solution.** The domain of the density is sketched below.



(i). Marginal density functions.

$$f_X(x) = \int_0^x \frac{1}{x} dy = \frac{1}{x} \times y \Big|_{y=0}^{y=x} = 1$$

$$f_Y(y) = \int_y^1 \frac{1}{x} dx = \ln(x) \Big|_y^1 = -\ln(y)$$

(ii). Since  $f_X(x)f_Y(y) = 1(-\ln(y)) \neq 1/x = f(x, y)$ , random variable X and Y are dependent on each other.

$$(iii). f(x|y) = \frac{f(x,y)}{f(y)} = \frac{1/x}{-\ln(y)} = -\frac{1}{x \ln(y)}, \quad 0 < y < x < 1.$$


#### Problem 4.

Let X and Y have density

$$f(x, y) = \begin{cases} xe^{-x(1+y)}, & \text{if } y > 0, x > 0 \\ 0, & \text{otherwise} \end{cases}$$

In addition to the (i)-(iii), compute the  $P(X \geq 2; Y \geq 4)$ .

**Solution:** We sketch the domain in the following.



(i). Marginal density functions.

$$f_X(x) = \int_0^{\infty} x e^{-x(1+y)} dy = - \int_0^{\infty} x e^{-x(1+y)} d[-x(1+y)] = e^{-x(1+y)} \Big|_{y=0}^{y=\infty} = e^{-x}$$

$$\begin{aligned} f_Y(y) &= \int_0^{\infty} x e^{-x(1+y)} dx = - \frac{1}{(1+y)} \int_0^{\infty} x e^{-x(1+y)} d[-x(1+y)] \\ &= - \frac{1}{(1+y)} \int_0^{\infty} x d e^{-x(1+y)} = - \frac{1}{(1+y)} \left[ x e^{-x(1+y)} \Big|_{x=0}^{x=\infty} - \int_0^{\infty} e^{-x(1+y)} dx \right] \\ &= \frac{1}{(1+y)} \left[ - \frac{1}{(1+y)} \int_0^{\infty} e^{-x(1+y)} d(-x(1+y)) \right] = - \frac{1}{(1+y)^2} e^{-x(1+y)} \Big|_{x=0}^{x=\infty} = \frac{1}{(1+y)^2} \end{aligned}$$

(ii). Since  $f_X(x)f_Y(y) = e^{-x} \frac{1}{(1+y)^2} \neq x e^{-x(1+y)} = f(x, y)$ , random variable X and Y are dependent on each other.

$$(iii). f(x|y) = \frac{f(x,y)}{f(y)} = \frac{x e^{-x(1+y)}}{\frac{1}{(1+y)^2}} = x(1+y)^2 e^{-x(1+y)}, \quad 0 < y < x < 1.$$

(iv).  $P(X > 2, Y > 4)$  is the volume between the sky-blue region and the surface of the density function. However, the sky-blue region is irregular. We can partition it into two regular regions: one type I and one type II (see (b) of the above figure). Therefore,

$$\begin{aligned} P(X > 2, Y > 4) &= \iint_{R_1} f(x, y) dA = \int_2^{\infty} \int_4^{\infty} x e^{-x(1+y)} dy dx \\ &= \int_2^{\infty} \int_4^{\infty} e^{-x(1+y)} dx (1+y) dx = - \int_2^{\infty} e^{-x(1+y)} \Big|_{y=4}^{y=\infty} dx \\ &= \int_2^{\infty} e^{-5x} dx = - \frac{e^{-5x}}{5} \Big|_{x=2}^{x=\infty} = \frac{e^{-10}}{5} \end{aligned}$$