

Measurements on Three Hawk Species

Description

Students and faculty at Cornell College in Mount Vernon, Iowa, collected data over many years at the hawk blind at Lake MacBride near Iowa City, Iowa. The data set that we are analyzing here is a subset of the original data set, using only those species for which there were more than 10 observations. Data were collected on random samples of three different species of hawks: Red tailed, Sharp-shinned, and Cooper's hawks.

Format

A data frame with 908 observations on the following 19 variables.

Month: 8=September to 12=December

Day: Date in the month

Year: Year: 1992-2003

CaptureTime: Time of capture (HH:MM)

ReleaseTime: Time of release (HH:MM)

BandNumber: ID band code

Species: CH=Cooper's, RT=Red-tailed, SS=Sharp-Shinned

Age: A=Adult or I=Immature

Sex: F=Female or M=Male

Wing: Length (in mm) of primary wing feather from tip to wrist it attaches to

Weight: Body weight (in gm)

Culmen: Length (in mm) of the upper bill from the tip to where it bumps into the fleshy part of the bird

Hallux: Length (in mm) of the killing talon

Tail: Measurement (in mm) related to the length of the tail (invented at the MacBride Raptor Center)

StandardTail: Standard measurement of tail length (in mm)

Tarsus: Length of the basic foot bone (in mm)

WingPitFat: Amount of fat in the wing pit

KeelFat: Amount of fat on the breastbone (measured by feel)

Crop: Amount of material in the crop, coded from 1=full to 0=empty