

STAT0008 Lecture 4

The Cramér-Rao Lower Bound and the Exponential Family of Distributions

Dr. Aidan O'Keeffe

Department of Statistical Science
University College London

22nd October 2018

- ▶ Score function – linear form and its relationship to the Cramér-Rao lower bound
- ▶ Minimum variance bound estimators (MVBES)
- ▶ Minimum variance bound unbiased estimators (MVBUES)
- ▶ Fisher information and the Cramér-Rao lower bound for k -dimensional parameters
- ▶ The exponential family of distributions

Cramér-Rao Lower Bound

Recall, from Lecture 3, the Cramér-Rao lower bound...

If T is an unbiased estimator for $m(\theta)$, then the variance of T is such that

$$\text{Var}(T) \geq \frac{[m'(\theta)]^2}{\mathcal{I}(\theta)}.$$

Also, from Lecture 3, the inequality above was derived from the assumption that the absolute value of the Pearson correlation between T and the score function $U(\theta; \mathbf{X})$ is ≤ 1 .

$$\text{i.e. } |\text{Corr}(T, U(\theta; \mathbf{X}))| \leq 1.$$

Cramér-Rao Lower Bound

We obtain equality where

$$\text{Corr}(T, U(\theta; \mathbf{X})) = 1.$$

In other words, the Cramér-Rao lower bound will be attained where the Pearson correlation between T and $U(\theta; \mathbf{X})$ is equal to 1.

A Pearson correlation of 1 means that there is a linear, deterministic relationship between T and $U(\theta; \mathbf{X})$.

We can show that the variance of T attains the Cramér-Rao lower bound if and only if T and $U(\theta; \mathbf{X})$ are linearly related such that

$$U(\theta; \mathbf{X}) = A(\theta) (T(\mathbf{X}) - m(\theta))$$

where $A(\theta)$ is a non-random function of θ .

Score Function: Linear Form

We see that, where the Cramér-Rao lower bound is attained by $\text{Var}(T(\mathbf{X}))$, the score function $U(\theta; \mathbf{X})$ can be written

$$U(\theta; \mathbf{X}) = A(\theta) (T(\mathbf{X}) - m(\theta))$$

$U(\theta; \mathbf{X})$, the score function, contains \mathbf{X} and θ .

$A(\theta)$ is a non-random function of θ . i.e. it has θ as an argument but **not** \mathbf{X} .

$T(\mathbf{X})$ is a statistic that is a function of \mathbf{X} only and does not contain θ .

$m(\theta)$ is a function of θ only and does not contain \mathbf{X} .

Note that the linear form must be as above so that $\mathbb{E}(U(\theta; \mathbf{X})) = 0$ holds.

Score Function: Linear Form

Why is this functional form useful...?

Well... if we can write the score function in this way we can:

1. Obtain an estimator of $m(\theta)$ that attains the Cramér-Rao lower bound (i.e. has the smallest possible variance).
2. Identify the Cramér-Rao lower bound (and Fisher information).

Point 1. is obvious. To verify Point 2., let us go back to the linear form of $U(\theta; \mathbf{X})$.

Score Function: Linear Form

Set

$$U(\theta; \mathbf{X}) = A(\theta) (T(\mathbf{X}) - m(\theta))$$

Score Function: Linear Form

Score Function: Linear Form

Hence, we see that

$$A(\theta) = \frac{\mathcal{I}(\theta)}{m'(\theta)}.$$

So if we write the score function in the form

$$\begin{aligned} U(\theta; \mathbf{X}) &= A(\theta) (T(\mathbf{X}) - m(\theta)) \\ &= \frac{\mathcal{I}(\theta)}{m'(\theta)} (T(\mathbf{X}) - m(\theta)). \end{aligned}$$

then we can easily recover the Fisher information and Cramér-Rao lower bound.

MVBE and MVBUE

Suppose that T is an estimator of $m(\theta)$.

If $\text{Var}(T)$ attains the Cramér-Rao lower bound, then T is known as a **minimum variance bound estimator**, with acronym **MVBE**.

If T is **unbiased** for $m(\theta)$ **and** $\text{Var}(T)$ attains the Cramér-Rao lower bound, then T is known as a **minimum variance bound unbiased estimator**, with acronym **MVBUE**.

Given a likelihood function, $\mathcal{L}(\theta \mid \mathbf{X})$, we can use the following steps to attempt to identify MVBUEs.

1. Write the score function $U(\theta; \mathbf{X}) = \frac{\partial}{\partial \theta} \log \mathcal{L}(\theta | \mathbf{X})$ in the form

$$U(\theta; \mathbf{X}) = A(\theta) (T(\mathbf{X}) - m(\theta)).$$

2. Then $T(\mathbf{X})$ is unbiased for $m(\theta)$ and $\text{Var}(T(\mathbf{X}))$ attains the Cramér-Rao lower bound.
3. The Cramér-Rao lower bound $\left(\frac{[m'(\theta)]^2}{\mathcal{I}(\theta)} \right)$ can be easily identified using the equation

$$A(\theta) = \frac{\mathcal{I}(\theta)}{m'(\theta)} \implies \text{CRLB} = \frac{m'(\theta)}{A(\theta)}.$$

T unbiased for θ

Note that when $T(\mathbf{X})$ is unbiased for θ , we have $m(\theta) = \theta$ and $m'(\theta) = 1$ so

$$U(\theta; \mathbf{X}) = \mathcal{I}(\theta) (T(\mathbf{X}) - \theta) .$$

In this case, the Cramér-Rao lower bound is

$$\text{CRLB} = \frac{1}{\mathcal{I}(\theta)} .$$

MVBUE: Example 1

Suppose that X_1, \dots, X_n are a set of iid random variables where each $X_i \sim \text{Bernoulli}(\theta)$. Find a MVBUE for θ and determine the variance of this estimator.

MVBUE: Example 1

MVBUE: Example 2

Suppose that X_1, \dots, X_n are a set of iid random variables where each $X_i \sim \text{Poi}(\lambda)$. Find a MVBUE for λ and determine the variance of this estimator.

MVBUE: Example 2

k —dimensional Parameter

Suppose now that $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k)^\top$ is a parameter of dimension k (i.e. a $k \times 1$ vector).

We define a sample X_1, \dots, X_n such that each $X_i \sim \mathcal{D}(\boldsymbol{\theta})$, with corresponding likelihood function $\mathcal{L}(\boldsymbol{\theta} \mid \mathbf{X})$. The log-likelihood function is

$$\ell(\boldsymbol{\theta} \mid \mathbf{X}) = \log[\mathcal{L}(\boldsymbol{\theta} \mid \mathbf{X})].$$

The score function takes the form of a $k \times 1$ vector, $U(\boldsymbol{\theta}; \mathbf{X})$, where

$$U(\boldsymbol{\theta}; \mathbf{X}) = \begin{pmatrix} \frac{\partial \ell(\boldsymbol{\theta} \mid \mathbf{X})}{\partial \theta_1} \\ \vdots \\ \frac{\partial \ell(\boldsymbol{\theta} \mid \mathbf{X})}{\partial \theta_k} \end{pmatrix}$$

k -dimensional Parameter

The score function is such that

$$\mathbb{E}[U(\boldsymbol{\theta}; \mathbf{X})] = \begin{pmatrix} \mathbb{E}\left[\frac{\partial \ell(\boldsymbol{\theta}|\mathbf{X})}{\partial \theta_1}\right] \\ \vdots \\ \mathbb{E}\left[\frac{\partial \ell(\boldsymbol{\theta}|\mathbf{X})}{\partial \theta_k}\right] \end{pmatrix} = \mathbf{0}.$$

As in the one-dimensional parameter case, the Fisher information is computed as the variance of the score function.

However, since $\boldsymbol{\theta}$ is a k -dimensional vector, the variance of the score function is a $k \times k$ variance-covariance matrix, known as the **Fisher information matrix**.

k —dimensional Parameter

The Fisher information matrix is

$$\begin{aligned}\mathcal{I}(\boldsymbol{\theta}) &= \text{Var}(U(\boldsymbol{\theta}; \mathbf{X})) \\ &= \mathbb{E}[U(\boldsymbol{\theta}; \mathbf{X})U(\boldsymbol{\theta}; \mathbf{X})^\top] \\ &= \begin{pmatrix} \mathbb{E}\left[\left(\frac{\partial \ell}{\partial \theta_1}\right)^2\right] & \cdots & \mathbb{E}\left[\frac{\partial \ell}{\partial \theta_1} \frac{\partial \ell}{\partial \theta_k}\right] \\ \vdots & \ddots & \vdots \\ \mathbb{E}\left[\frac{\partial \ell}{\partial \theta_k} \frac{\partial \ell}{\partial \theta_1}\right] & \cdots & \mathbb{E}\left[\left(\frac{\partial \ell}{\partial \theta_k}\right)^2\right] \end{pmatrix}\end{aligned}$$

where, for brevity, $\ell = \ell(\boldsymbol{\theta} \mid \mathbf{X})$. We note that the $(r, s)^{\text{th}}$ element of $\mathcal{I}(\boldsymbol{\theta})$ is

$$[\mathcal{I}(\boldsymbol{\theta})]_{rs} = \mathbb{E}\left[\frac{\partial \ell(\boldsymbol{\theta} \mid \mathbf{X})}{\partial \theta_r} \frac{\partial \ell(\boldsymbol{\theta} \mid \mathbf{X})}{\partial \theta_s}\right].$$

Under certain regularity conditions, the following result holds

$$[\mathcal{I}(\boldsymbol{\theta})]_{rs} = \mathbb{E} \left[-\frac{\partial^2 \ell(\boldsymbol{\theta} \mid \mathbf{X})}{\partial \theta_r \partial \theta_s} \right].$$

If $T(\mathbf{X})$ is an estimator for the k —dimensional vector, $\boldsymbol{\theta}$, we can establish the Cramér-Rao lower bound for the variance-covariance matrix of $\text{Var}(T(\mathbf{X}))$.

Unlike in the one-dimensional case, we shall concentrate only on unbiased estimators of $\boldsymbol{\theta}$.

k -dimensional Parameter: Cramér-Rao lower bound

Suppose that $T(\mathbf{X})$ is an estimator (in the form of a $k \times 1$ random variable) of the k -dimensional parameter $\boldsymbol{\theta}$. Then the variance-covariance matrix of $T(\mathbf{X})$ is such that

$$\text{Var}(T(\mathbf{X})) \geq \mathcal{I}^{-1}(\boldsymbol{\theta}).$$

Here, the matrix inequality \geq is such that $A \geq B$ implies that $A - B$ is positive semi-definite for square matrices A and B of the same dimension.

k —dimensional Parameter: Cramér-Rao lower bound

Moreover, if $T(\mathbf{X})$ is such that $T_j(\mathbf{X})$ is an unbiased estimator for θ_j (for some $j \in \{1, \dots, k\}$) then the Cramér-Rao lower bound for the variance of $T_j(\mathbf{X})$ is

$$\text{Var}(T_j(\mathbf{X})) \geq [\mathcal{I}^{-1}(\boldsymbol{\theta})]_{jj}.$$

Here, $T_j(\mathbf{X})$ is the j^{th} component of $T(\mathbf{X}) = (T_1(\mathbf{X}), \dots, T_k(\mathbf{X}))^T$ and θ_j is the j^{th} component of $\boldsymbol{\theta}$.

$[\mathcal{I}^{-1}(\boldsymbol{\theta})]_{jj}$ is the $(j, j)^{\text{th}}$ element of the inverse of the Fisher information matrix.

The condition for equality is that $T_j(\mathbf{X})$ is a linear function of $U_1(\boldsymbol{\theta}; \mathbf{X}), \dots, U_k(\boldsymbol{\theta}; \mathbf{X})$ where $U_j(\boldsymbol{\theta}; \mathbf{X})$ denotes the j^{th} component of the $k \times 1$ score function $U(\boldsymbol{\theta}; \mathbf{X})$ ($j \in \{1, \dots, k\}$).

k -dimensional Parameter: Example

Suppose that X_1, \dots, X_n are iid $\mathcal{N}(\mu, \sigma^2)$ random variables.

- (a) Compute the Fisher information matrix for the parameters $(\mu, \sigma^2)^\top$.
- (b) Hence
 - (i) Determine the Cramér-Rao lower bound for the variance of unbiased estimators of μ
 - (ii) Determine the Cramér-Rao lower bound for the variance of unbiased estimators of σ^2

k -dimensional Parameter: Example

k -dimensional Parameter: Example

k -dimensional Parameter: Example

Exponential Family

We have seen that, if an estimator $T(\mathbf{X})$ attains the Cramér-Rao lower bound, then the score function is of the form

$$\begin{aligned}U(\theta; \mathbf{X}) &= A(\theta)(T(\mathbf{X}) - m(\theta)) \\&= A(\theta)T(\mathbf{X}) + B(\theta) \quad \text{where} \quad B(\theta) = -A(\theta)m(\theta).\end{aligned}$$

Then

$$\frac{\partial}{\partial \theta} \log \mathcal{L}(\theta \mid \mathbf{X}) = A(\theta)T(\mathbf{X}) + B(\theta)$$

and integrating the above with respect to θ , we obtain

$$\log \mathcal{L}(\theta \mid \mathbf{X}) = a(\theta)T(\mathbf{X}) + b(\theta) + c(\mathbf{X})$$

with

$$a(\theta) = \int_{\Theta} A(\theta) d\theta, \quad b(\theta) = \int_{\Theta} B(\theta) d\theta.$$

Exponential Family

Hence, the likelihood function is

$$\begin{aligned}\mathcal{L}(\theta \mid \mathbf{X}) &= \exp \{a(\theta)T(\mathbf{X}) + b(\theta) + c(\mathbf{X})\} \\ &= [\exp \{a(\theta)T(\mathbf{X}) + b(\theta)\}] [\exp \{c(\mathbf{X})\}]\end{aligned}$$

and, by the factorisation criterion, we see that $T(\mathbf{X})$ is sufficient for θ .

Recall that the likelihood function, $\mathcal{L}(\theta \mid \mathbf{X})$ is the joint probability density/mass function of \mathbf{X} where θ is a distributional parameter.

Thus, the joint density/mass $f(\mathbf{x}; \theta)$ is such that

$$f(\mathbf{x}; \theta) = \exp \{a(\theta)T(\mathbf{x}) + b(\theta) + c(\mathbf{x})\}.$$

Exponential Family

A family of probability distributions whose density/mass function has the form

$$f(x; \theta) = \exp \{a(\theta)T(x) + b(\theta) + c(x)\}$$

for θ in some parameter space Θ and $X \sim \mathcal{D}(\theta)$, is said to be an **exponential family** of distributions.

We note that the support of $f(x; \theta)$ (i.e. the values of x for which $f(x; \theta)$ is defined) must not depend on θ .

In addition, $c(x)$ is not arbitrary because the pdf/pmf, $f(x; \theta)$, must integrate/sum to 1.

Exponential Family

If $\mathbf{X} = \{X_1, \dots, X_n\}$ are independent random variables from the same univariate exponential family then the likelihood function is given by

$$\mathcal{L}(\theta \mid \mathbf{x}) = \exp \{a(\theta)T(\mathbf{x}) + nb(\theta) + c(\mathbf{x})\}$$

where

$$T(\mathbf{x}) = \sum_{i=1}^n T(x_i) \quad \text{and} \quad c(\mathbf{x}) = \sum_{i=1}^n c(x_i).$$

Notice that the score function of the above implies that $\frac{T(\mathbf{X})}{n}$ is a MVBUE for $-\frac{b'(\theta)}{a'(\theta)}$.

Exponential Family: k -dimensional parameter

Suppose now that θ is a k -dimensional parameter ($k > 1$) where

$$\theta = (\theta_1, \dots, \theta_k)^\top.$$

A family of probability distributions $\{f(x; \theta), \theta \in \Theta\}$ belongs to the exponential family of distributions if $f(x; \theta)$ has the form

$$f(x; \theta) = \exp \left\{ \sum_{r=1}^k a_r(\theta) T_r(x) + b(\theta) + c(x) \right\}.$$

As in the one-dimensional parameter case, the support of $f(x; \theta)$ must not depend on θ .

Exponential Family: k -dimensional parameter

If $\mathbf{X} = \{X_1, \dots, X_n\}$ are independent random variables from the an exponential family parameterised by $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k)^\top \in \Theta$ then the joint density of \mathbf{X} is given by

$$f(\mathbf{x}; \boldsymbol{\theta}) = \exp \left\{ \sum_{r=1}^k a_r(\boldsymbol{\theta}) T_r(\mathbf{x}) + nb(\boldsymbol{\theta}) + c(\mathbf{x}) \right\}$$

where

$$T_r(\mathbf{x}) = \sum_{i=1}^n T_r(x_i) \quad \text{and} \quad c(\mathbf{x}) = \sum_{i=1}^n c(x_i).$$

It follows from the factorisation criterion that

$$\{T_1(\mathbf{X}), \dots, T_k(\mathbf{X})\}$$

are jointly sufficient for $\boldsymbol{\theta}$.

Exponential Family: k -dimensional parameter

Here, where the number jointly sufficient statistics is equal to the dimension of $\boldsymbol{\theta}$, the jointly sufficient statistics $T_1(\mathbf{X}), \dots, T_k(\mathbf{X})$ may be expressed as functions of

$$U_j(\boldsymbol{\theta}; \mathbf{X}) = \frac{\partial \ell(\boldsymbol{\theta}; \mathbf{X})}{\partial \theta_j} \quad j = 1, \dots, k.$$

Hence, estimators that are linear functions of $T_1(\mathbf{X}), \dots, T_k(\mathbf{X})$ will have variances that attain the corresponding Cramér-Rao lower bound.

Exponential Family

At this point, we might ask '*why is an exponential family important?*'

In answer

- ▶ If a distribution is from an exponential family, then sufficient statistics for the distributional parameters are guaranteed to exist.
- ▶ In a Bayesian framework, if a likelihood function/joint density belongs to an exponential family, then there exists a **conjugate prior** distribution for the distributional parameters.

Conjugate Prior

A conjugate prior is a prior whereby, when combined with a given likelihood function, the posterior distribution belongs to the same family as the prior.

For example, if we have a likelihood function from a $\text{Bin}(\theta)$ distribution and we choose a Beta prior for θ then we will obtain a Beta distribution for the posterior of θ .

This is a situation where the prior is a conjugate prior.

Exponential Families

Many familiar classes of distributions belong to the exponential family of distributions. For example

- ▶ Binomial, Poisson;
- ▶ Exponential, gamma;
- ▶ Normal, Beta.

Exponential Family: Example 1

Suppose that $X \sim \text{Poi}(\theta)$. Show that X belongs to the exponential family of distributions and identify the functions $T(x)$, $a(\theta)$, $b(\theta)$ and $c(x)$.

Exponential Family: Example 2

Suppose that $X \sim \text{Beta}(\alpha, \beta)$. Show that X belongs to the exponential family of distributions and identify the functions $T(x)$, $a(\boldsymbol{\theta})$, $b(\boldsymbol{\theta})$ and $c(x)$, where $\boldsymbol{\theta} = (\alpha, \beta)^\top$. (Note: $T(x)$ and $a(\boldsymbol{\theta})$ have dimension 2)

Exponential Family: Example 2

Learning Outcomes

- ▶ Understand how to identify the Fisher information and CRLB when the score function has been written in a linear form.
- ▶ Understand what is meant by Minimum variance bound estimators (MVBES) and minimum variance bound unbiased estimators (MVBUEs), and how to determine such estimators.
- ▶ Learn how to calculate the Fisher information matrix and Cramér lower bounds for k -dimensional parameters.
- ▶ Understand what is meant by the exponential family of distributions and learn how to determine whether or not a given distribution belongs to the exponential family.