

# Introduction to structural equation modelling - advanced modelling techniques

---

Frank Pennekamp

05.11.2021

Department of Evolutionary Biology and Environmental Sciences

University of Zurich

- Advanced SEM topics covered today:


- Advanced SEM topics covered today:
  - Latent variables

- Advanced SEM topics covered today:
  - Latent variables
  - Composite variables

- Advanced SEM topics covered today:
  - Latent variables
  - Composite variables
  - Interactions


- Advanced SEM topics covered today:
  - Latent variables
  - Composite variables
  - Interactions
  - Complex survey designs

## Revisiting the meta-model



- Meta-model are conceptual models that allow to link data with theory.


## Revisiting the meta-model



- Meta-model are conceptual models that allow to link data with theory.
- So far, we have only worked with manifest (measured) variables.




## Revisiting the meta-model




- Meta-model are conceptual models that allow to link data with theory.
- So far, we have only worked with manifest (measured) variables.
- However, some aspects of a meta-model may be difficult to quantify and measure directly.

## Revisiting the meta-model




- Meta-model are conceptual models that allow to link data with theory.
- So far, we have only worked with manifest (measured) variables.
- However, some aspects of a meta-model may be difficult to quantify and measure directly.
- This is because they represent an abstract, multifaceted concept (e.g., general intelligence).

## Revisiting the meta-model



- Meta-model are conceptual models that allow to link data with theory.
- So far, we have only worked with manifest (measured) variables.
- However, some aspects of a meta-model may be difficult to quantify and measure directly.
- This is because they represent an abstract, multifaceted concept (e.g., general intelligence).
- Or, because we measure variables with error.

## Revisiting the meta-model




- Meta-model are conceptual models that allow to link data with theory.
- So far, we have only worked with manifest (measured) variables.
- However, some aspects of a meta-model may be difficult to quantify and measure directly.
- This is because they represent an abstract, multifaceted concept (e.g., general intelligence).
- Or, because we measure variables with error.
- This is where latent and composite variables are needed.

## Latent variables


---

# Latent variables




- Latent variables are unobserved, but their influence is captured by indicator variable(s).

# Latent variables



- Latent variables are unobserved, but their influence is captured by indicator variable(s).
- Latents can capture complex or conceptual properties that are difficult to quantify or measure directly.


# Latent variables



- Latent variables are unobserved, but their influence is captured by indicator variable(s).
- Latents can capture complex or conceptual properties that are difficult to quantify or measure directly.
- Graphically, latent variables are often represented by an oval node shapes (ellipses).




# Latent variables




- Direction of causality reversed: from latent variables to the observed variables (reflective indicators).

# Latent variables




- Direction of causality reversed: from latent variables to the observed variables (reflective indicators).
- Indicator variables are emergent manifestation of the underlying phenomenon represented by the latent variable.

# Latent variables



- Direction of causality reversed: from latent variables to the observed variables (reflective indicators).
- Indicator variables are emergent manifestation of the underlying phenomenon represented by the latent variable.
- All indicators should be positively correlated to the latent variable (i.e., driver).

# Latent variables



- Direction of causality reversed: from latent variables to the observed variables (reflective indicators).
- Indicator variables are emergent manifestation of the underlying phenomenon represented by the latent variable.
- All indicators should be positively correlated to the latent variable (i.e., driver).
- In contrast to indicators, latent variables are free of random or systematic measurement errors.

- Latents can be both endogenous and exogenous.

- Latents can be both endogenous and exogenous.
- *Measurement model* focuses solely on relating indicators to latent variables.

- Latents can be both endogenous and exogenous.
- *Measurement model* focuses solely on relating indicators to latent variables.
- *Structural model* is one with directed paths between latent variables.

Latents often fitted in two steps:

- 1) Confirmatory factor analysis



Latents often fitted in two steps:

- 1) Confirmatory factor analysis

- Multi-indicator latent variables test hypothesis that multiple indicator variables are generated by the same underlying process.

Latents often fitted in two steps:

1) Confirmatory factor analysis

- Multi-indicator latent variables test hypothesis that multiple indicator variables are generated by the same underlying process.
- Precursor to evaluation of any structural models in which the latent variables appear.

Latents often fitted in two steps:

1) Confirmatory factor analysis

- Multi-indicator latent variables test hypothesis that multiple indicator variables are generated by the same underlying process.
- Precursor to evaluation of any structural models in which the latent variables appear.

2) Full model including latent variable

Latents often fitted in two steps:

1) Confirmatory factor analysis

- Multi-indicator latent variables test hypothesis that multiple indicator variables are generated by the same underlying process.
- Precursor to evaluation of any structural models in which the latent variables appear.


2) Full model including latent variable

- To estimate effect of latent variable on other variables.

## Composite variables


---

## Composite variables




- Composite variables specify the influence of collections of other variables (e.g., landuse).

# Composite variables



- Composite variables specify the influence of collections of other variables (e.g., landuse).
- In comparison to latent variables, they arise from the indicators.


# Composite variables



- Composite variables specify the influence of collections of other variables (e.g., landuse).
- In comparison to latent variables, they arise from the indicators.
- Visualized by arrows pointing from the indicators to the composite (formative or causal indicators).




## Composite variables




- Composite variables specify the influence of collections of other variables (e.g., landuse).
- In comparison to latent variables, they arise from the indicators.
- Visualized by arrows pointing from the indicators to the composite (formative or causal indicators).
- Values are determined by its causes (indicators), thus error variance is set to 0.

## Composite variables




- Composite variables specify the influence of collections of other variables (e.g., landuse).
- In comparison to latent variables, they arise from the indicators.
- Visualized by arrows pointing from the indicators to the composite (formative or causal indicators).
- Values are determined by its causes (indicators), thus error variance is set to 0.
- Composites are often shown as hexangular shapes or ellipses (latent composites).

## Key differences between latent and composite variables




- Flow of causation in a latent variables is from the construct to its indicators, i.e., indicators are driven by an underlying, unmeasured process.

## Key differences between latent and composite variables




- Flow of causation in a latent variables is from the construct to its indicators, i.e., indicators are driven by an underlying, unmeasured process.
- As a latent variable represents an underlying, data-generating process, its indicators need to be correlated to each other.

## Key differences between latent and composite variables



- Flow of causation in a latent variables is from the construct to its indicators, i.e., indicators are driven by an underlying, unmeasured process.
- As a latent variable represents an underlying, data-generating process, its indicators need to be correlated to each other.
- For composite variables, the flow of causation is reversed and the indicators are independent entities.

## Key differences between latent and composite variables



- Flow of causation in a latent variables is from the construct to its indicators, i.e., indicators are driven by an underlying, unmeasured process.
- As a latent variable represents an underlying, data-generating process, its indicators need to be correlated to each other.
- For composite variables, the flow of causation is reversed and the indicators are independent entities.
- In the case of a composite variable, there is no assumption about the relation between the indicators.

# Key differences between latent and composite variables

How distinguish latent and composites:

- Rule of thumb 1: if the indicators are redundant, they likely belong to a latent variable.

# Key differences between latent and composite variables

How distinguish latent and composites:


- Rule of thumb 1: if the indicators are redundant, they likely belong to a latent variable.
- Rule of thumb 2: If the meaning of the construct changes after dropping one of the indicators, the construct likely is a composite.



## Interactions


---

# Interactions




- In nature, things often are contingent on each other.

# Interactions




- In nature, things often are contingent on each other.
- For instance, the effect of nutrients on plant growth, may depend on how disturbed the environment is.

# Interactions




- In nature, things often are contingent on each other.
- For instance, the effect of nutrients on plant growth, may depend on how disturbed the environment is.
- Such a behaviour is called an interaction, which indicates that the effect of the two main effects are different when combined.

# Interactions



- In nature, things often are contingent on each other.
- For instance, the effect of nutrients on plant growth, may depend on how disturbed the environment is.
- Such a behaviour is called an interaction, which indicates that the effect of the two main effects are different when combined.
- Both positive and negative interactions are possible.

# Interactions



- In nature, things often are contingent on each other.
- For instance, the effect of nutrients on plant growth, may depend on how disturbed the environment is.
- Such a behaviour is called an interaction, which indicates that the effect of the two main effects are different when combined.
- Both positive and negative interactions are possible.
- In regression, the interaction is represented by a coefficient that estimates the effect of the product of the two predictors.

Interactions can be modelled in different ways in lavaan:

- 1) Multiple groups

Interactions can be modelled in different ways in lavaan:

- 1) Multiple groups
- 2) Composites



```
mod <- sem(model, group = "age_class", data = dd)
```

- Multigroup fitting allows coefficients to vary among groups.

```
mod <- sem(model, group = "age_class", data = dd)
```

- Multigroup fitting allows coefficients to vary among groups.
- Lavaan offers the “group” argument to specify for which groups coefficients should be estimated.

```
mod <- sem(model, group = "age_class", data = dd)
```

- Multigroup fitting allows coefficients to vary among groups.
- Lavaan offers the “group” argument to specify for which groups coefficients should be estimated.
- Importantly, groups have to be categorical (e.g., sex, age class).

## Interactions (multi group)

Lavaan allows to introduce equality constraints on various aspects via the `group.equal` argument:

```
mod <- sem(model, group = "age_class", group.equal =  
c("regressions"), data = dd)
```

Additional constraints could be:

```
group.equal=c(  
  "intercepts",  
  "means",  
  "regressions",  
  "residuals",  
  "residual.covariances")
```

## Interactions (multi group)

- Even more control by having the same name for different parameters:

```
model <- '  
y ~ c("b1", "b1") * x1 + c("b2", "b2") * x3  
x2 ~ c("b3", "b4") * x1  
x3 ~ c("b5", "b5") * x2  
,
```

Same coefficients for all but the effect of x1 on x2.

Heywood cases: when things go  
wrong

---

## Heywood cases : when things go wrong

- Improper solutions:

## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances



## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances
  - Correlations larger than 1

## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances
  - Correlations larger than 1
- Reasons for the warnings are:

## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances
  - Correlations larger than 1
- Reasons for the warnings are:
  - 1) slightly negative error estimates (not really a problem)

## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances
  - Correlations larger than 1
- Reasons for the warnings are:
  - 1) slightly negative error estimates (not really a problem)
  - 2) indicators for latent variables need to be positively correlated

## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances
  - Correlations larger than 1
- Reasons for the warnings are:
  - 1) slightly negative error estimates (not really a problem)
  - 2) indicators for latent variables need to be positively correlated
  - 3) local non-identification

## Heywood cases : when things go wrong

- Improper solutions:
  - Negative variances
  - Correlations larger than 1
- Reasons for the warnings are:
  - 1) slightly negative error estimates (not really a problem)
  - 2) indicators for latent variables need to be positively correlated
  - 3) local non-identification
  - 4) general misspecification

## Complex sampling structure

---

- Needed when data is nested:



- Needed when data is nested:
  - within sites

## Complex sampling structure

- Needed when data is nested:
  - within sites
  - within groups such as families, nests

- Needed when data is nested:
  - within sites
  - within groups such as families, nests
- Nesting violates the principle of being independent and identically distributed.

## Complex sampling structure

- Needed when data is nested:
  - within sites
  - within groups such as families, nests
- Nesting violates the principle of being independent and identically distributed.
- Necessary to account for this structure in the data in the model.

```
library("lavaan.survey")
```

- The add-on package `lavaan.survey` allows the analysis of stratified, clustered or weighted data.

```
library("lavaan.survey")
```

- The add-on package `lavaan.survey` allows the analysis of stratified, clustered or weighted data.
- `lavaan` objects can be processed further with a specific data structure:

```
library("lavaan.survey")
```

- The add-on package lavaan.survey allows the analysis of stratified, clustered or weighted data.
- lavaan objects can be processed further with a specific data structure:
  - 1) initialize the design

```
library("lavaan.survey")
```

- The add-on package `lavaan.survey` allows the analysis of stratified, clustered or weighted data.
- `lavaan` objects can be processed further with a specific data structure:
  - 1) initialize the design
  - 2) post-process the `lavaan` object and compute the adjusted results.



```
library("lavaan.survey")
```

- The add-on package `lavaan.survey` allows the analysis of stratified, clustered or weighted data.
- `lavaan` objects can be processed further with a specific data structure:
  - 1) initialize the design
  - 2) post-process the `lavaan` object and compute the adjusted results.
  - 3) Result is a corrected `lavaan` object.

```
design <- svydesign(ids = ~ plot, strata = ~ field, nest = TRUE,  
summary(design)
```

```
fit.nested <- lavaan.survey(lavaan.fit = model, survey.design =
```

- Needed to specify the study design

## Complex sampling structure

```
design <- svydesign(ids = ~ plot, strata = ~ field, nest = TRUE,  
summary(design)
```

```
fit.nested <- lavaan.survey(lavaan.fit = model, survey.design =
```

- Needed to specify the study design
- Here we have plots (**ids**) nested in fields (**strata**)

## Complex sampling structure

```
design <- svydesign(ids = ~ plot, strata = ~ field, nest = TRUE,  
summary(design)
```

```
fit.nested <- lavaan.survey(lavaan.fit = model, survey.design =
```

- Needed to specify the study design
- Here we have plots (**ids**) nested in fields (**strata**)
- Next, we can refit the **simple** model from before with **lavaan.survey** using the specified study design as an argument.

Questions?

---

## Live coding session

---

Your turn: working with the  
Seabloom dataset

---

## Exercise 1

Start with the following model:

```
library("lavaan")

simple <-
"mass.above ~ nadd + disk + rich + even + precip.mm
rich ~ nadd + precip.mm
even ~ nadd + precip.mm

rich ~~ even"

fit.simple <- sem(simple, data = seabloom, estimator = "MLM")
summary(fit.simple)
```



- Construct latent variable diversity based on richness, evenness and  
ens.pie

- Construct latent variable diversity based on richness, evenness and  
ens.pie
  - Run confirmatory factor analysis

- Construct latent variable diversity based on richness, evenness and  
ens.pie
  - Run confirmatory factor analysis
  - What do you conclude?

## Exercise 3

- Construct latent variable diversity only based on richness and evenness

- Construct latent variable diversity only based on richness and evenness
  - Run confirmatory factor analysis

- Construct latent variable diversity only based on richness and evenness
  - Run confirmatory factor analysis
  - Include latent variable into full model

- Construct latent variable diversity only based on richness and evenness
  - Run confirmatory factor analysis
  - Include latent variable into full model
  - What do you conclude?

- Construct composite variable landuse based on nutrients and disturbance (disk):



- Construct composite variable landuse based on nutrients and disturbance (disk):
  - What do you conclude?

- Construct composite variable landuse based on nutrients and disturbance (disk):
  - What do you conclude?
  - Build composite manually.

- Investigate possible interaction between disturbance and nutrient addition on AGB:

- Investigate possible interaction between disturbance and nutrient addition on AGB:
  - Construct composite variable to capture a possible interaction:

- Investigate possible interaction between disturbance and nutrient addition on AGB:
  - Construct composite variable to capture a possible interaction:
  - What do you conclude?

- Investigate possible interaction between disturbance and nutrient addition on AGB:
  - Construct composite variable to capture a possible interaction:
  - What do you conclude?
  - Incorporate composite into full model

- Investigate possible interaction between disturbance and nutrient addition on AGB:

- Investigate possible interaction between disturbance and nutrient addition on AGB:
  - Use multi-group fitting to explore interaction.



- Investigate possible interaction between disturbance and nutrient addition on AGB:
  - Use multi-group fitting to explore interaction.
  - What do you conclude?

- Account for nested experimental design with the `lavaan.survey` package:

- Account for nested experimental design with the `lavaan.survey` package:
  - Add individual plots nested in fields.