

Artificial Intelligence I 2023/2024

Week 8 Tutorial and Additional Exercises

Hierarchical Clustering and Evaluation of Clustering Algorithms

School of Computer Science

4th of March 2024

In this tutorial...

In this tutorial we will be covering

- Hierarchical Clustering.
- Cutting the Dendrogram.
- Supervised and unsupervised clustering validation criteria.
- Silhouette coefficient.
- Classification-oriented validation criteria
- Similarity-oriented validation criteria.

Inter-Cluster Dissimilarity Metrics

- Distance metrics can be generalised for clusters to define inter-cluster dissimilarity measures. Let C_1 and C_2 be clusters containing n_1 and n_2 examples respectively. Some examples of distance metrics between C_1 and C_2 are:

- 1 *Single linkage* is defined as

$$d_{SL}(C_1, C_2) := \min_{\mathbf{x}^{(1)} \in C_1, \mathbf{x}^{(2)} \in C_2} \text{Dist}(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}).$$

- 2 *Complete linkage* is defined as

$$d_{CL}(C_1, C_2) := \max_{\mathbf{x}^{(1)} \in C_1, \mathbf{x}^{(2)} \in C_2} \text{Dist}(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}).$$

- 3 *Group Average linkage* is defined as

$$d_{GL}(C_1, C_2) := \frac{1}{n_1 n_2} \sum_{\mathbf{x}^{(1)} \in C_1} \sum_{\mathbf{x}^{(2)} \in C_2} \text{Dist}(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}).$$

- $\text{Dist}(\cdot)$ can be any distance function between vectors.

Hierarchical Clustering

Recall the formal algorithm of *Hierarchical Clustering*.

Algorithm 1: Hierarchical clustering.

Input: Distance matrix corresponding to the data set $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}$

Output: Dendrogram.

```
1 repeat
2   Find two clusters  $C_1, C_2$  with the smallest inter-cluster dissimilarity. That is,

                                     
$$\arg \min_{C_1, C_2} d_A(C_1, C_2)$$


       where  $A \in \{SL, CL, GL\}$  denotes single linkage (SL), complete linkage (CL)
       or group linkage (GL);
3   Merge together  $C_1, C_2$  into a single cluster;
4   Note the clusters merged and their corresponding linkage  $d_A(\cdot, \cdot)$  in a
       dendrogram.
5 until Only one cluster remains.;
6 return Dendrogram.
```

Exercise 1

- Consider the set with 6 examples and the following distance matrix (for some choice of distance function).

	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$\mathbf{x}^{(3)}$	$\mathbf{x}^{(4)}$	$\mathbf{x}^{(5)}$	$\mathbf{x}^{(6)}$
$\mathbf{x}^{(1)}$	0	0.20	0.15	0.76	0.54	0.31
$\mathbf{x}^{(2)}$	0.20	0	0.89	0.18	0.66	0.27
$\mathbf{x}^{(3)}$	0.15	0.89	0	0.82	0.73	0.56
$\mathbf{x}^{(4)}$	0.76	0.18	0.82	0	0.42	0.39
$\mathbf{x}^{(5)}$	0.54	0.66	0.73	0.42	0	0.51
$\mathbf{x}^{(6)}$	0.31	0.27	0.56	0.39	0.51	0

- Use Hierarchical clustering in algorithm 1 to merge all examples into a single cluster.
- Use **single linkage** as the inter-cluster dissimilarity metric.
- Sketch the dendrogram you found along the way clearly depicting the height at which two clusters fuse.

Exercise 1: Solution

- We start with each example in its own cluster and calculate the distance matrix for these clusters.

	$\{\mathbf{x}^{(1)}\}$	$\{\mathbf{x}^{(2)}\}$	$\{\mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}\}$	0	0.20	0.15	0.76	0.54	0.31
$\{\mathbf{x}^{(2)}\}$	0.20	0	0.89	0.18	0.66	0.27
$\{\mathbf{x}^{(3)}\}$	0.15	0.89	0	0.82	0.73	0.56
$\{\mathbf{x}^{(4)}\}$	0.76	0.18	0.82	0	0.42	0.39
$\{\mathbf{x}^{(5)}\}$	0.54	0.66	0.73	0.42	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.31	0.27	0.56	0.39	0.51	0

- The closest clusters are $\{\mathbf{x}^{(1)}\}$ and $\{\mathbf{x}^{(3)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}, \{\mathbf{x}^{(2)}\}, \{\mathbf{x}^{(4)}\}, \{\mathbf{x}^{(5)}\}, \{\mathbf{x}^{(6)}\}.$$

Exercise 1: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(2)}\}$	$\{\mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	0	0.20	0.76	0.54	0.31
$\{\mathbf{x}^{(2)}\}$	0.20	0	0.18	0.66	0.27
$\{\mathbf{x}^{(4)}\}$	0.76	0.18	0	0.42	0.39
$\{\mathbf{x}^{(5)}\}$	0.54	0.66	0.42	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.31	0.27	0.39	0.51	0

- The closest clusters are $\{\mathbf{x}^{(2)}\}$ and $\{\mathbf{x}^{(4)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}, \{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}, \{\mathbf{x}^{(5)}\}, \{\mathbf{x}^{(6)}\}.$$

Exercise 1: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	0	0.20	0.54	0.31
$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}$	0.20	0	0.42	0.27
$\{\mathbf{x}^{(5)}\}$	0.54	0.42	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.31	0.27	0.51	0

- The closest clusters are $\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$ and $\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}\}, \{\mathbf{x}^{(5)}\}, \{\mathbf{x}^{(6)}\}.$$

Exercise 1: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}\}$	0	0.42	0.27
$\{\mathbf{x}^{(5)}\}$	0.42	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.27	0.51	0

- The closest clusters are $\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}\}$ and $\{\mathbf{x}^{(6)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}, \{\mathbf{x}^{(5)}\}.$$

Exercise 1: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}$	$\{\mathbf{x}^{(5)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}$	0	0.42
$\{\mathbf{x}^{(5)}\}$	0.42	0


- The closest clusters are $\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}$ and $\{\mathbf{x}^{(5)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(5)}, \mathbf{x}^{(6)}\}.$$

- Finally, we construct the dendrogram.

Exercise 1: Solution (continued)

The dendrogram is the following:



Exercise 2

- Reconsider the set with 6 examples and the following distance matrix (for some choice of distance function).

	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$\mathbf{x}^{(3)}$	$\mathbf{x}^{(4)}$	$\mathbf{x}^{(5)}$	$\mathbf{x}^{(6)}$
$\mathbf{x}^{(1)}$	0	0.20	0.15	0.76	0.54	0.31
$\mathbf{x}^{(2)}$	0.20	0	0.89	0.18	0.66	0.27
$\mathbf{x}^{(3)}$	0.15	0.89	0	0.82	0.73	0.56
$\mathbf{x}^{(4)}$	0.76	0.18	0.82	0	0.42	0.39
$\mathbf{x}^{(5)}$	0.54	0.66	0.73	0.42	0	0.51
$\mathbf{x}^{(6)}$	0.31	0.27	0.56	0.39	0.51	0

- Use Hierarchical clustering in algorithm 1 to merge all examples into a single cluster.
- Use **complete linkage** as the inter-cluster dissimilarity metric.
- Sketch the dendrogram you found along the way.

Exercise 2: Solution

- We start with each example in its own cluster and calculate the distance matrix for these clusters.

	$\{\mathbf{x}^{(1)}\}$	$\{\mathbf{x}^{(2)}\}$	$\{\mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}\}$	0	0.20	0.15	0.76	0.54	0.31
$\{\mathbf{x}^{(2)}\}$	0.20	0	0.89	0.18	0.66	0.27
$\{\mathbf{x}^{(3)}\}$	0.15	0.89	0	0.82	0.73	0.56
$\{\mathbf{x}^{(4)}\}$	0.76	0.18	0.82	0	0.42	0.39
$\{\mathbf{x}^{(5)}\}$	0.54	0.66	0.73	0.42	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.31	0.27	0.56	0.39	0.51	0

- The closest clusters are $\{\mathbf{x}^{(1)}\}$ and $\{\mathbf{x}^{(3)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}, \{\mathbf{x}^{(2)}\}, \{\mathbf{x}^{(4)}\}, \{\mathbf{x}^{(5)}\}, \{\mathbf{x}^{(6)}\}.$$

Exercise 2: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(2)}\}$	$\{\mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	0	0.89	0.82	0.73	0.56
$\{\mathbf{x}^{(2)}\}$	0.89	0	0.18	0.66	0.27
$\{\mathbf{x}^{(4)}\}$	0.82	0.18	0	0.42	0.39
$\{\mathbf{x}^{(5)}\}$	0.73	0.66	0.42	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.56	0.27	0.39	0.51	0

- The closest clusters are $\{\mathbf{x}^{(2)}\}$ and $\{\mathbf{x}^{(4)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}, \{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}, \{\mathbf{x}^{(5)}\}, \{\mathbf{x}^{(6)}\}.$$

Exercise 2: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}$	$\{\mathbf{x}^{(5)}\}$	$\{\mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	0	0.89	0.73	0.56
$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}$	0.89	0	0.66	0.39
$\{\mathbf{x}^{(5)}\}$	0.73	0.66	0	0.51
$\{\mathbf{x}^{(6)}\}$	0.56	0.39	0.51	0

- The closest clusters are $\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}\}$ and $\{\mathbf{x}^{(6)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}, \{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}, \{\mathbf{x}^{(5)}\}.$$

Exercise 2: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}$	$\{\mathbf{x}^{(5)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	0	0.89	0.73
$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}$	0.89	0	0.66
$\{\mathbf{x}^{(5)}\}$	0.73	0.66	0

- The closest clusters are $\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}$ and $\{\mathbf{x}^{(5)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}, \{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(5)}, \mathbf{x}^{(6)}\}.$$

Exercise 2: Solution (continued)

- We then recalculate the distance matrix for the new clusters.

	$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(5)}, \mathbf{x}^{(6)}\}$
$\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$	0	0.89
$\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(5)}, \mathbf{x}^{(6)}\}$	0.89	0


- The closest clusters are $\{\mathbf{x}^{(1)}, \mathbf{x}^{(3)}\}$ and $\{\mathbf{x}^{(2)}, \mathbf{x}^{(4)}, \mathbf{x}^{(5)}, \mathbf{x}^{(6)}\}$.
- The new clusters are

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(5)}, \mathbf{x}^{(6)}\}.$$

- Finally, we construct the dendrogram.

Exercise 2: Solution (continued)

The dendrogram is the following:



Cutting the Dendrogram

- In Hierarchical Clustering, we can impose a threshold on the inter-cluster distance or on the number of clusters.
- When this threshold is surpassed, the algorithm terminates without forming any further clusters, and returns the clusters formed so far.
- Different thresholds can result in different clusters.

Exercise 3

- Reconsider the set with 6 examples and the following distance matrix (for some choice of distance function).

	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$\mathbf{x}^{(3)}$	$\mathbf{x}^{(4)}$	$\mathbf{x}^{(5)}$	$\mathbf{x}^{(6)}$
$\mathbf{x}^{(1)}$	0	0.20	0.15	0.76	0.54	0.31
$\mathbf{x}^{(2)}$	0.20	0	0.89	0.18	0.66	0.27
$\mathbf{x}^{(3)}$	0.15	0.89	0	0.82	0.73	0.56
$\mathbf{x}^{(4)}$	0.76	0.18	0.82	0	0.42	0.39
$\mathbf{x}^{(5)}$	0.54	0.66	0.73	0.42	0	0.51
$\mathbf{x}^{(6)}$	0.31	0.27	0.56	0.39	0.51	0

- Use Hierarchical clustering but impose a threshold of 0.35 on the inter-cluster distance. Write the final clusters.
- Use Hierarchical clustering but impose a threshold of 3 on the number of clusters. Write the final clusters.
- Use **single linkage** as the inter-cluster dissimilarity metric.

Exercise 3: Solution


- If we impose a threshold of 0.35 on the inter-cluster distance, the final clusters will be

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}, \mathbf{x}^{(6)}\}, \{\mathbf{x}^{(5)}\}.$$

- We next draw the dendrogram with the horizontal cut.

Exercise 3: Solution (continued)

The dendrogram with the horizontal cut is the following:



Exercise 3: Solution (continued)


- If we impose a threshold of 3 on the number of clusters, the final clusters will be

$$\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \mathbf{x}^{(3)}, \mathbf{x}^{(4)}\}, \{\mathbf{x}^{(5)}\}, \{\mathbf{x}^{(6)}\}.$$

- We next draw the dendrogram with the horizontal cut.

Exercise 3: Solution (continued)

The dendrogram with the horizontal cut is the following:



Silhouette Coefficient

- Let $\mathbf{x}^{(i)}$ be an example in cluster C , and define
 - a_i to be the average distance of $\mathbf{x}^{(i)}$ to all other examples in C , i.e.,

$$a_i := \frac{\sum_{\mathbf{x} \in C, \mathbf{x} \neq \mathbf{x}^{(i)}} d(\mathbf{x}^{(i)}, \mathbf{x})}{(\text{no. of examples in cluster } C) - 1}.$$

- b_i to be the minimum of the average distance of $\mathbf{x}^{(i)}$ to examples in other clusters, i.e.

$$b_i := \min_{\substack{k=1, \dots, K \\ C_k \neq C}} \frac{\sum_{\mathbf{x} \in C_k} d(\mathbf{x}^{(i)}, \mathbf{x})}{\text{no. of examples in } C_k}.$$

- The SC for $\mathbf{x}^{(i)}$ is defined as

$$s_i := \frac{b_i - a_i}{\max\{a_i, b_i\}}.$$

Silhouette Coefficient (continued)

- The SC of a cluster C is defined as

$$s_C := \frac{\sum_{\{i: \mathbf{x}^{(i)} \in C\}} s_i}{\text{no. of examples in cluster } C}.$$

- The SC of a clustering structure \mathcal{C} with N examples is defined as

$$s_{\mathcal{C}} := \frac{\sum_{i=1}^N s_i}{N}.$$

Exercise 4

- Consider a dataset with 4 examples, clustered by an algorithm as

$$C_1 = \{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}\}, \quad C_2 = \{\mathbf{x}^{(3)}, \mathbf{x}^{(4)}\}.$$

- The distance matrix for these examples is the following

	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$\mathbf{x}^{(3)}$	$\mathbf{x}^{(4)}$
$\mathbf{x}^{(1)}$	0	0.10	0.65	0.55
$\mathbf{x}^{(2)}$	0.10	0	0.70	0.60
$\mathbf{x}^{(3)}$	0.65	0.70	0	0.90
$\mathbf{x}^{(4)}$	0.55	0.60	0.90	0

- Compute the SC for each point, for each cluster, and for the overall clustering structure $\mathcal{C} = \{C_1, C_2\}$.
- Comment on the suitability of examples assigned to C_1 .

Exercise 4: Solution

- We first find the a_i 's and the b_i 's

$$a_1 = 0.1, a_2 = 0.1, a_3 = 0.9, a_4 = 0.9.$$

$$b_1 = 0.6, b_2 = 0.65, b_3 = 0.675, b_4 = 0.575.$$

- We then find the SC of each example

$$s_1 = 0.8333, s_2 = 0.8461, s_3 = -0.25, s_4 = -0.3611.$$

- We then find the SC of each cluster

$$s_{C_1} = 0.8397, s_{C_2} = -0.3055.$$

- We then find the SC of the clustering structure

$$s_C = 0.2670.$$

Classification-oriented validation criteria

- Consider a set of L different classes, clustered into K clusters.
- *Precision* of cluster i with respect to class j

$$precision(i, j) := \frac{\text{no. of examples of class } j \text{ in cluster } i}{\text{no. of examples in cluster } i}.$$

- *Recall* of cluster i with respect to class j

$$recall(i, j) := \frac{\text{no. of examples of class } j \text{ in cluster } i}{\text{no. of examples in class } j}.$$

- *F-measure* of cluster i with respect to class j

$$F(i, j) := \frac{2 \cdot precision(i, j) \cdot recall(i, j)}{precision(i, j) + recall(i, j)}.$$

Classification-oriented validation criteria (continued)

- The *entropy* of cluster i is defined as

$$e_i := - \sum_{j=1}^L \text{precision}(i, j) \cdot \log_2(\text{precision}(i, j)),$$

where $-x \log_2 x := 0$, when $x = 0$.

- The *total entropy* of the set of clusters is defined as

$$e := \sum_{i=1}^K \frac{\text{no. of examples in cluster } i}{\text{total no. of examples}} e_i.$$

- We want a low entropy.

Classification-oriented validity measures (continued)

- The *purity* of cluster i is defined as

$$p_i := \max_j \text{precision}(i, j).$$

- The *overall purity* of the set of clusters is defined as

$$p := \sum_{i=1}^K \frac{\text{no. of examples in cluster } i}{\text{total no. of examples}} p_i.$$

- We want a high purity.

Exercise 5

- Consider the set with 10 examples and 3 classes, clustered into 3 clusters (**classes and clusters are not the same**)

Example	Class	Cluster	Example	Class	Cluster
$\mathbf{x}^{(1)}$	1	1	$\mathbf{x}^{(6)}$	3	1
$\mathbf{x}^{(2)}$	3	2	$\mathbf{x}^{(7)}$	2	2
$\mathbf{x}^{(3)}$	2	3	$\mathbf{x}^{(8)}$	2	2
$\mathbf{x}^{(4)}$	1	1	$\mathbf{x}^{(9)}$	1	3
$\mathbf{x}^{(5)}$	3	2	$\mathbf{x}^{(10)}$	2	1

- Write down the confusion matrix.
- Compute the following
 - 1 $precision(1, 3)$.
 - 2 $recall(1, 3)$.
 - 3 $F(1, 3)$.
 - 4 e_2 .
 - 5 p_2 .

Exercise 5: Solution

- The confusion matrix is the following

	Cluster 1	Cluster 2	Cluster 3	Total
Class 1	2	0	1	3
Class 2	1	2	1	4
Class 3	1	2	0	3
Total	4	4	2	10

- We also compute the following

- ① $\text{precision}(1, 3) = 1/4$.
- ② $\text{recall}(1, 3) = 1/3$.
- ③ $F(1, 3) = 2/7$.
- ④ $e_2 = 1$.
- ⑤ $p_2 = 1/2$.

Similarity-oriented validation criteria

- Consider a set of N examples of different classes, clustered into clusters.
- The *ideal cluster similarity matrix* is an $N \times N$ matrix whose ij -th element equals 1 if examples i and j are in the same cluster, and 0 otherwise.
- The *ideal class similarity matrix* is an $N \times N$ matrix whose ij -th element equals 1 if examples i and j are in the same class, and 0 otherwise.
- We can compute the correlation between these two matrices.
- We can also use binary similarity-based measures.

Binary similarity-based measures

- Consider a set of N examples of different classes, clustered into clusters and define the following
 - 1 $f_{00} :=$ no. of pairs having different class and different cluster.
 - 2 $f_{01} :=$ no. of pairs having different class and same cluster.
 - 3 $f_{10} :=$ no. of pairs having same class and different cluster.
 - 4 $f_{11} :=$ no. of pairs having same class and same cluster.
- The *Rand statistic* is defined as

$$\text{Rand statistic} = \frac{f_{00} + f_{11}}{f_{00} + f_{01} + f_{10} + f_{11}}.$$

- The *Jaccard coefficient* is defined as

$$\text{Jaccard coefficient} = \frac{f_{11}}{f_{01} + f_{10} + f_{11}}.$$

Exercise 6

- Reconsider the first five examples of the previous set with 3 classes, clustered into 3 clusters

Example	Class	Cluster
$\mathbf{x}^{(1)}$	1	1
$\mathbf{x}^{(2)}$	3	2
$\mathbf{x}^{(3)}$	2	3
$\mathbf{x}^{(4)}$	1	1
$\mathbf{x}^{(5)}$	3	2

- Write down the ideal cluster similarity matrix and the ideal class similarity matrix.
- Compute the Rand statistic and the Jaccard coefficient.

Exercise 6: Solution

- The ideal cluster similarity matrix is the following

	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$\mathbf{x}^{(3)}$	$\mathbf{x}^{(4)}$	$\mathbf{x}^{(5)}$
$\mathbf{x}^{(1)}$	1	0	0	1	0
$\mathbf{x}^{(2)}$	0	1	0	0	1
$\mathbf{x}^{(3)}$	0	0	1	0	0
$\mathbf{x}^{(4)}$	1	0	0	1	0
$\mathbf{x}^{(5)}$	0	1	0	0	1

- The ideal class similarity matrix is the following

	$\mathbf{x}^{(1)}$	$\mathbf{x}^{(2)}$	$\mathbf{x}^{(3)}$	$\mathbf{x}^{(4)}$	$\mathbf{x}^{(5)}$
$\mathbf{x}^{(1)}$	1	0	0	1	0
$\mathbf{x}^{(2)}$	0	1	0	0	1
$\mathbf{x}^{(3)}$	0	0	1	0	0
$\mathbf{x}^{(4)}$	1	0	0	1	0
$\mathbf{x}^{(5)}$	0	1	0	0	1

Exercise 6: Solution (continued)

- We first compute the following
 - 1 $f_{00} = 8.$
 - 2 $f_{01} = 0.$
 - 3 $f_{10} = 0.$
 - 4 $f_{11} = 2.$
- Therefore, *Rand statistic* = 1.
- Also, *Jaccard coefficient* = 1.

Advanced Material

(OPTIONAL) Advanced Exercise 1


- Let C_1 , C_2 and C_3 be clusters. Prove the following:
 - ① $d_{SL}(C_1, C_2 \cup C_3) = \min\{d_{SL}(C_1, C_2), d_{SL}(C_1, C_3)\}$.
 - ② $d_{CL}(C_1, C_2 \cup C_3) = \max\{d_{CL}(C_1, C_2), d_{CL}(C_1, C_3)\}$.
- Recall that
 - ① $C \cup C' = \{\mathbf{x} : \mathbf{x} \in C \vee \mathbf{x} \in C'\}$.
 - ② $d_{SL}(C, C') = \min_{\mathbf{x} \in C, \mathbf{x}' \in C'} \text{Dist}(\mathbf{x}, \mathbf{x}')$.
 - ③ $d_{CL}(C, C') = \max_{\mathbf{x} \in C, \mathbf{x}' \in C'} \text{Dist}(\mathbf{x}, \mathbf{x}')$.
 - ④ $\text{Dist}(\cdot, \cdot)$ is some distance function for vectors.
- Hint: $\min_{\mathbf{x} \in C \vee \mathbf{x} \in C'} f(\mathbf{x}) = \min\{\min_{\mathbf{x} \in C} f(\mathbf{x}), \min_{\mathbf{x} \in C'} f(\mathbf{x})\}$, for any sets C, C' , and real-valued function $f(\cdot)$. The same holds if we replace all the min's with max's.

(OPTIONAL) Advanced Exercise 1: Solution

- ①
$$\begin{aligned} d_{SL}(C_1, C_2 \cup C_3) &= \min_{\mathbf{x} \in C_1, \mathbf{x}' \in C_2 \cup C_3} \text{Dist}(\mathbf{x}, \mathbf{x}') = \\ &= \min_{\mathbf{x} \in C_1, \mathbf{x}' \in C_2 \vee \mathbf{x}' \in C_3} \text{Dist}(\mathbf{x}, \mathbf{x}') = \\ &= \min\{\min_{\mathbf{x} \in C_1, \mathbf{x}' \in C_2} \text{Dist}(\mathbf{x}, \mathbf{x}'), \min_{\mathbf{x} \in C_1, \mathbf{x}' \in C_3} \text{Dist}(\mathbf{x}, \mathbf{x}')\} = \\ &= \min\{d_{SL}(C_1, C_2), d_{SL}(C_1, C_3)\}. \end{aligned}$$
- ②
$$\begin{aligned} d_{CL}(C_1, C_2 \cup C_3) &= \max_{\mathbf{x} \in C_1, \mathbf{x}' \in C_2 \cup C_3} \text{Dist}(\mathbf{x}, \mathbf{x}') = \\ &= \max_{\mathbf{x} \in C_1, \mathbf{x}' \in C_2 \vee \mathbf{x}' \in C_3} \text{Dist}(\mathbf{x}, \mathbf{x}') = \\ &= \max\{\max_{\mathbf{x} \in C_1, \mathbf{x}' \in C_2} \text{Dist}(\mathbf{x}, \mathbf{x}'), \max_{\mathbf{x} \in C_1, \mathbf{x}' \in C_3} \text{Dist}(\mathbf{x}, \mathbf{x}')\} = \\ &= \max\{d_{CL}(C_1, C_2), d_{CL}(C_1, C_3)\}. \end{aligned}$$

Any questions?

Until the next time...



Thank you for your attention!