

Cite as: J. V. Moreno-Mayar *et al.*, *Science* 10.1126/science.aav2621 (2018).

# Early human dispersals within the Americas

**J. Víctor Moreno-Mayar<sup>1\*</sup>, Lasse Vinner<sup>1\*</sup>, Peter de Barros Damgaard<sup>1\*</sup>, Constanza de la Fuente<sup>1\*</sup>, Jeffrey Chan<sup>2\*</sup>, Jeffrey P. Spence<sup>3\*</sup>, Morten E. Allentoft<sup>1</sup>, Tharsika Vimala<sup>1</sup>, Fernando Racimo<sup>1</sup>, Thomaz Pinotti<sup>4</sup>, Simon Rasmussen<sup>5</sup>, Ashot Margaryan<sup>1,6</sup>, Miren Iraeta Orbegozo<sup>1</sup>, Dorothea Mylopotamitaki<sup>1</sup>, Matthew Wooller<sup>7</sup>, Clement Bataille<sup>8</sup>, Lorena Becerra-Valdivia<sup>9</sup>, David Chivall<sup>9</sup>, Daniel Comeskey<sup>9</sup>, Thibaut Devière<sup>9</sup>, Donald K. Grayson<sup>10</sup>, Len George<sup>11</sup>, Harold Harry<sup>12</sup>, Verner Alexandersen<sup>13</sup>, Charlotte Primeau<sup>13</sup>, Jon Erlandson<sup>14</sup>, Claudia Rodrigues-Carvalho<sup>15</sup>, Silvia Reis<sup>15</sup>, Murilo Q. R. Bastos<sup>15</sup>, Jerome Cybulski<sup>16,17,18</sup>, Carlos Vullo<sup>19</sup>, Flavia Morello<sup>20</sup>, Miguel Vilar<sup>21</sup>, Spencer Wells<sup>22</sup>, Kristian Gregersen<sup>1</sup>, Kasper Lykke Hansen<sup>1</sup>, Niels Lynnerup<sup>13</sup>, Marta Mirazón Lahr<sup>23</sup>, Kurt Kjær<sup>1</sup>, André Strauss<sup>24,25</sup>, Marta Alfonso-Durruty<sup>26</sup>, Antonio Salas<sup>27,28</sup>, Hannes Schroeder<sup>1</sup>, Thomas Higham<sup>9</sup>, Ripan S. Malhi<sup>29</sup>, Jeffrey T. Rasic<sup>30</sup>, Luiz Souza<sup>31</sup>, Fabricio R. Santos<sup>4</sup>, Anna-Sapfo Malaspina<sup>32</sup>, Martin Sikora<sup>1</sup>, Rasmus Nielsen<sup>1,33,34</sup>, Yun S. Song<sup>2,33,35†</sup>, David J. Meltzer<sup>1,36†</sup>, Eske Willerslev<sup>1,37,38‡</sup>**

<sup>1</sup>Centre for GeoGenetics, Natural History Museum of Denmark, University of Copenhagen, 1350 Copenhagen, Denmark. <sup>2</sup>Computer Science Division, University of California, Berkeley, CA 94720, USA. <sup>3</sup>Computational Biology Graduate Group, University of California, Berkeley, CA 94720, USA. <sup>4</sup>Departamento de Genética, Ecología e Evolución, Instituto de Ciencias Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil. <sup>5</sup>Department of Bio and Health Informatics, Technical University of Denmark, Lyngby, Denmark. <sup>6</sup>Institute of Molecular Biology, National Academy of Sciences, 7 Hasratian St., 0014, Yerevan, Armenia. <sup>7</sup>Alaska Stable Isotope Facility, Water and Environmental Research Center, University of Alaska Fairbanks, Fairbanks, AK 99775, USA. <sup>8</sup>Department of Earth and Environmental Sciences, University of Ottawa, Ottawa, Canada. <sup>9</sup>Oxford Radiocarbon Accelerator Unit, Research Lab for Archaeology and the History of Art, School of Archaeology, University of Oxford, 1-2 South Parks Road, Oxford, OX1 3TG, UK. <sup>10</sup>Department of Anthropology and Quaternary Research Center, University of Washington, Seattle, WA 98185, USA. <sup>11</sup>Fallon Paiute-Shoshone Tribe, 565 Rio Vista Drive, Fallon, NV 89406, USA. <sup>12</sup>Stswecem'c/Xgat'tem Band, Dog Creek, BC V0L 1J0, Canada. <sup>13</sup>Laboratory of Biological Anthropology, Department of Forensic Medicine, University of Copenhagen, Copenhagen, Denmark. <sup>14</sup>University of Oregon Museum of Natural and Cultural History, Eugene, OR 97403, USA.

<sup>15</sup>Departamento de Antropología, Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil. <sup>16</sup>Canadian Museum of History, Gatineau, Québec K1A 0M8, Canada. <sup>17</sup>Department of Anthropology, University of Western Ontario, London, Ontario, Canada. <sup>18</sup>Department of Archaeology, Simon Fraser University, Burnaby, BC, Canada. <sup>19</sup>Equipo Argentino de Antropología Forense, Independencia 644-3A, Edif. EME1, Córdoba, Argentina. <sup>20</sup>Universidad de Magallanes, Instituto de la Patagonia, Av. Manuel Bulnes, 01890, Punta Arenas, Región de Magallanes y Antártica, Chile. <sup>21</sup>National Geographic Society, 1145 17th Street NW, Washington, DC 20036, USA. <sup>22</sup>Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA. <sup>23</sup>Leverhulme Centre for Human Evolutionary Studies, Department of Archaeology, University of Cambridge, Cambridge, UK. <sup>24</sup>Museu de Arqueología e Etnología, Universidade de São Paulo. Av. Prof. Almeida Prado, 1466 Butantã, 05508-070, São Paulo, Brazil. <sup>25</sup>Institute for Archaeological Sciences, Palaeoanthropology and Senckenberg Centre for Human Evolution and Palaeoenvironment, University of Tübingen, 72070 Tübingen, Germany. <sup>26</sup>SASW Department, Kansas State University, 1603 Old Claflin Place, Manhattan, KS 66506, USA. <sup>27</sup>Unidade de Xenética, Departamento de Anatomía Patológica e Ciencias Forenses, Instituto de Ciencias Forenses, Facultade de Medicina, Universidade de Santiago de Compostela (USC), Spain. <sup>28</sup>GenPoB Research Group, Instituto de Investigaciones Sanitarias (IDIS), Hospital Clínico Universitario de Santiago (SERGAS), Galicia, Spain. <sup>29</sup>Department of Anthropology, School of Integrative Biology, Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. <sup>30</sup>U.S. National Park Service, 4175 Geist Rd., Fairbanks, AK 99709, USA. <sup>31</sup>LACICOR – Conservation Science Laboratory, CECOR – Center for Conservation and Restoration of Cultural Properties, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil. <sup>32</sup>Department of Computational Biology, University of Lausanne, Swiss Institute of Bioinformatics, Lausanne, Switzerland. <sup>33</sup>Department of Statistics, University of California, Berkeley, CA 94720, USA. <sup>34</sup>Department of Integrative Biology, University of California, Berkeley, CA 94720, USA. <sup>35</sup>Chan Zuckerberg Biohub, San Francisco, CA 94158, USA. <sup>36</sup>Department of Anthropology, Southern Methodist University, Dallas, TX 75275, USA. <sup>37</sup>Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK. <sup>38</sup>Wellcome Trust Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge CB10 1SA, UK.

\*These authors contributed equally to this work.

†Corresponding author. Email: ewillerslev@smm.ku.dk (E.W.); dmeltzer@smu.edu (D.J.M.); yss@berkeley.edu (Y.S.S.)

**Studies of the peopling of the Americas have focused on the timing and number of initial migrations. Less attention has been paid to the subsequent spread of people within the Americas. We sequenced 15 ancient human genomes spanning Alaska to Patagonia; six are  $\geq 10,000$  years old (up to  $\sim 18\times$  coverage). All are most closely related to Native Americans, including an Ancient Beringian individual, and two morphologically distinct “Paleoamericans.” We find evidence of rapid dispersal and early diversification, including previously unknown groups, as people moved south. This resulted in multiple independent, geographically uneven migrations, including one that provides clues of a Late Pleistocene Australasian genetic signal, and a later Mesoamerican-related expansion. These led to complex and dynamic population histories from North to South America.**

Previous genomic studies have estimated that ancestral Native Americans (NA) diverged from Siberian and East Asian populations around  $25,000 \pm 1100$  years ago ( $25 \pm 1.1$  ka) (1, 2), followed by a split  $22\text{--}18.1$  ka within that ancestral

lineage between later NA and Ancient Beringians (AB). Subsequently, NA diverged into two branches, Northern Native Americans (NNA) and Southern Native Americans (SNA)  $\sim 17.5\text{--}14.6$  ka (2–4), a process inferred to have taken

place south of eastern Beringia (present day Alaska and western Yukon Territory). All contemporary and ancient NA individuals for whom genome-wide data have been generated prior to this study, derive from either the NNA or SNA branch.

However, there is disagreement over claims of earlier migrations into the Americas, possibly related to Australasians or by bearers of a distinctive cranial form (“Paleoamericans”) (5, 6). Whether there were additional splits within the Americas, how many migratory movements north and south took place, and the speed of human dispersal at different times and regions, are contentious. In contrast to models based on contemporary and Pleistocene age genetic data (3, 4), genomic studies of later Holocene human remains indicate there was post-divergence admixture between basal Native American groups (7). Overall, the degree of population isolation, admixture, or continuity in different geographic regions of the Americas after initial settlement is poorly understood (7–9).

Genome sequences from Late Pleistocene/Early Holocene are rare. If we are to resolve how the peopling process occurred, more are needed beyond the three currently available: Anzick1, Montana (~12.8 ka) (3); Kennewick Man/Ancient One, Washington (~9 ka) (10), and USR1, Alaska (~11.5 ka) (1).

### Dataset and method summary

We engaged and sought feedback from Indigenous groups linked to the ancestral individuals analyzed in this study using the recommendations for genomics research with Indigenous communities (11–13). We obtained genome sequences from 15 ancient human remains (Fig. 1A). These include Trail Creek Cave 2, Alaska (radiocarbon dated to ~9 ka, ~0.4× genomic depth of coverage); Big Bar Lake, British Columbia (~5.6 ka, ~1.2×); Spirit Cave, Nevada (~10.7 ka, ~18×); four individuals from Lovelock Cave, Nevada (ranging in age from ~1.95 to 0.6 ka, and ~0.5× to ~18.7×); five individuals from Lagoa Santa, Brazil (~10.4 to ~9.8 ka, ~0.18× to ~15.5×); two individuals from the Punta Santa Ana and Ayayema sites in Patagonian Chile (~7.2 ka and ~5.1 ka, ~1.5× and ~10.6×, respectively); and, an Inca mummy from Mendoza, Argentina (estimated to be ~0.5 ka, ~2.5×) (14) [all  $^{14}\text{C}$  ages are in calibrated years (13, 15)] (Fig. 1, A and B). We also sequenced a ~15.9× genome from a ~19th century Andaman islander, used as a proxy for Australasian ancestry in models involving admixture into Native Americans (2, 6, 13, 16). All DNA extracts were confirmed to contain fragments with characteristic ancient DNA misincorporation patterns and low contamination levels (<3%) (13). The Spirit Cave, Lovelock 2, and Lovelock 3 genomes were generated solely from USER-treated extracts, confirmed to contain characteristic ancient DNA misincorporation patterns prior to treatment (13, 17).

To assess the genetic relationships between these and

other ancient and contemporary human genomes, we compiled a whole-genome comparative dataset of 378 individuals (13). Additionally, we merged these data with a SNP panel of 167 worldwide populations genotyped for 199,285 SNPs, enriched in NA populations whose European and African ancestry components have been identified and masked (2, 4, 13, 18) (Fig. 1A). Of particular interest are the Mixe, a Mesoamerican reference group representing an early internal branch within SNA, prior to the divergence of South Americans (4), and that lacks the Australasian ancestry signal documented among some Amazonian groups (2, 6, 16).

We explored the ancient individuals’ broad genetic affinities initially using model-based clustering (19) and multidimensional scaling (MDS) (Fig. 1, C to F (13)). MDS was applied to both the identity-by-state distance matrix for all individuals (20) and the  $f_3$ -distance matrix over populations included in the SNP array dataset (13, 21, 22). We then tested specific hypotheses by computing error-corrected and genotype-based  $D$ -statistics (13, 21, 23) and fitting admixture graphs (4, 13, 21, 24) (Figs. 2, 3, and 4). Furthermore, we inferred demographic and temporal parameters using the joint site frequency spectrum (SFS) (25, 26) and linkage disequilibrium (27, 28) information (Fig. 5). These efforts enabled us to explore finer scale complex models, using whole-genome data (13). Average depth of coverage of the genomes presented in this study ranges widely, which meant that not all genomes could be used in all analyses, as specified (13).

Our aim is to understand broad patterns in the dispersal, divergence and admixture of people throughout the Americas. Given the highly uneven distribution of genome samples in time and space, our results are expressed as much as possible chronologically from oldest to youngest, and geographically from north to south, to mirror how the peopling of the Americas proceeded.

### Alaskan genome informs on early eastern Beringian populations

While the earliest archaeological evidence for a human presence in eastern Beringia remains disputed, people were present in Alaska by at least 14.4 ka (29). Genomic insights from the USR1 genome indicate that AB (1) remained isolated in interior Alaska until at least the terminal Pleistocene, and was an outgroup to NNA and SNA. It was inferred the NNA/SNA population split occurred outside of eastern Beringia (1, 2). In contrast, recent findings suggest the ancestral population of NNA may have existed north of the continental ice sheets (9).

The Trail Creek Cave genome is from a tooth of a young child recovered from Alaska’s Seward Peninsula (13). This individual clusters adjacent to USR1 in MDS analyses (Fig. 1C) (13) and carries a similar distribution of ancestry components (Fig. 1F) (13). Trail Creek and USR2 (found with and a close relative to USR1), harbor the same mtDNA haplogroup, B2,

but not the derived B2 variant found elsewhere in the Americas (1, 13). Genotype-based *D*-statistics of the form *D*(*Aymara, NA; TrailCreek, Yoruba*) and *D*(*USR1, TrailCreek; NA, Yoruba*) suggest that Trail Creek forms a clade with USR1 that represents an outgroup to other Native Americans (13). This placement was supported by fitting *f*-statistic-based admixture graphs (13, 21).

This procedure used here and for other samples, relies on a “seed graph” that incorporates the formation of ancestral NA and their three basal branches (AB, NNA, SNA) (1, 30, 31). The seed graph includes the following leaves: Yoruba (representing Africans), Mal’ta (Ancient North Eurasians), Andaman (Australasians), Han (East Asians), USR1 (AB), Athabascan (NNA) and Spirit Cave (SNA, see below) (13). We then enumerated all possible extensions of the seed graph where an individual genome, Trail Creek in this case, was added as either a non-admixed or an admixed population (32). We optimized the parameters for each topology using qpGraph (21) and favored the graph producing the best likelihood and the lowest residuals between observed and predicted *f*-statistics. Given that admixed models yield better likelihood scores (because of the additional parameters being optimized), we considered an admixed model to be an improvement compared to its non-admixed counterpart only if the absolute difference between fit scores (log-likelihoods) was >~4.6, corresponding to a *p*-value of ~0.01 in a standard likelihood ratio test (30). In agreement with the exploratory analyses, we found the model in which Trail Creek and USR1 form a clade to be the most likely (Fig. 2A) (13).

These results suggest that USR1 and Trail Creek were members of an AB metapopulation that occupied eastern Beringia and remained isolated from other NA populations during the Late Pleistocene/Early Holocene. Finding two members of the AB population, from sites ~750 km apart, with a similar artifact technology (13), supports the inference the SNA/NNA split occurred south of eastern Beringia (1, 9). The alternative, that NNA/SNA split in Alaska, seems less likely; it would have required several thousand years of strong population structure prior to ~16 ka to differentiate those groups from each other and from AB, as well as a separate SNA presence, which has yet to be found (1). These data indicate that the Athabascans and Inuit, who inhabit Alaska today and are NNA, but with additional Siberian-related ancestry (1, 4, 18, 33), presumably moved north into the region sometime after ~9 ka, the age of the Trail Creek individual (1, 13).

### Rapid dispersal of the SNA population across the Rockies and into South America

The NNA/SNA split is estimated to have taken place ~17.5–14.6 ka (1, 2). Members of the SNA branch ultimately reached southern South America, and based on mtDNA, Y chromosome and genome-wide evidence this likely occurred quickly

(2, 7, 8, 34, 35). This movement gave rise to serial splitting and early population structure, with Mesoamericans being the most deeply divergent group, followed by South Americans east and west of the Andes (4, 36). However, genomic data from Spirit Cave (10.7 ka) and Lagoa Santa (10.4 ka), the oldest sites in this study, show that the SNA dispersal pattern south of the continental ice sheets involved complex admixture events between earlier-established populations.

MDS and ADMIXTURE, as well as a TreeMix tree focused on SNA genomes, reveal that the Spirit Cave and Lagoa Santa individuals were members of the SNA branch (Fig. 1, C and F) (13). Within that branch, Spirit Cave is closest to Anzick1, while Lagoa Santa is closest to southern SNA groups. Interestingly, two of the Lagoa Santa individuals carry the same mtDNA haplogroup (D4h3a) as Anzick1, yet three of the Lagoa Santa individuals harbor the same Y chromosome haplogroup as Spirit Cave (Q-M848) (13). Nonetheless, MDS transformations restricted to SNA (Fig. 1, D and E) (13) together with TreeMix graphs including admixture (13), suggest these North and South American ancient individuals are closely related, regardless of Lagoa Santa’s affinity to present-day South American groups.

We formally tested this scenario by fitting *f*-statistics-based admixture graphs, and found that even though Anzick1, Spirit Cave and Lagoa Santa are separated by ~2000 years and thousands of kilometers, these three individuals can be modeled as a clade to the exclusion of the Mesoamerican Mixe (13). While we did not find evidence rejecting this clade using TreeMix and *D*-statistics (13), further SFS-based modeling indicates that Mixe most likely carry gene flow from an unsampled outgroup, and form a clade with Lagoa Santa. Including nonzero outgroup admixture into Mixe when fitting an *f*-statistics-based admixture graph resulted in a significantly better fit (likelihood ratio test, *P* < 0.05) (Fig. 3, A and B) (13). Hereafter we refer to that outgroup as ‘Unsampled population A’ (UPopA), which is neither AB, NNA or SNA, and which we infer split off from Native Americans ~24.7 ka, ranging from 30–22 ka (95% CI; this large range is a result of the analytical challenge of estimating divergence and admixture times in the absence of UPopA genome data). This age range overlaps with the inferred split of Native Americans from Siberians and East Asians 26.1–23.9 ka (1), and the divergence of USR1 from other NA (23.3–21.2 ka). This temporal overlap, which cannot be fully resolved into a relative sequence with current data, suggests there were multiple splits that took place in Beringia close in time. Depending on how close they ultimately prove to be, they could imply there was moderate structure within Beringia (1, 37), possibly along with indirect gene flow from Siberians, perhaps via other NA populations. Under a model with a pulse-like gene flow, we inferred a probability of ~11% gene flow from UPopA into Mixe ~8.7 ka (95% CI: 0.4–13.9 ka; the wide interval

potentially reflects unmodeled continuous migration) (Fig. 5) (13). Thus, we favor a model where the common ancestor of Anzick1 and Spirit Cave diverged from the common ancestor of Lagoa Santa and Mixe ~14.1 ka (95% CI: 13.2–14.9 ka), perhaps as the latter were moving southward. We infer that the Lagoa Santa population diverged from Mixe shortly thereafter, ~13.9 ka (95% CI: 12.8–14.8 ka) (Fig. 5) (13). The proximity of these estimated divergence times suggests the dispersal process was very rapid on an archaeological time scale, as populations expanded across North America perhaps in a matter of centuries, and then into eastern South America within a millennium or two.

### Australasian ancestry in Early Holocene South America and claims of “Paleoamericans”

Both the Spirit Cave and Lagoa Santa individuals have been identified as “Paleoamericans” (38, 39), connoting a cranial morphology distinct from that of modern Native Americans. Interpretations of this pattern range from it being the result of a separate earlier migration into the Americas, to population continuity and *in situ* differentiation owing to factors such as isolation and drift (13, 40–42). We examined if this morphology might be associated with ancient Australasian genetic ancestry found in present-day Amazonian groups (2, 6). However, no morphometric data are available for present-day peoples with this genetic signal (6), nor has this signal been detected in any ancient skeleton with this morphology (2, 10).

To test for the Australasian genetic signal in Native Americans, we computed *D*-statistics of the form  $D(NA, NA; Eurasian, Yoruba)$ , where NA represents all newly sequenced and reference high-depth Native American genomes (13). In agreement with previous results (6), we find that the Amazonian Suruí share a larger proportion of alleles with Australasian groups (represented by Papuans, Australians and Andaman Islanders), with respect to Mixe (13). Lagoa Santa yielded similar results to those obtained for the Suruí. When compared to Mesoamerican groups (Mixe and Huichol), Lagoa Santa also shares a larger proportion of alleles with Australasian groups but not with other Eurasians (Fig. 4) (13). However, the Australasian signal is not present in the Spirit Cave individual, and we include this distinction in the admixture graph modeling (Figs. 3A and 4A) (13). We inferred less than 3% European contamination in the Lagoa Santa genome (<3%) (13) and show that this finding is robust to potential European contamination by computing “contamination-corrected”  $f_4$ -statistics (Fig. 4B) (13). The fact that the Australasian genomic signature was present in Brazil 10.4 ka, but absent in all genomes tested to date as old or older and further north, presents a challenge in accounting for its presence in Lagoa Santa.

Importantly, all sequenced “Paleoamericans” (including Kennewick/Ancient One) (2, 10), are genetically closer to

contemporary Native Americans, than to any other ancient or contemporary group sequenced to date.

### Multiple dispersals into South America

Genome-wide data from contemporary populations suggested a single expansion wave into South America with little gene flow between groups (4), but see (36). In contrast, analysis of later Holocene genomes suggests South Americans derived from one or more admixture events between two ancestral Native American groups, possibly via multiple movements into South America (7).

To test these competing scenarios, we performed an exhaustive admixture graph search, as described above. Here we fitted a “seed graph” involving Yoruba, Mal’ta, Andaman, Han, Anzick1, Spirit Cave, Lagoa Santa and Mixe (present-day Mesoamericans), and tested all possible “non-admixed” and “admixed” models for SNA: the Mesoamerican Maya and Yukpa of Venezuela; groups east (Suruí, Karitiana, Piapoco, Chane) and west of the spine of the Andes (Aymara, Quechua); six ancient Patagonians [Ayayema, Punta Santa Ana and four individuals from (43)], the ancient Taino (44) and the Aconcagua Inca mummy (14) (Fig. 1B) (13). This analysis indicates most present-day South American populations do not form a clade with Lagoa Santa, but instead derive from a mixture of Lagoa Santa and Mesoamerican-related ancestries (Fig. 3A) (13). We confirmed these results by computing standard and error-corrected *D*-statistics of the form  $D(LagoaSanta, SNA; Mixe, Yoruba)$  and  $D(Mixe, SNA; LagoaSanta, Yoruba)$  (Fig. 3B) (13). For most groups, these statistics are inconsistent with a simple tree and indicate multiple dispersals into South America.

The ~5.1 ka Patagonian Ayayema genome is an exception; it forms a clade with the Lagoa Santa population. This suggests the arrival of the Mesoamerican-related ancestry occurred post-5.1 ka, and/or that it did not reach the remote region inhabited by the Ayayema individual’s ancestors (Fig. 3C) (13). This result is qualitatively mirrored by the 7.2 ka Punta Santa Ana individual (both cluster with present-day Patagonians, and form a clade with Lagoa Santa). However, the low coverage of Punta Santa Ana may reduce our power to detect possible Mesoamerican-related admixture (Fig. 3C) (13).

We further explored the fit of the model (Fig. 3A) for each South American group by fixing the Australasian contribution into Lagoa Santa and the Mesoamerican contribution (Fig. 3, D and E) into the test SNA population, across a range of values (13). While an Australasian contribution <1% and >~6% results in a significant decrease in likelihood (likelihood ratio test,  $P < 0.05$ ), the Mesoamerican contribution has a wider range of plausible values (Fig. 3E) (13). Yet, modelling each SNA group with little to no Mesoamerican-related admixture consistently yields significantly lower fit scores ( $P < 0.05$ ) (13), except for the Ayayema individual (Fig. 3D) (13).

The Australasian contribution into Lagoa Santa is consistently nonzero when modeling South Americans, although we did not observe in every case a significant improvement when modeling Australasian admixture into SNA groups through Lagoa Santa (13). This result suggests this ancestry may have been widespread among early South Americans. Although we are unable to estimate the Lagoa Santa-related admixture proportion for these groups with confidence, we observe a general trend for populations east of the Andes (e.g., Suruí) to bear more of this ancestry than Andean groups (e.g., Aymara) (Fig. 3F) (13). A possible explanation for this difference is that on the western side of the Andes there was greater Mesoamerican-related admixture.

Finally, we explore the demographic history of present day South Americans using both joint SFS (*momi2*) (25, 26) and linkage disequilibrium information [*smc++* (27) and *diCal2* (28)]. We seek to understand these groups' relationships to Lagoa Santa – which also provides an indirect means of assessing the impact of admixture on the Australasian signature. For the SFS analysis, we selected the Karitiana, the only SNA population for which there are sufficient unadmixed publicly-available genomes ( $n = 5$ ); for the *diCal* analysis, we used the Karitiana, Aymara ( $n = 1$ ) and Suruí ( $n = 2$ ) (13). From SFS analysis we infer that the ancestors of Lagoa Santa and Karitiana diverged from each other ~12.9 ka (95% CI: 10.4–14.0 ka). Subsequently, the latter received gene flow from a Mesoamerican-related population, which already carried admixture from the outgroup UPopA (Fig. 5) (13). Assuming pulse-like migration, this points to recent gene flow (~35%) from the Mesoamerican-related group into Karitiana (Fig. 5), possibly suggesting ongoing admixture over an extended period. Indeed, when we allowed for two pulses, we inferred substantial gene flow in both the recent and distant past (13). The *diCal2* results are consistent for the Karitiana, Aymara, and Suruí showing their demographic histories involved a mixture between a Lagoa Santa-related and a Mixe-related source (13).

Overall, our findings suggest that soon after arrival, South Americans diverged along multiple geographic paths (36). That process was further complicated by the arrival of a second independent migration and gene flow in Middle to Late Holocene times. Later admixture potentially reduced the Australasian signature that might have been carried by earlier inhabitants.

#### **Long-term population continuity in the North American Great Basin and the Numic Expansion**

Mesoamerican-related expansion possibly had a bearing on a later, unresolved pattern seen in North America. In the western Great Basin of North America paleoenvironmental evidence indicates there was decreased effective precipitation and increased aridity during the Middle Holocene, which led to a human population decline (45, 46). By ~5 ka regional

populations were rebounding, but whether these were descendants of the previous inhabitants is not known. Unclear also is the relationship between those later Holocene groups and the people present in the region at the time of European Contact and today. There is linguistic evidence that ancestors of Numic speakers presently inhabiting the region today arrived recently, perhaps ~1 ka. There is also archaeological evidence of changes in material culture around that time, though how those relate to the linguistic turnover is uncertain. Nor is it known if these changes are related to population admixture or replacement. Patterns and changes in language, material culture and genetics need not be congruent or causally linked (47). Thus, the question of what has been called the Numic Expansion has been highly debated (46, 48); we address the population aspect by comparing genomes from Spirit Cave and Lovelock Cave (Fig. 1A) (13).

MDS and ADMIXTURE analyses, as well as *D*-statistics of the form  $D(SpiritCave, Lovelock2/3; NA, Yoruba)$  suggest that despite the ~9,000 years separating the Spirit Cave and Lovelock individuals, they form a clade with respect to other NA (Fig. 1, C to F) (13). We tested that topology through the same admixture graph search implemented for SNA (13). We were not able to reject the model without Mesoamerican-related admixture for Lovelock2 (~1.9 ka). However, the ~0.7 ka Lovelock3 individual received Mesoamerican-admixture, from a group that was likely not present in the region just ~1.2 ka earlier in Lovelock2. Because we do not know the language(s) that may have been involved, we cannot securely attribute this admixture to arriving Numic speakers [the Mixe, whom we use as a proxy for Mesoamerican ancestry, fall in a separate language family from Numic (49)]. Importantly, we also observe genetic continuity, suggesting that there was not a complete population replacement.

Present-day Pima from northern Mexico can also best be modeled as a Mesoamerican-related mixture. However, the Pima require admixture from a branch splitting above the Mixe-Spirit Cave divergence, likely an NNA population (13). We cannot specify a particular source population. These patterns indicate that complex population movements and mixture occurred after the initial settlements of the Great Basin and Southwest from both the north and south.

#### **Long-term complex population history in the Pacific Northwest**

Pacific Northwest groups had a late Pleistocene demographic history argued to be distinct from early SNA groups (1, 2, 9, 18, 33). To explore the population history and the relationship of regional populations to NNA and SNA, we assessed the genetic affinities between the 5.6 ka Big Bar Lake individual from the Fraser Plateau of central British Columbia, and other NA. Given their relative geographic proximity, we included the 939, 302, and 443 individuals from coastal British Columbia (9), and the Kennewick Man/Ancient One

(10). Since these genomes have been deemed representatives of NNA, we also included genomic data from Southwestern Ontario (ASO) individuals, who are closely related to Algonquin (NNA) populations (7).

These ancient North American individuals clustered separately from SNA populations in both MDS transformations, and their ancestry component distribution closely resembles that of NNA populations (Fig. 1, C and F) (13). However, we observed genetic differentiation between these individuals and other North American populations. While the coastal British Columbia ancient individuals clustered together with present-day Athabaskan and Tsimshian speakers from the region, ASO and Kennewick were placed in an intermediate position between NNA and SNA. Although the Big Bar individual was placed close to NNA populations not carrying recent Siberian admixture (Fig. 1, C and F) (13), *D*-statistics of the form  $D(Aymara, NA; BigBar, Yoruba)$  and  $D(USR1, NA; BigBar, Yoruba)$  suggest Big Bar represents a previously undetected outgroup to non-AB Native Americans, one that diverged prior to the NNA/SNA split (13).

In order to describe the genetic ancestry of these individuals, we used the admixture graph search strategy (13). In agreement with previous results, the ancient coastal British Columbia individuals are best modeled as a clade to Athabascans, who bear Siberian-related admixture (Fig. 2C). However, the best-fitting model suggests the Big Bar individual represents a population that split prior to the NNA-SNA divergence, but after AB and without Siberian admixture (Fig. 2B) (13). Finally, in accordance with their placement in both MDS transformations, we observed that Kennewick and ASO individuals are best modeled as deriving a fraction of their ancestry from an SNA-related source, represented by Spirit Cave in this case (Fig. 2, D and E) (13). We confirm this through error-corrected *D*-statistics (Fig. 2F) (13), suggesting gene flow between ASO and an SNA group that diverged after the split of Anzick1, and that did not bear recent Mesoamerican-related ancestry.

Thus, the broader population history in this region was evidently marked by admixture between the NNA and SNA branches which most likely gave rise to the ancestors of Kennewick and ASO, and by isolation between groups in coastal (e.g., the 939 individual) and interior (Big Bar) British Columbia.

## Discussion

The genomes described here do not undermine the previously established tree of AB splitting from ancestral Native Americans, followed by the basal NNA/SNA split south of eastern Beringia. However, they show it is at best a rough outline of the process. We now find that once south of eastern Beringia, Native Americans radiated rapidly and gave rise to multiple populations, some of which are visible in the genetic record only as unsampled populations, and who at different

times expanded to different portions of the continent, though not as extensively as the initial peopling (Fig. 6).

Rapid movement from North to South America is evident genetically (Fig. 6, A and B), and had been anticipated by the ‘archaeologically-instantaneous’ appearance of sites throughout the hemisphere just after 13 ka (50, 51). It suggests the mechanism of movement was not simply gradual population growth and incremental geographic expansion, but rather was likely more akin to leap-frogging across large portions of the diverse intervening landscape (52). If this result holds, it predicts that additional terminal Pleistocene samples will fit on a star-like pattern as observed here.

That the early population evidently spread widely, rapidly, but if somewhat unevenly across the Americas, in turn suggests their access to large portions of the hemisphere was essentially unrestricted (52). Yet, there are hints in the genetic record of early unsampled populations (6) (Fig. 5), and the material culture associated with that rapid spread (Clovis and later) is distinct from and post-dates the earliest secure archaeological presence in the Americas at 14.6 ka (53). How these early groups are related, particularly ones with excess Australasian ancestry, and their degree of structure, remains largely unknown.

The Australasian signal is not present in USR1 or Spirit Cave, but only appears in Lagoa Santa. None of these individuals has UPopA/Mesoamerican-related admixture, which apparently damped the Australasian signature in South American groups, such as the Karitiana (Figs. 4 and 5). These findings suggest the Australasian signal, possibly present in a structured ancestral NA population (16), was absent in NA prior to the Spirit Cave/Lagoa Santa split. Groups carrying this signal were either already present in South America when the ancestors of Lagoa Santa reached the region, or Australasian-related groups arrived later but before 10.4 ka (the Lagoa Santa  $^{14}\text{C}$  age). That this signal has not been previously documented in North America implies that an earlier group possessing it had disappeared, or a later-arriving group passed through North America without leaving any genetic trace (Fig. 6, A and C). If such a signal is ultimately detected in North America it could help determine when groups bearing Australasian ancestry arrived, relative to the divergence of SNA groups.

Although we detect the Australasian signal in one of the Lagoa Santa individuals identified as a “Paleoamerican,” it is absent in other “Paleoamericans” (2, 10), including Spirit Cave with its strong genetic affinities to Lagoa Santa. This indicates the “Paleoamerican” cranial form is not associated with the Australasian genetic signal, as previously suggested (6), or any other specific NA clade (2). The cause of this cranial form, if it is representative of broader population patterns, evidently did not result from separate ancestry, but likely multiple factors, including isolation and drift and non-

stochastic mechanisms (2, 10, 13, 54).

The attenuating effects of distance, compounded (in places) by geographic barriers, led to cultural drift and regional adaptations, even early in the peopling process (52, 55). It was previously surmised that Clovis (Anzick1) and Western Stemmed (Spirit Cave) technologies (46, 56) represented “genetically divergent, founding groups” (57). It appears instead that the divergence is principally cultural, between genetically close populations living on opposite sides of the Rocky Mountains. This result affirms the point that archaeological, anatomical, and genetic records are not necessarily congruent (47).

That one of the principal isolating mechanisms was likely geographic helps explain the long-term population continuity in the Great Basin. Continuity occurred despite fluctuating human population densities and the cultural and linguistic changes that occurred over a 9,000 year span (46). In the Pacific NW, geographic barriers were less formidable, but we surmise that the region’s natural richness and diversity may have led groups to inhabit different environmental niches, which resulted in the emergence of social boundaries that maintained populations separation. In this region’s long history, we find evidence that groups on the coast (e.g., 939) and their contemporaries in the interior (Big Bar), were as genetically distinct as are present-day groups (18) (Fig. 6A). It is not known how or whether such differences map on to the region’s rich linguistic complexity and material culture differences (13, 58). Previous research on mtDNA and Y sequences hypothesized a shared origin for Pacific Northwest populations, followed by divergence due to isolation and drift (18). That Big Bar represents a previously unseen, isolated population supports the latter, but implies that the initial peopling of the region was complex and structured.

We also find evidence of a later Mesoamerican-admixture, which while geographically extensive, was not associated with a “wave” throughout the Americas, nor did it inevitably lead to replacement. Rather, it appears to mark the movement north and south (Fig. 6C), of what may have been relatively small groups that did not necessarily swamp local populations genetically or culturally, as illustrated by admixture in the Lovelock3 individual. Regardless of whether this marks the “Numic Expansion,” it was associated with evidence of cultural continuity as well as change; it was not an instance of population replacement. How or whether this Mesoamerican-related expansion is expressed culturally in South America is not known.

The genomes reported here fill gaps in our temporal and spatial coverage, and are valuable anchor points that reveal that the human population history of the Americas. As has long been expected (52), and is characteristic of human population histories around the world (59), the peopling process was marked by complex local and long-range demographic

processes over time. The peopling of the Americas will likely prove more complicated still. As we have found, there was a previously unknown population in the Americas (UPopA), as well as one that harbored an Australasian signal in the Late Pleistocene and reached South America, yet left no apparent traces in North America. In addition, all of our evidence of the peopling process is from archaeologically known groups: Clovis (Anzick) and later populations. Yet, there is archaeological evidence of an earlier, pre-Clovis presence in the Americas, one for which we have yet to recover any ancient DNA. How these various population threads may ultimately come together, and how these populations were related to Native Americans past and present, remains to be resolved.

## Materials and methods

### Laboratory procedures

Ancient DNA work was performed in dedicated clean laboratory facilities at the Centre for GeoGenetics, Natural History Museum, University of Copenhagen. Extraction, treatment and library build protocols followed for each sample are detailed in (13). Sequencing was carried out in Illumina HiSeq instruments.

### Data processing

Sequencing reads were trimmed for Illumina adaptors using AdapterRemoval (60) and mapped to the human reference genome build 37 using BWA v.0.6.2-r126 (61) with disabled seeding (-l parameter) (62). Reads with mapping quality lower than 30 were discarded, PCR duplicates were identified using MarkDuplicates (63) and local realignment was carried out using GATK (64). Genotype calls for high coverage samples were generated using SAMtools mpileup (65) and filtered following (2). Called genotypes were phased with impute2 (66, 67) using the 1,000 genomes phased variant panel (phase 3) as a reference and the HapMap recombination rates. The final callset was masked using a 35-mer “snpability” mask with a stringency of 0.5 (68), and the strict accessible regions from the 1000 Genomes Project (69).

### Ancient DNA data authentication

We examined the fragment length distributions and the base substitution patterns using bamdamage (20). We estimated mtDNA, X-chromosome and nuclear contamination using contamMix (70), ANGSD (71) and DICE (72), respectively.

### Population structure analyses

We investigated the broad relationships between ancient and present-day genomes using model-based clustering as implemented in ADMIXTURE (19) and multidimensional scaling applied to the identity-by-state (20) and  $f_3$ -distance matrices (21, 22).

### D-statistics

We computed  $D$ -statistics to formally test hypotheses of treeness and gene-flow. Genotype-based  $D$ -statistics were computed as detailed in (21), and error-corrected  $D$ -statistics

were computed following (23). For both approaches, standard errors were estimated through a weighted block jackknife approach over 5-Mb blocks.

### Admixture graph fitting

We used qpGraph to fit *f*-statistics-based admixture graphs (4, 21). We implemented an exhaustive admixture graph search where we considered a “seed graph” onto which a “test population” was added either as a “non-admixed” or an “admixed” group in every possible position. Extensions of the “seed graph” were enumerated using the admixturegraph R package (32). We evaluated each topology based on its fit score, the *Z*-score of the worst residual between the observed and predicted *D*-statistics, and the presence of zero-length internal edges, and carried out likelihood-ratio tests following (30). For all tests, we only considered transversion polymorphisms.

### Demographic inference

We estimated marginal population sizes over time for different Native American groups using SMC++ (27). We then used momi2 (25, 26) to infer demographic parameters for a number of models, using the joint site frequency spectrum. Confidence intervals were obtained through a nonparametric bootstrap procedure. We confirmed the SFS-based inference by using diCal2 (28), which relies on linkage disequilibrium information, to infer key demographic parameters relating pairs of Native American populations. A detailed description of laboratory and analytical methods is provided in (13).

### REFERENCES AND NOTES

1. J. V. Moreno-Mayar, B. A. Potter, L. Vinner, M. Steinrücken, S. Rasmussen, J. Terhorst, J. A. Kamm, A. Albrechtsen, A.-S. Malaspina, M. Sikora, J. D. Reuther, J. D. Irish, R. S. Malhi, L. Orlando, Y. S. Song, R. Nielsen, D. J. Meltzer, E. Willerslev, Terminal Pleistocene Alaskan genome reveals first founding population of Native Americans. *Nature* **553**, 203–207 (2018). doi:10.1038/nature25173 Medline
2. M. Raghavan, M. Steinrücken, K. Harris, S. Schiffels, S. Rasmussen, M. DeGiorgio, A. Albrechtsen, C. Valdiosera, M. C. Ávila-Arcos, A.-S. Malaspina, A. Eriksson, I. Moltke, M. Metspalu, J. R. Homburger, J. Wall, O. E. Cornejo, J. V. Moreno-Mayar, T. S. Korneliussen, T. Pierre, M. Rasmussen, P. F. Campos, P. de Barros Damgaard, M. E. Allentoft, J. Lindo, E. Metspalu, R. Rodríguez-Varela, J. Mansilla, C. Henrickson, A. Seguin-Orlando, H. Malmström, T. Stafford Jr., S. S. Shringarpure, A. Moreno-Estrada, M. Karmin, K. Tambets, A. Bergström, Y. Xue, V. Warmuth, A. D. Friend, J. Singarayer, P. Valdes, F. Balloux, I. Leboreiro, J. L. Vera, H. Rangel-Villalobos, D. Pettener, D. Luiselli, L. G. Davis, E. Heyer, C. P. E. Zollikofer, M. S. Ponce de León, C. I. Smith, V. Grimes, K.-A. Pike, M. Deal, B. T. Fuller, B. Arriaza, V. Standen, M. F. Luz, F. Ricaut, N. Guidon, L. Osipova, M. I. Voevodka, O. L. Posukh, O. Balanovsky, M. Lavryashina, Y. Bogunov, E. Khusnutdinova, M. Gubina, E. Balanovska, S. Fedorova, S. Litvinov, B. Malyarchuk, M. Derenko, M. J. Mosher, D. Archer, J. Cybulski, B. Petzelt, J. Mitchell, R. Worl, P. J. Norman, P. Parham, B. M. Kemp, T. Kivisild, C. Tyler-Smith, M. S. Sandhu, M. Crawford, R. Villems, D. G. Smith, M. R. Waters, T. Goebel, J. R. Johnson, R. S. Malhi, M. Jakobsson, D. J. Meltzer, A. Manica, R. Durbin, C. D. Bustamante, Y. S. Song, R. Nielsen, E. Willerslev, Genomic evidence for the Pleistocene and recent population history of Native Americans. *Science* **349**, aab3884–aab3884 (2015). doi:10.1126/science.aab3884 Medline
3. M. Rasmussen, S. L. Anzick, M. R. Waters, P. Skoglund, M. DeGiorgio, T. W. Stafford Jr., S. Rasmussen, I. Moltke, A. Albrechtsen, S. M. Doyle, G. D. Poznik, V. Guðmundsdóttir, R. Yadav, A.-S. Malaspina, S. S. White 5th, M. E. Allentoft, O. E. Cornejo, K. Tambets, A. Eriksson, P. D. Heintzman, M. Karmin, T. S. Korneliussen, D. J. Meltzer, T. L. Pierre, J. Stenderup, L. Saag, V. M. Warmuth, M. C. Lopes, R. S. Malhi, S. Brunak, T. Sicheritz-Ponten, I. Barnes, M. Collins, L. Orlando, F. Balloux, A. Manica, R. Gupta, M. Metspalu, C. D. Bustamante, M. Jakobsson, R. Nielsen, E. Willerslev, The genome of a Late Pleistocene human from a Clovis burial site in western Montana. *Nature* **506**, 225–229 (2014). doi:10.1038/nature13025 Medline
4. D. Reich, N. Patterson, D. Campbell, A. Tandon, S. Mazieres, N. Ray, M. V. Parra, W. Rojas, C. Duque, N. Mesa, L. F. García, O. Triana, S. Blair, A. Maestre, J. C. Dib, C. M. Bravi, G. Bailliet, D. Corach, T. Hünemeier, M. C. Bortolini, F. M. Salzano, M. L. Petzl-Erler, V. Acuña-Alonso, C. Aguilar-Salinas, S. Canizales-Quinteros, T. Tusié-Luna, L. Riba, M. Rodríguez-Cruz, M. Lopez-Alarcón, R. Coral-Vazquez, T. Canto-Cetina, I. Silva-Zolezzi, J. C. Fernández-López, A. V. Contreras, G. Jimenez-Sánchez, M. J. Gómez-Vázquez, J. Molina, A. Carracedo, A. Salas, C. Gallo, G. Poletti, D. B. Witonsky, G. Alkorta-Aranburu, R. I. Sukernik, L. Osipova, S. A. Fedorova, R. Vasquez, M. Villena, C. Moreau, R. Barrantes, D. Pauls, L. Excoffier, G. Bedoya, F. Rothhammer, J.-M. Dugoujon, G. Larrouy, W. Klitz, D. Labuda, J. Kidd, K. Kidd, A. Di Renzo, N. B. Freimer, A. L. Price, A. Ruiz-Linares, Reconstructing Native American population history. *Nature* **488**, 370–374 (2012). doi:10.1038/nature1258 Medline
5. W. A. Neves, M. Hubbe, Cranial morphology of early Americans from Lagoa Santa, Brazil: Implications for the settlement of the New World. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 18309–18314 (2005). doi:10.1073/pnas.0507185102 Medline
6. P. Skoglund, S. Mallick, M. C. Bortolini, N. Chennagiri, T. Hünemeier, M. L. Petzl-Erler, F. M. Salzano, N. Patterson, D. Reich, Genetic evidence for two founding populations of the Americas. *Nature* **525**, 104–108 (2015). Medline
7. C. L. Scheib, H. Li, T. Desai, V. Link, C. Kendall, G. Dewar, P. W. Griffith, A. Morseburg, J. R. Johnson, A. Potter, S. L. Kerr, P. Endicott, J. Lindo, M. Haber, Y. Xue, C. Tyler-Smith, M. S. Sandhu, J. G. Lorenz, T. D. Randall, Z. Faltskova, L. Pagani, P. Danecek, T. C. O’Connell, P. Martz, A. S. Boraas, B. F. Byrd, A. Leventhal, R. Cambra, R. Williamson, L. Lesage, B. Holguin, E. Ygnacio-De Soto, J. Rosas, M. Metspalu, J. T. Stock, A. Manica, A. Scally, D. Wegmann, R. S. Malhi, T. Kivisild, Ancient human parallel lineages within North America contributed to a coastal expansion. *Science* **360**, 1024–1027 (2018). doi:10.1126/science.aar6851 Medline
8. B. Llamas, L. Fehren-Schmitz, G. Valverde, J. Soubrier, S. Mallick, N. Rohland, S. Nordenfelt, C. Valdiosera, S. M. Richards, A. Rohrlach, M. I. B. Romero, I. F. Espinoza, E. T. Cagigao, L. W. Jiménez, K. Makowski, I. S. L. Reyna, J. M. Lory, J. A. B. Torrez, M. A. Rivera, R. L. Burger, M. C. Ceruti, J. Reinhard, R. S. Wells, G. Politis, C. M. Santoro, V. G. Standen, C. Smith, D. Reich, S. Y. W. Ho, A. Cooper, W. Haak, Ancient mitochondrial DNA provides high-resolution time scale of the peopling of the Americas. *Sci. Adv.* **2**, e1501385–e1501385 (2016). doi:10.1126/sciadv.1501385 Medline
9. J. Lindo, A. Achilli, U. A. Perego, D. Archer, C. Valdiosera, B. Petzelt, J. Mitchell, R. Worl, E. J. Dixon, T. E. Fifeield, M. Rasmussen, E. Willerslev, J. S. Cybulski, B. M. Kemp, M. DeGiorgio, R. S. Malhi, Ancient individuals from the North American Northwest Coast reveal 10,000 years of regional genetic continuity. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 4093–4098 (2017). doi:10.1073/pnas.1620410114 Medline
10. M. Rasmussen, M. Sikora, A. Albrechtsen, T. S. Korneliussen, J. V. Moreno-Mayar, G. D. Poznik, C. P. E. Zollikofer, M. P. de León, M. E. Allentoft, I. Moltke, H. Jónsson, C. Valdiosera, R. S. Malhi, L. Orlando, C. D. Bustamante, T. W. Stafford Jr., D. J. Meltzer, R. Nielsen, E. Willerslev, The ancestry and affiliations of Kennewick Man. *Nature* **523**, 455–458 (2015). Medline
11. J. Bardill, A. C. Bader, N. A. Garrison, D. A. Bolnick, J. A. Raff, A. Walker, R. S. Malhi; Summer internship for Indigenous peoples in Genomics (SING) Consortium, Advancing the ethics of paleogenomics. *Science* **360**, 384–385 (2018). doi:10.1126/science.aaq1131 Medline
12. K. G. Claw, M. Z. Anderson, R. L. Begay, K. S. Tsosie, K. Fox, N. A. Garrison; Summer internship for Indigenous peoples in Genomics (SING) Consortium, A framework for enhancing ethical genomic research with Indigenous communities. *Nat. Commun.* **9**, 2957 (2018). doi:10.1038/s41467-018-05188-3 Medline
13. See supplementary materials.
14. A. Gómez-Carballa, L. Catelli, J. Pardo-Seco, F. Martinón-Torres, L. Roewer, C. Vullo, A. Salas, The complete mitogenome of a 500-year-old Inca child mummy. *Sci. Rep.* **5**, 16462 (2015). doi:10.1038/srep16462 Medline
15. P. J. Reimer, E. Bard, A. Bayliss, J. W. Beck, P. G. Blackwell, C. B. Ramsey, C. E. Buck, H. Cheng, R. L. Edwards, M. Friedrich, P. M. Grootes, T. P. Guilderson, H. Haflidason, I. Hajdas, C. Hatté, T. J. Heaton, D. L. Hoffmann, A. G. Hogg, K. A.

- Hughen, K. F. Kaiser, B. Kromer, S. W. Manning, M. Niu, R. W. Reimer, D. A. Richards, E. M. Scott, J. R. Southon, R. A. Staff, C. S. M. Turney, J. van der Plicht, IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* **55**, 1869–1887 (2013). doi:10.2458/azu\_js\_rc.55.16947
16. P. Skoglund, D. Reich, A genomic view of the peopling of the Americas. *Curr. Opin. Genet. Dev.* **41**, 27–35 (2016). doi:10.1016/j.gde.2016.06.016 Medline
17. A. W. Briggs, U. Stenzel, M. Meyer, J. Krause, M. Kircher, S. Pääbo, Removal of deaminated cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res.* **38**, e87–e87 (2010). doi:10.1093/nar/gkp1163 Medline
18. P. Verdu, T. J. Pemberton, R. Laurent, B. M. Kemp, A. Gonzalez-Oliver, C. Gorodezky, C. E. Hughes, M. R. Shattuck, B. Petzelt, J. Mitchell, H. Harry, T. William, R. Worl, J. S. Cybulski, N. A. Rosenberg, R. S. Malhi, Patterns of admixture and population structure in native populations of Northwest North America. *PLOS Genet.* **10**, e1004530 (2014). doi:10.1371/journal.pgen.1004530 Medline
19. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009). doi:10.1101/gr.094052.109 Medline
20. A.-S. Malaspina, O. Tange, J. V. Moreno-Mayar, M. Rasmussen, M. DeGiorgio, Y. Wang, C. E. Valdiosera, G. Politis, E. Willerslev, R. Nielsen, bammds: A tool for assessing the ancestry of low-depth whole-genome data using multidimensional scaling (MDS). *Bioinformatics* **30**, 2962–2964 (2014). doi:10.1093/bioinformatics/btu410 Medline
21. N. Patterson, P. Moorjani, Y. Luo, S. Mallick, N. Rohland, Y. Zhan, T. Genschoreck, T. Webster, D. Reich, Ancient admixture in human history. *Genetics* **192**, 1065–1093 (2012). doi:10.1534/genetics.112.145037 Medline
22. Q. Fu, C. Posth, M. Hajdinjak, M. Petr, S. Mallick, D. Fernandes, A. Furtwängler, W. Haak, M. Meyer, A. Mittnik, B. Nickel, A. Peltzer, N. Rohland, V. Slon, S. Talamo, I. Lazaridis, M. Lipson, I. Mathieson, S. Schiffels, P. Skoglund, A. P. Derevianko, N. Drozdov, V. Slavinsky, A. Tsybarkov, R. G. Cremonesi, F. Mallegni, B. Gély, E. Vacca, M. R. G. Morales, L. G. Straus, C. Neugebauer-Maresch, M. Teschler-Nicola, S. Constantin, O. T. Moldovan, S. Benazzi, M. Peresani, D. Coppola, M. Lari, S. Ricci, A. Ronchitelli, F. Valentini, C. Thevenet, K. Wehrberger, D. Grigorescu, H. Rougier, I. Crevecoeur, D. Flas, P. Sernal, M. A. Mannino, C. Cupillard, H. Bocherens, N. J. Conard, K. Harvati, V. Moiseyev, D. G. Drucker, J. Svoboda, M. P. Richards, D. Caramelli, R. Pinhasi, J. Kelso, N. Patterson, J. Krause, S. Pääbo, D. Reich, The genetic history of Ice Age Europe. *Nature* **534**, 200–205 (2016). doi:10.1038/nature17993 Medline
23. S. Soraggi, C. Wiuf, A. Albrechtsen, Powerful inference with the D-Statistic on low-coverage whole-genome data. *G3 (Bethesda)* **8**, 551–566 (2018). doi:10.1534/g3.117.300192 Medline
24. J. K. Pickrell, J. K. Pritchard, Inference of population splits and mixtures from genome-wide allele frequency data. *PLOS Genet.* **8**, e1002967 (2012). doi:10.1371/journal.pgen.1002967 Medline
25. J. A. Kamm, J. Terhorst, Y. S. Song, Efficient computation of the joint sample frequency spectra for multiple populations. *J. Comput. Graph. Stat.* **26**, 182–194 (2017). doi:10.1080/10618600.2016.1159212 Medline
26. J. A. Kamm, J. Terhorst, R. Durbin, Y. S. Song, Efficiently inferring the demographic history of many populations with allele count data. *bioRxiv* 287268 [Preprint]. 23 March 2018. https://doi.org/10.1101/287268.
27. J. Terhorst, J. A. Kamm, Y. S. Song, Robust and scalable inference of population history from hundreds of unphased whole genomes. *Nat. Genet.* **49**, 303–309 (2017). doi:10.1038/ng.3748 Medline
28. M. Steinrücken, J. A. Kamm, Y. S. Song, Inference of complex population histories using whole-genome sequences from multiple populations. *bioRxiv* 026591 [Preprint]. 16 September 2015. https://doi.org/10.1101/026591.
29. C. E. Holmes, in *From the Yenisei to the Yukon: Interpreting Lithic Assemblage Variability in Late Pleistocene/Early Holocene Beringia*, T. Goebel, I. Buvit, Eds. (Peopling of the Americas, Texas A&M Univ., ed. 1, 2011).
30. M. Lipson, D. Reich, A working model of the deep relationships of diverse modern human genetic lineages outside of Africa. *Mol. Biol. Evol.* **34**, 889–902 (2017). Medline
31. M. Raghavan, P. Skoglund, K. E. Graf, M. Metspalu, A. Albrechtsen, I. Moltke, S. Rasmussen, T. W. Stafford Jr., L. Orlando, E. Metspalu, M. Karmin, K. Tambets, S. Roots, R. Mägi, P. F. Campos, E. Balanovska, O. Balanovsky, E. Khusnutdinova, S. Litvinov, L. P. Osipova, S. A. Fedorova, M. I. Voevodova, M. DeGiorgio, T. Sicheritz-Ponten, S. Brunak, S. Demeshchenko, T. Kivisild, R. Villems, R. Nielsen, M. Jakobsson, E. Willerslev, Upper Palaeolithic Siberian genome reveals dual ancestry of Native Americans. *Nature* **505**, 87–91 (2014). doi:10.1038/nature12736 Medline
32. K. Leppälä, S. V. Nielsen, T. Mailund, admixturegraph: An R package for admixture graph manipulation and fitting. *Bioinformatics* **33**, 1738–1740 (2017). doi:10.1093/bioinformatics/btx048 Medline
33. P. Flegontov et al., Na-Dene populations descend from the Paleo-Eskimo migration into America. *bioRxiv* 074476 [Preprint]. 13 September 2016. https://doi.org/10.1101/074476
34. E. Tamm, T. Kivisild, M. Reidla, M. Metspalu, D. G. Smith, C. J. Mulligan, C. M. Bravi, O. Rickards, C. Martinez-Labarga, E. K. Khusnutdinova, S. A. Fedorova, M. V. Golubenko, V. A. Stepanov, M. A. Zubova, S. I. Zhadanov, L. P. Ossipova, L. Damba, M. I. Voevodova, J. E. Dipierrri, R. Villems, R. S. Malhi, Beringian standstill and spread of Native American founders. *PLOS ONE* **2**, e829 (2007). doi:10.1371/journal.pone.0000829 Medline
35. M. C. Bortolini, R. González-José, S. L. Bonatto, F. R. Santos, Reconciling pre-Columbian settlement hypotheses requires integrative, multidisciplinary, and model-bound approaches. *Proc. Natl. Acad. Sci. U.S.A.* **111**, E213–E214 (2014). doi:10.1073/pnas.1321197111 Medline
36. D. N. Harris, W. Song, A. C. Shetty, K. S. Levano, O. Cáceres, C. Padilla, V. Borda, D. Tarazona, O. Trujillo, C. Sanchez, M. D. Kessler, M. Galarza, S. Capristano, H. Montejo, P. O. Flores-Villanueva, E. Tarazona-Santos, T. D. O'Connor, H. Guio, Evolutionary genomic dynamics of Peruvians before, during, and after the Inca Empire. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E6526–E6535 (2018). doi:10.1073/pnas.1720798115 Medline
37. D. A. Bolnick, J. A. Raff, L. C. Springs, A. W. Reynolds, A. T. Miró-Herrans, Native American Genomics and Population Histories. *Annu. Rev. Anthropol.* **45**, 319–340 (2016). doi:10.1146/annurev-anthro-102215-100036
38. W. A. Neves, M. Hubbe, L. B. Piló, Early Holocene human skeletal remains from Sumidouro Cave, Lagoa Santa, Brazil: History of discoveries, geological and chronological context, and comparative cranial morphology. *J. Hum. Evol.* **52**, 16–30 (2007). doi:10.1016/j.jhevol.2006.07.012 Medline
39. R. L. Jantz, D. W. Owsley, Variation among early North American crania. *Am. J. Phys. Anthropol.* **114**, 146–155 (2001). doi:10.1002/1096-8644(200102)114:2<146::AID-AJPA1014>3.0.CO;2-6 Medline
40. N. von Cramon-Taubadel, A. Strauss, M. Hubbe, Evolutionary population history of early Paleoamerican cranial morphology. *Sci. Adv.* **3**, e1602289 (2017). doi:10.1126/sciadv.1602289 Medline
41. W. A. Neves, J. F. Powell, A. Prous, E. G. Ozolins, M. Blum, Lapa Vermelha IV Hominid 1: Morphological affinities of the earliest known American. *Genet. Mol. Biol.* **22**, 461–469 (1999). doi:10.1590/S1415-4757199000400001
42. R. González-José, M. C. Bortolini, F. R. Santos, S. L. Bonatto, The peopling of America: Craniofacial shape variation on a continental scale and its interpretation from an interdisciplinary view. *Am. J. Phys. Anthropol.* **137**, 175–187 (2008). doi:10.1002/ajpa.20854 Medline
43. C. de la Fuente, M. C. Ávila-Arcos, J. Galimany, M. L. Carpenter, J. R. Homburger, A. Blanco, P. Contreras, D. Cruz Dávalos, O. Reyes, M. San Roman, A. Moreno-Estrada, P. F. Campos, C. Eng, S. Huntsman, E. G. Burchard, A.-S. Malaspina, C. D. Bustamante, E. Willerslev, E. Llop, R. A. Verdugo, M. Moraga, Genomic insights into the origin and diversification of late maritime hunter-gatherers from the Chilean Patagonia. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E4006–E4012 (2018). doi:10.1073/pnas.1715688115 Medline
44. H. Schroeder, M. Sikora, S. Gopalakrishnan, L. M. Cassidy, P. Maisano Delser, M. Sandoval Velasco, J. G. Schraiber, S. Rasmussen, J. R. Homburger, M. C. Ávila-Arcos, M. E. Allentoft, J. V. Moreno-Mayar, G. Renaud, A. Gómez-Carballa, J. E. Laffoon, R. J. A. Hopkins, T. F. G. Higham, R. S. Carr, W. C. Schaffer, J. S. Day, M. Hoogland, A. Salas, C. D. Bustamante, R. Nielsen, D. G. Bradley, C. L. Hofman, E. Willerslev, Origins and genetic legacies of the Caribbean Taíno. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 2341–2346 (2018). doi:10.1073/pnas.1716839115 Medline
45. L. A. Louderback, D. K. Grayson, M. Llobera, Middle-Holocene climates and human population densities in the Great Basin, western USA. *Holocene* **21**, 366–373 (2011). doi:10.1177/0959683610374888
46. D. K. Grayson, *The Great Basin: A Natural Prehistory* (Univ. of California, revised and expanded ed., 2011).
47. N. N. Johannsen, G. Larson, D. J. Meltzer, M. Vander Linden, A composite window

- into human history. *Science* **356**, 1118–1120 (2017). doi:10.1126/science.aan0737 Medline
48. D. B. Madsen, D. Rhode, Eds., *Across the West: Human Population Movement and the Expansion of the Numa* (Univ. of Utah, 1994).
49. L. Campbell, *American Indian Languages: The Historical Linguistics of Native America* (Oxford Univ., 1997).
50. L. Bueno, G. Politis, L. Prates, J. Steele, Eds., A Late Pleistocene/Early Holocene Archaeological 14C Database for Central and South America: Palaeoenvironmental Contexts and Demographic Interpretations. *Quat. Int.* **301**, 1–158 (2013).
51. M. R. Waters, T. W. Stafford Jr., Redefining the age of Clovis: Implications for the peopling of the Americas. *Science* **315**, 1122–1126 (2007). doi:10.1126/science.1137166 Medline
52. D. J. Meltzer, *First Peoples in a New World: Colonizing Ice Age America* (Univ. of California, 2009).
53. T. D. Dillehay, C. Ramírez, M. Pino, M. B. Collins, J. Rossen, J. D. Pino-Navarro, Monte Verde: Seaweed, food, medicine, and the peopling of South America. *Science* **320**, 784–786 (2008). doi:10.1126/science.1156533 Medline
54. G. R. Scott, K. Schmitz, K. N. Heim, K. S. Paul, R. Schomberg, M. A. Pilloud, Sinodonty, Sundadonty, and the Beringian Standstill model: Issues of timing and migrations into the New World. *Quat. Int.* **466**, 233–246 (2018). doi:10.1016/j.quaint.2016.04.027
55. M. I. Eren, B. Buchanan, M. J. O'Brien, Social learning and technological evolution during the Clovis colonization of the New World. *J. Hum. Evol.* **80**, 159–170 (2015). doi:10.1016/j.jhevol.2015.01.002 Medline
56. L. Becerra-Valdivia, M. R. Waters, T. W. Stafford Jr., S. L. Anzick, D. Comeskey, T. Devière, T. Higham, Reassessing the chronology of the archaeological site of Anzick. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 7000–7003 (2018). doi:10.1073/pnas.1803624115 Medline
57. D. L. Jenkins, L. G. Davis, T. W. Stafford Jr., P. F. Campos, B. Hockett, G. T. Jones, L. S. Cummings, C. Yost, T. J. Connolly, R. M. Yohe 2nd, S. C. Gibbons, M. Raghavan, M. Rasmussen, J. L. A. Pajimans, M. Hofreiter, B. M. Kemp, J. L. Barta, C. Monroe, M. T. P. Gilbert, E. Willerslev, Clovis age Western Stemmed projectile points and human coprolites at the Paisley Caves. *Science* **337**, 223–228 (2012). doi:10.1126/science.1218443 Medline
58. C. E. Hughes, M. P. Rogers, A. C. Owings, B. Petzelt, J. Mitchell, H. Harry, T. Williams, D. Goldberg, D. Labuda, D. G. Smith, J. S. Cybulski, R. S. Malhi, Genetic Structure of First Nation Communities in the Pacific Northwest. *Hum. Biol.* **88**, 251–263 (2016). doi:10.13110/humanbiology.88.4.0251 Medline
59. R. Nielsen, J. M. Akey, M. Jakobsson, J. K. Pritchard, S. Tishkoff, E. Willerslev, Tracing the peopling of the world through genomics. *Nature* **541**, 302–310 (2017). doi:10.1038/nature21347 Medline
60. S. Lindgreen, AdapterRemoval: Easy cleaning of next-generation sequencing reads. *BMC Res. Notes* **5**, 337 (2012). doi:10.1186/1756-0500-5-337 Medline
61. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009). doi:10.1093/bioinformatics/btp324 Medline
62. M. Schubert, A. Ginolhac, S. Lindgreen, J. F. Thompson, K. A. S. Al-Rasheid, E. Willerslev, A. Krogh, L. Orlando, Improving ancient DNA read mapping against modern reference genomes. *BMC Genomics* **13**, 178 (2012). doi:10.1186/1471-2164-13-178 Medline
63. Picard, <http://picard.sourceforge.net>.
64. M. A. DePristo, E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire, C. Hartl, A. A. Philippakis, G. del Angel, M. A. Rivas, M. Hanna, A. McKenna, T. J. Fennell, A. M. Kernytsky, A. Y. Sivachenko, K. Cibulskis, S. B. Gabriel, D. Altshuler, M. J. Daly, A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011). doi:10.1038/ng.806 Medline
65. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin; 1000 Genome Project Data Processing Subgroup, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009). doi:10.1093/bioinformatics/btp352 Medline
66. B. N. Howie, P. Donnelly, J. Marchini, A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLOS Genet.* **5**, e1000529 (2009). doi:10.1371/journal.pgen.1000529 Medline
67. B. Howie, J. Marchini, M. Stephens, Genotype imputation with thousands of genomes. *G3 (Bethesda)* **1**, 457–470 (2011). doi:10.1534/g3.111.001198 Medline
68. SNPable Regions, <http://lh3lh3.users.sourceforge.net/snitable.shtml>.
69. The 1000 Genomes Project Consortium, A global reference for human genetic variation. *Nature* **526**, 68–74 (2015). doi:10.1038/nature15393 Medline
70. Q. Fu, A. Mitnik, P. L. F. Johnson, K. Bos, M. Lari, R. Bollongino, C. Sun, L. Giemsch, R. Schmitz, J. Burger, A. M. Ronchitelli, F. Martini, R. G. Cremonesi, J. Svoboda, P. Bauer, D. Caramelli, S. Castellano, D. Reich, S. Pääbo, J. Krause, A revised timescale for human evolution based on ancient mitochondrial genomes. *Curr. Biol.* **23**, 553–559 (2013). doi:10.1016/j.cub.2013.02.044 Medline
71. T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014). doi:10.1186/s12859-014-0356-4 Medline
72. F. Racimo, G. Renaud, M. Slatkin, Joint Estimation of Contamination, Error and Demography for Nuclear DNA from Ancient Humans. *PLOS Genet.* **12**, e1005972 (2016). doi:10.1371/journal.pgen.1005972 Medline
73. A. S. Dyke, A. Moore, L. Robertson, “Deglaciation of North America” (Geological Survey of Canada Open File 1547, 2003).
74. J. Lindo et al., The genetic prehistory of the Andean highlands: 7000 years BP though European contact. *Sci. Adv.* 10.1126/sciadv.aau4921 (2018).
75. C. Posth et al., Reconstructing the Deep Population History of Central and South America. *Cell* 10.1016/j.cell.2018.10.027 (2018).
76. H. Larsen, Trail Creek: Final report on the excavation of two caves on Seward Peninsula, Alaska. *Acta Arctica* **15**, 7–79 (1968).
77. K. Pasda, *Seward Peninsula, Alaska: Trail Creek Caves 2 and 9 Revisited: The Skeletal Remains* (BAR International Series, Archaeopress, 2012).
78. S. J. AlQahtani, M. P. Hector, H. M. Liversidge, Brief communication: The London atlas of human tooth development and eruption. *Am. J. Phys. Anthropol.* **142**, 481–490 (2010). doi:10.1002/ajpa.21258 Medline
79. H. M. Liversidge, T. Molleson, Variation in crown and root formation and eruption of human deciduous teeth. *Am. J. Phys. Anthropol.* **123**, 172–180 (2004). doi:10.1002/ajpa.10318 Medline
80. J. S. Cybulski et al., The Big Bar Lake burial: middle period human remains from the Canadian Plateau. *Can. J. Archaeol.* 55–78 (2007).
81. R. S. Malhi, B. M. Kemp, J. A. Eshleman, J. Cybulski, D. G. Smith, S. Cousins, H. Harry, Mitochondrial haplogroup M discovered in prehistoric North Americans. *J. Archaeol. Sci.* **34**, 642–648 (2007). doi:10.1016/j.jas.2006.07.004
82. D. L. Pokotyo, D. Mitchell, in *Plateau*, D. E. Walker, Ed. (Handbook of North American Indians, Smithsonian Institution, 1998), pp. 81–102.
83. S. M. Wheeler, Cave burials near Fallon, Nevada. *Nev. Hist. Soc. Q.* **40**, 15–23 (1997).
84. B. Hockett, E. Palus, A Brief History and Perspective on Spirit Cave, Nevada. *PaleoAmerica* **4**, 1–7 (2018). doi:10.1080/2055563.2017.1412211
85. A. Dansie, Early Holocene burials in Nevada: Overview of localities, research and legal issues. *Nev. Hist. Soc. Q.* **40**, 4–14 (1997).
86. C. S. Fowler, E. M. Hattori, in *The Great Basin: People and Place in Ancient Times*, C. S. Fowler, D. D. Fowler, Eds. (Popular Southwestern Archaeology Books, School for Advanced Research Press, 2008), pp. 60–67.
87. P. Barker, C. Ellis, S. Damadio, “Determination of cultural affiliation of ancient human remains from Spirit Cave, Nevada” (Unpublished report on file, Bureau of Land Management, Nevada State Office, Reno, 2000).
88. H. J. Hecht Edgar, Paleopathology of the Wizards Beach Man (Ahur 2023) and the Spirit Cave mummy (Ahur 2064). *Nev. Hist. Soc. Q.* **40**, 57–61 (1997).
89. R. L. Jantz, D. W. Owsley, Pathology, taphonomy, and cranial morphometrics of the Spirit Cave mummy. *Nev. Hist. Soc. Q.* **40**, 62–82 (1997).
90. J. F. Powell, *The First Americans: Race, Evolution, and the Origin of Native Americans* (Cambridge Univ., 2005).
91. J. F. Powell, W. A. Neves, Craniofacial morphology of the first Americans: Pattern and process in the peopling of the New World. *Am. J. Phys. Anthropol.* **110** (suppl. 29), 153–188 (1999). Medline
92. D. W. Owsley, R. L. Jantz, Eds., *Kennewick Man: The Scientific Investigation of an Ancient American Skeleton* (Texas A&M Univ., ed. 1, 2014).
93. B. M. Auerbach, Skeletal variation among early Holocene North American humans: Implications for origins and diversity in the Americas. *Am. J. Phys. Anthropol.* **149**, 525–536 (2012). doi:10.1002/ajpa.22154 Medline
94. R. F. Heizer, L. K. Napton, *Archaeology and the Prehistoric Great Basin Lacustrine Subsistence Regime as Seen from Lovelock Cave, Nevada*, vol. 10 of Contributions

- of the University of California Archaeological Research Facility* (Univ. of California, 1970).
95. W. A. Neves et al., in *Paleoamerican Odyssey*, K. E. Graf, C. V. Ketron, M. R. Waters, Eds. (Texas A&M Univ., Texas A&M Univ. ed. 1., 2014), pp. 397–412.
96. P. W. Lund, Carta escripta da Lagôa Santa (Minas Geraes), ao Sr. 1.o Secretario do Instituto, pelo socio honorario Sr. Dr. Lund. *Revista trimensal de Historia e Geographia* **4**, 80–87 (1842).
97. P. W. Lund, Carta do Dr. Lund escripta de Lagôa Santa (Minas Geraes) a 21 de abril de 1844. *Revista trimensal de Historia e Geographia* **6**, 326–334 (1844).
98. P. W. Lund, Notice sur des ossements humains fossiles, trouvés dans une grotte du Brésil. *Mémoires de la Société Royale des Antiquaires du Nord* **2**, 49–77 (1843).
99. A. Hrdlička, *Early Man in South America* (Bureau of American Ethnology Bulletin 52, Government Printing Office, 1912).
100. O. R. Ortiz-Troncoso, Los yacimientos de Punta Santa Ana y Bahía Buena (Patagonia Austral). Excavaciones y fechados radiocarbónicos. *An. Inst. Patagon.* **7**, 93–122 (1975).
101. D. Legoupil, P. Sellier, La sepultura de la cueva Ayayema (isla Madre de Dios, archipiélagos occidentales de Patagonia). *Magallania* **32**, 115–124 (2004).
102. F. Morello, J. Torres, I. Martínez, K. Rodriguez, M. Arroyo-Kalin, C. French, V. Sierpe, M. San Román, Arqueología de la Punta Santa Ana: Reconstrucción de secuencias de ocupación de cazadores-recolectores marinos del Estrecho de Magallanes, Patagonia. *Magallania* **40**, 129–149 (2012). doi:[10.4067/S0718-22442012000200008](https://doi.org/10.4067/S0718-22442012000200008)
103. A. G. Hogg, Q. Hua, P. G. Blackwell, M. Niu, C. E. Buck, T. P. Guilderson, T. J. Heaton, J. G. Palmer, P. J. Reimer, R. W. Reimer, C. S. M. Turney, S. R. H. Zimmerman, SHCal13 Southern Hemisphere Calibration, 0–50,000 Years cal BP. *Radiocarbon* **55**, 1889–1903 (2013). doi:[10.2458/azu\\_js\\_rc.55.16783](https://doi.org/10.2458/azu_js_rc.55.16783)
104. M. Stuiver, P. J. Reimer, Extended 14C Data Base and Revised CALIB 3.0 14C Age Calibration Program. *Radiocarbon* **35**, 215–230 (1993). doi:[10.1017/S003382200013904](https://doi.org/10.1017/S003382200013904)
105. J. Schobinger, Los santuarios de altura incaicos y el Aconcagua: Aspectos generales e interpretativos. *Relaciones de la Sociedad Argentina de Antropología* **24**, 7–27 (1999).
106. S. Quevedo, E. Durán, Ofrendas a los dioses en las montañas: Santuarios de altura de la cultura inka. *Bol. Mus. Nac. Hist. Nat.* **43**, 193–206 (1992).
107. A. Salas, L. Catelli, J. Pardo-Seco, A. Gómez-Carballa, F. Martinón-Torres, J. Roberto-Barcena, C. Vullo, Y-chromosome Peruvian origin of the 500-year-old Inca child mummy sacrificed in Cerro Aconcagua (Argentina). *Sci. Bull.* (2018). doi:[10.1016/j.scib.2018.08.009](https://doi.org/10.1016/j.scib.2018.08.009)
108. A. R. Radcliffe-Brown, *The Andaman Islanders: A Study in Social Anthropology* (Forgotten Books, 2015).
109. R. S. Mahi, J. S. Cybulski, R. Y. Tito, J. Johnson, H. Harry, C. Dan, Brief communication: Mitochondrial haplotype C4c confirmed as a founding genome in the Americas. *Am. J. Phys. Anthropol.* **141**, 494–497 (2010). Medline
110. J. Ruhs, “Memorandum: Spirit Cave human remains and associated funerary objects – conclusions and determination concerning Native American status, cultural affiliation and disposition” (Unpublished report on file, Bureau of Land Management, Nevada State Office, Reno, 2016).
111. Department of the Interior, Notice of Inventory Completion: U.S. Department of the Interior, Bureau of Land Management, Nevada State Office, Reno, NV. *Fed. Regist.* **81**, 71760–71762 (2016).
112. E. Callaway, North America’s oldest mummy returned to US tribe after genome sequencing. *Nature* **540**, 178–179 (2016). doi:[10.1038/540178a](https://doi.org/10.1038/540178a)
113. F. Brock, T. Higham, P. Ditchfield, C. B. Ramsey, Current Pretreatment Methods for AMS Radiocarbon Dating at the Oxford Radiocarbon Accelerator Unit (Orau). *Radiocarbon* **52**, 103–112 (2010). doi:[10.1017/S003382200045069](https://doi.org/10.1017/S003382200045069)
114. T. Deviese, D. Comeskey, J. McCullagh, C. Bronk Ramsey, T. Higham, New protocol for compound-specific radiocarbon analysis of archaeological bones. *Rapid Commun. Mass Spectrom.* **32**, 373–379 (2018). doi:[10.1002/rctm.8047](https://doi.org/10.1002/rctm.8047) Medline
115. M. T. P. Gilbert, H.-J. Bandelt, M. Hofreiter, I. Barnes, Assessing ancient DNA studies. *Trends Ecol. Evol.* **20**, 541–544 (2005). doi:[10.1016/j.tree.2005.07.005](https://doi.org/10.1016/j.tree.2005.07.005) Medline
116. E. Willerslev, A. Cooper, Ancient DNA. *Proc. Biol. Sci.* **272**, 3–16 (2005). doi:[10.1098/rspb.2004.2813](https://doi.org/10.1098/rspb.2004.2813) Medline
117. P. B. Damgaard, A. Margaryan, H. Schroeder, L. Orlando, E. Willerslev, M. E. Allentoft, Improving access to endogenous DNA in ancient bones and teeth. *Sci. Rep.* **5**, 11184 (2015). doi:[10.1038/srep11184](https://doi.org/10.1038/srep11184) Medline
118. M. E. Allentoft, M. Sikora, K.-G. Sjögren, S. Rasmussen, M. Rasmussen, J. Stenderup, P. B. Damgaard, H. Schroeder, T. Ahlström, L. Vinner, A.-S. Malaspina, A. Margaryan, T. Higham, D. Chivall, N. Lynnerup, L. Harvig, J. Baron, P. Della Casa, P. Dąbrowski, P. R. Duffy, A. V. Ebel, A. Epimakhov, K. Frei, M. Furmanek, T. Gralak, A. Gromov, S. Gronkiewicz, G. Grupe, T. Hajdu, R. Jarzysz, V. Khartanovich, A. Khokhlov, V. Kiss, J. Kolář, A. Kriiska, I. Lasak, C. Longhi, G. McGlynn, A. Merkevicius, I. Merkyte, M. Metspalu, R. Mkrtchyan, V. Moiseyev, L. Paja, G. Pálfi, D. Pokutta, Ł. Pospiszny, T. D. Price, L. Saag, M. Sablin, N. Shishlina, V. Smrčka, V. I. Soenov, V. Szeverényi, G. Tóth, S. V. Trifanova, L. Varul, M. Vicze, L. Yepiskoposyan, V. Zhitenev, L. Orlando, T. Sicheritz-Pontén, S. Brunak, R. Nielsen, K. Kristiansen, E. Willerslev, Population genomics of Bronze Age Eurasia. *Nature* **522**, 167–172 (2015). doi:[10.1038/nature14507](https://doi.org/10.1038/nature14507) Medline
119. I. Glocke, M. Meyer, Extending the spectrum of DNA sequences retrieved from ancient bones and teeth. *Genome Res.* **27**, 1230–1237 (2017). doi:[10.1101/gr.219675.116](https://doi.org/10.1101/gr.219675.116) Medline
120. M. Meyer, M. Kircher, Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb. Protoc.* **2010**, t5448 (2010). doi:[10.1101/pdb.prot5448](https://doi.org/10.1101/pdb.prot5448) Medline
121. J. Dabney, M. Knapp, I. Glocke, M.-T. Gansauge, A. Weihmann, B. Nickel, C. Valdiosera, N. García, S. Pääbo, J.-L. Arsuaga, M. Meyer, Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 15758–15763 (2013). doi:[10.1073/pnas.1314445110](https://doi.org/10.1073/pnas.1314445110) Medline
122. L. Orlando, A. Ginolhac, G. Zhang, D. Froese, A. Albrechtsen, M. Stiller, M. Schubert, E. Cappellini, B. Petersen, I. Moltke, P. L. F. Johnson, M. Fumagalli, J. T. Vilstrup, M. Raghavan, T. Korneliussen, A.-S. Malaspina, J. Vogt, D. Szklarczyk, C. D. Kelstrup, J. Vinther, A. Dolocan, J. Stenderup, A. M. V. Velazquez, J. Cahill, M. Rasmussen, X. Wang, J. Min, G. D. Zazula, A. Seguin-Orlando, C. Mortensen, K. Magnussen, J. F. Thompson, J. Weinstock, K. Gregersen, K. H. Røed, V. Eisenmann, C. J. Rubin, D. C. Miller, D. F. Antczak, M. F. Bertelsen, S. Brunak, K. A. S. Al-Rasheid, O. Ryder, L. Andersson, J. Mundy, A. Krogh, M. T. P. Gilbert, K. Kjær, T. Sicheritz-Pontén, L. J. Jensen, J. V. Olsen, M. Hofreiter, R. Nielsen, B. Shapiro, J. Wang, E. Willerslev, Recalibrating Equus evolution using the genome sequence of an early Middle Pleistocene horse. *Nature* **499**, 74–78 (2013). doi:[10.1038/nature12323](https://doi.org/10.1038/nature12323) Medline
123. C. Carøe, S. Gopalakrishnan, L. Vinner, S. S. T. Mak, M. H. S. Sinding, J. A. Samaniego, N. Wales, T. Sicheritz-Pontén, M. T. P. Gilbert, Single-tube library preparation for degraded DNA. *Methods Ecol. Evol.* **9**, 410–419 (2018). doi:[10.1111/2041-210X.12871](https://doi.org/10.1111/2041-210X.12871)
124. S. S. T. Mak, S. Gopalakrishnan, C. Carøe, C. Geng, S. Liu, M. S. Sinding, L. F. K. Kuderna, W. Zhang, S. Fu, F. G. Vieira, M. Germonpré, H. Bocherens, S. Fedorov, B. Petersen, T. Sicheritz-Pontén, T. Marques-Bonet, G. Zhang, H. Jiang, M. T. P. Gilbert, Comparative performance of the BGISEQ-500 vs Illumina HiSeq2500 sequencing platforms for palaeogenomic sequencing. *Gigascience* **6**, 1–13 (2017). doi:[10.1093/gigascience/gix049](https://doi.org/10.1093/gigascience/gix049) Medline
125. M. Kircher, S. Sawyer, M. Meyer, Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res.* **40**, e3–e3 (2012). doi:[10.1093/nar/gkr771](https://doi.org/10.1093/nar/gkr771) Medline
126. M. Meyer, M. Kircher, M.-T. Gansauge, H. Li, F. Racimo, S. Mallick, J. G. Schraiber, F. Jay, K. Prüfer, C. de Filippo, P. H. Sudmant, C. Alkan, Q. Fu, R. Do, N. Rohland, A. Tandon, M. Siebauer, R. E. Green, K. Bryc, A. W. Briggs, U. Stenzel, J. Dabney, J. Shendure, J. Kitzman, M. F. Hammer, M. V. Shunkov, A. P. Derevianko, N. Patterson, A. M. Andrés, E. E. Eichler, M. Slatkin, D. Reich, J. Kelso, S. Pääbo, A high-coverage genome sequence from an archaic Denisovan individual. *Science* **338**, 222–226 (2012). doi:[10.1126/science.1224344](https://doi.org/10.1126/science.1224344) Medline
127. A.-S. Malaspina, M. C. Westaway, C. Muller, V. C. Sousa, O. Lao, I. Alves, A. Bergström, G. Athanasiadis, J. Y. Cheng, J. E. Crawford, T. H. Heupink, E. Macholdt, S. Peischl, S. Rasmussen, S. Schiffels, S. Subramanian, J. L. Wright, A. Albrechtsen, C. Barbieri, I. Dupanloup, A. Eriksson, A. Margaryan, I. Moltke, I. Pugach, T. S. Korneliussen, I. P. Levkivskyi, J. V. Moreno-Mayar, S. Ni, F. Racimo, M. Sikora, Y. Xue, F. A. Aghakhanian, N. Brucato, S. Brunak, P. F. Campos, W. Clark, S. Ellingvåg, G. Fourmile, P. Gerbault, D. Injie, G. Koki, M. Leavesley, B. Logan, A.

- Lynch, E. A. Matisoo-Smith, P. J. McAllister, A. J. Mentzer, M. Metspalu, A. B. Migliano, L. Murgha, M. E. Phipps, W. Pormat, D. Reynolds, F.-X. Ricaut, P. Siba, M. G. Thomas, T. Wales, C. M. Wall, S. J. Oppenheimer, C. Tyler-Smith, R. Durbin, J. Dortsch, A. Manica, M. H. Schierup, R. A. Foley, M. M. Lahr, C. Bowern, J. D. Wall, T. Mailund, M. Stoneking, R. Nielsen, M. S. Sandhu, L. Excoffier, D. M. Lambert, E. Willerslev, A genomic history of Aboriginal Australia. *Nature* **538**, 207–214 (2016). doi:10.1038/nature18299 Medline
128. A. W. Briggs, U. Stenzel, P. L. F. Johnson, R. E. Green, J. Kelso, K. Prüfer, M. Meyer, J. Krause, M. T. Ronan, M. Lachmann, S. Pääbo, Patterns of damage in genomic DNA sequences from a Neandertal. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 14616–14621 (2007). doi:10.1073/pnas.0704665104 Medline
129. W. J. Kent, C. W. Sugnet, T. S. Furey, K. M. Roskin, T. H. Pringle, A. M. Zahler, D. Haussler, The human genome browser at UCSC. *Genome Res.* **12**, 996–1006 (2002). doi:10.1101/gr.229102 Medline
130. K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002). doi:10.1093/nar/gkf436 Medline
131. K. Katoh, M. C. Frith, Adding unaligned sequences into an existing alignment using MAFFT and LAST. *Bioinformatics* **28**, 3144–3146 (2012). doi:10.1093/bioinformatics/bts578 Medline
132. A. Gelman, D. B. Rubin, Inference from iterative simulation using multiple sequences. *Stat. Sci.* **7**, 457–472 (1992). doi:10.1214/ss/1177011136
133. M. Rasmussen, X. Guo, Y. Wang, K. E. Lohmueller, S. Rasmussen, A. Albrechtsen, L. Skotte, S. Lindgreen, M. Metspalu, T. Jombart, T. Kivisild, W. Zhai, A. Eriksson, A. Manica, L. Orlando, F. M. De La Vega, S. Tridico, E. Metspalu, K. Nielsen, M. C. Ávila-Arcos, J. V. Moreno-Mayar, C. Müller, J. Dortsch, M. T. P. Gilbert, O. Lund, A. Wesolowska, M. Karmin, L. A. Weinert, B. Wang, J. Li, S. Tai, F. Xiao, T. Hanihara, G. van Driem, A. R. Jha, F.-X. Ricaut, P. de Knijff, A. B. Migliano, I. Gallego Romero, K. Kristiansen, D. M. Lambert, S. Brunak, P. Forster, B. Brinkmann, O. Nehlich, M. Bunce, M. Richards, R. Gupta, C. D. Bustamante, A. Krogh, R. A. Foley, M. M. Lahr, F. Balloux, T. Sicheritz-Pontén, R. Villemans, R. Nielsen, J. Wang, E. Willerslev, An Aboriginal Australian genome reveals separate human dispersals into Asia. *Science* **334**, 94–98 (2011). doi:10.1126/science.1211777 Medline
134. D. M. Altshuler, R. A. Gibbs, L. Peltonen, D. M. Altshuler, R. A. Gibbs, L. Peltonen, E. Dermitzakis, S. F. Schaffner, F. Yu, L. Peltonen, E. Dermitzakis, P. E. Bonnen, D. M. Altshuler, R. A. Gibbs, P. I. de Bakker, P. Deloukas, S. B. Gabriel, R. Gwilliam, S. Hunt, M. Inouye, X. Jia, A. Palotie, M. Parkin, P. Whittaker, F. Yu, K. Chang, A. Hawes, L. R. Lewis, Y. Ren, D. Wheeler, R. A. Gibbs, D. M. Muzny, C. Barnes, K. Darvishi, M. Hurles, J. M. Korn, K. Kristiansson, C. Lee, S. A. McCarron, J. Nemesh, E. Dermitzakis, A. Keinan, S. B. Montgomery, S. Pollack, A. L. Price, N. Soranzo, P. E. Bonnen, R. A. Gibbs, C. Gonzaga-Jauregui, A. Keinan, A. L. Price, F. Yu, V. Anttila, W. Brodeur, M. J. Daly, S. Leslie, G. McVean, L. Moutsianas, H. Nguyen, S. F. Schaffner, Q. Zhang, M. J. Ghori, R. McGinnis, W. McLaren, S. Pollack, A. L. Price, S. F. Schaffner, F. Takeuchi, S. R. Grossman, I. Shlyakhter, E. B. Hostetter, P. C. Sabeti, C. A. Adebamowo, M. W. Foster, D. R. Gordon, J. Licinio, M. C. Manca, P. A. Marshall, I. Matsuda, D. Ngare, V. O. Wang, D. Reddy, C. N. Rotimi, C. D. Royal, R. R. Sharp, C. Zeng, L. D. Brooks, J. E. McEwen; International HapMap 3 Consortium, Integrating common and rare genetic variation in diverse human populations. *Nature* **467**, 52–58 (2010). doi:10.1038/nature09298 Medline
135. P. Skoglund, J. Storå, A. Götherström, M. Jakobsson, Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* **40**, 4477–4482 (2013). doi:10.1016/j.jas.2013.07.004
136. E. Cappellini, A. Prohaska, F. Racimo, F. Welker, M. W. Pedersen, M. E. Allentoft, P. de Barros Damgaard, P. Gutenbrunner, J. Dunne, S. Hammann, M. Roffet-Salque, M. Ilardo, J. V. Moreno-Mayar, Y. Wang, M. Sikora, L. Vinner, J. Cox, R. P. Evershed, E. Willerslev, Ancient Biomolecules and Evolutionary Inference. *Annu. Rev. Biochem.* **87**, 1029–1060 (2018). doi:10.1146/annurev-biochem-062917-012002 Medline
137. T. Kivisild, The study of human Y chromosome variation through ancient DNA. *Hum. Genet.* **136**, 529–546 (2017). doi:10.1007/s00439-017-1773-z Medline
138. P. de Barros Damgaard, R. Martiniano, J. Kamm, J. V. Moreno-Mayar, G. Kroonen, M. Peyrot, G. Barjamovic, S. Rasmussen, C. Zacho, N. Bairukhanov, V. Zaibert, V. Merz, A. Biddanda, I. Merz, V. Loman, V. Evdokimov, E. Usmanova, B. Hemphill, A. Seguin-Orlando, F. E. Yediay, I. Ullah, K.-G. Sjögren, K. H. Iversen, J. Choin, C. de la Fuente, M. Ilardo, H. Schroeder, V. Moiseyev, A. Gromov, A. Polyakov, S. Omura, S. Y. Senyurt, H. Ahmad, C. McKenzie, A. Margaryan, A. Hameed, A. Samad, N. Gul, M. H. Khokhar, O. I. Goriunova, V. I. Bazaliiskii, J. Novembre, A. W. Weber, L. Orlando, M. E. Allentoft, R. Nielsen, K. Kristiansen, M. Sikora, A. K. Outram, R. Durbin, E. Willerslev, The first horse herders and the impact of early Bronze Age steppe expansions into Asia. *Science* **360**, eaar7711 (2018). doi:10.1126/science.aar7711 Medline
139. G. D. Poznik, B. M. Henn, M.-C. Yee, E. Slivierska, G. M. Euskirchen, A. A. Lin, M. Snyder, L. Quintana-Murci, J. M. Kidd, P. A. Underhill, C. D. Bustamante, Sequencing Y chromosomes resolves discrepancy in time to common ancestor of males versus females. *Science* **341**, 562–565 (2013). doi:10.1126/science.1237619 Medline
140. E. Garrison, G. Marth, Haplotype-based variant detection from short-read sequencing. *arXiv:1207.3907* [q-bio.GN] (20 July 2012).
141. G. D. Poznik, Y. Xue, F. L. Mendez, T. F. Willems, A. Massaia, M. A. Wilson Sayres, Q. Ayub, S. A. McCarthy, A. Narechania, S. Kashin, Y. Chen, R. Banerjee, J. L. Rodriguez-Flores, M. Cerezo, H. Shao, M. Gymrek, A. Malhotra, S. Louzada, R. Desalle, G. R. S. Ritchie, E. Cerveira, T. W. Fitzgerald, E. Garrison, A. Marcketta, D. Mittelman, M. Romanovitch, C. Zhang, X. Zheng-Bradley, G. R. Abecasis, S. A. McC Carroll, P. Flicek, P. A. Underhill, L. Coin, D. R. Zerbino, F. Yang, C. Lee, L. Clarke, A. Auton, Y. Erlich, R. E. Handsaker, C. D. Bustamante, C. Tyler-Smith; 1000 Genomes Project Consortium, Punctuated bursts in human male demography inferred from 1,244 worldwide Y-chromosome sequences. *Nat. Genet.* **48**, 593–599 (2016). doi:10.1038/ng.3559 Medline
142. A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014). doi:10.1093/bioinformatics/btu033 Medline
143. M. S. Jota, D. R. Lacerda, J. R. Sandoval, P. P. R. Vieira, D. Ohasi, J. E. Santos-Júnior, O. Acosta, C. Cuellar, S. Revollo, C. Paz-Y-Miño, R. Fujita, G. A. Vallejo, T. G. Schurr, E. M. Tarazona-Santos, S. Dj. Pena, Q. Ayub, C. Tyler-Smith, F. R. Santos; Genographic Consortium, New native South American Y chromosome lineages. *J. Hum. Genet.* **61**, 593–603 (2016). doi:10.1038/jhg.2016.26 Medline
144. M. Rasmussen, Y. Li, S. Lindgreen, J. S. Pedersen, A. Albrechtsen, I. Moltke, M. Metspalu, E. Metspalu, T. Kivisild, R. Gupta, M. Bertalan, K. Nielsen, M. T. P. Gilbert, Y. Wang, M. Raghavan, P. F. Campos, H. M. Kamp, A. S. Wilson, A. Gledhill, S. Tridico, M. Bunce, E. D. Lorenzen, J. Binladen, X. Guo, J. Zhao, X. Zhang, H. Zhang, Z. Li, M. Chen, L. Orlando, K. Kristiansen, M. Bak, N. Tommerup, C. Bendixen, T. L. Pierre, B. Grønnow, M. Meldgaard, C. Andreassen, S. A. Fedorova, L. P. Osipova, T. F. G. Higham, C. B. Ramsey, T. V. Hansen, F. C. Nielsen, M. H. Crawford, S. Brunak, T. Sicheritz-Pontén, R. Villemans, R. Nielsen, A. Krogh, J. Wang, E. Willerslev, Ancient human genome sequence of an extinct Palaeo-Eskimo. *Nature* **463**, 757–762 (2010). doi:10.1038/nature08835 Medline
145. M. C. Dulik, A. C. Owings, J. B. Gaieski, M. G. Vilar, A. Andre, C. Lennie, M. A. Mackenzie, I. Kritschs, S. Snowshoe, R. Wright, J. Martin, N. Gibson, T. D. Andrews, T. G. Schurr, S. Adhikarla, C. J. Adler, E. Balanovska, O. Balanovsky, J. Bertranpetti, A. C. Clarke, D. Comas, A. Cooper, C. S. I. Der Sarkissian, A. GaneshPrasad, W. Haak, M. Haber, A. Hobbs, A. Javed, L. Jin, M. E. Kaplan, S. Li, B. Martinez-Cruz, E. A. Matisoo-Smith, M. Mele, N. C. Merchant, R. J. Mitchell, L. Parida, R. Pitchappan, D. E. Platt, L. Quintana-Murci, C. Renfrew, D. R. Lacerda, A. K. Royyuru, F. R. Santos, H. Soodyall, D. F. Soria Hernanz, P. Swamikrishnan, C. Tyler-Smith, A. V. Santhakumari, P. P. Vieira, R. S. Wells, P. A. Zalloua, J. S. Ziegler; Genographic Consortium, Y-chromosome analysis reveals genetic divergence and new founding native lineages in Athabaskan- and Eskimoan-speaking populations. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 8471–8476 (2012). doi:10.1073/pnas.1118760109 Medline
146. S. D. J. Pena, F. R. Santos, N. O. N. Bianchi, C. M. Bravi, F. R. Carnese, F. Rothhammer, T. Gerelsaikhan, B. Munkhtuja, T. Oyunsuren, A major founder Y-chromosome haplotype in Amerindians. *Nat. Genet.* **11**, 15–16 (1995). doi:10.1038/ng0995-15 Medline
147. T. D. Dillehay, C. Ocampo, J. Saavedra, A. O. Sawakuchi, R. M. Vega, M. Pino, M. B. Collins, L. Scott Cummings, I. Arregui, X. S. Villagrán, G. A. Hartmann, M. Mella, A. González, G. Dix, New Archaeological Evidence for an Early Human Presence at Monte Verde, Chile. *PLOS ONE* **10**, e0141923 (2015). doi:10.1371/journal.pone.0141923 Medline
148. J. F. O'Connell, J. Allen, The process, biotic impact, and global implications of the human colonization of Sahul about 47,000 years ago. *J. Archaeol. Sci.* **56**, 73–84 (2015). doi:10.1016/j.jas.2015.02.020

149. J. F. O'Connell, J. Allen, The restaurant at the end of the universe: Modelling the colonisation of Sahul. *Aust. Archaeol.* **74**, 5–17 (2012). [doi:10.1080/03122417.2012.11681932](https://doi.org/10.1080/03122417.2012.11681932)
150. M. Hubbe, W. A. Neves, K. Harvati, Testing evolutionary and dispersion scenarios for the settlement of the new world. *PLOS ONE* **5**, e11105 (2010). [doi:10.1371/journal.pone.0011105](https://doi.org/10.1371/journal.pone.0011105) Medline
151. M. Hubbe, A. Strauss, A. Hubbe, W. A. Neves, Early South Americans Cranial Morphological Variation and the Origin of American Biological Diversity. *PLOS ONE* **10**, e0138090 (2015). [doi:10.1371/journal.pone.0138090](https://doi.org/10.1371/journal.pone.0138090) Medline
152. Q. Fu, H. Li, P. Moorjani, F. Jay, S. M. Slepchenko, A. A. Bondarev, P. L. F. Johnson, A. Aximu-Petri, K. Prüfer, C. de Filippo, M. Meyer, N. Zwyns, D. C. Salazar-García, Y. V. Kuzmin, S. G. Keates, P. A. Kosintsev, D. I. Razhev, M. P. Richards, N. V. Peristov, M. Lachmann, K. Douka, T. F. G. Higham, M. Slatkin, J.-J. Hublin, D. Reich, J. Kelso, T. B. Viola, S. Pääbo, Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* **514**, 445–449 (2014). [doi:10.1038/nature13810](https://doi.org/10.1038/nature13810) Medline
153. Q. Fu, M. Hajdinjak, O. T. Moldovan, S. Constantin, S. Mallick, P. Skoglund, N. Patterson, N. Rohland, I. Lazaridis, B. Nickel, B. Viola, K. Prüfer, M. Meyer, J. Kelso, D. Reich, S. Pääbo, An early modern human from Romania with a recent Neanderthal ancestor. *Nature* **524**, 216–219 (2015). [doi:10.1038/nature14558](https://doi.org/10.1038/nature14558) Medline
154. T. M. Karafet, F. L. Mendez, H. Sudoyo, J. S. Lansing, M. F. Hammer, Improved phylogenetic resolution and rapid diversification of Y-chromosome haplogroup K-M526 in Southeast Asia. *Eur. J. Hum. Genet.* **23**, 369–373 (2015). [doi:10.1038/ejhg.2014.106](https://doi.org/10.1038/ejhg.2014.106) Medline
155. A. Bergström, N. Nagle, Y. Chen, S. McCarthy, M. O. Pollard, Q. Ayub, S. Wilcox, L. Wilcox, R. A. H. van Oorschot, P. McAllister, L. Williams, Y. Xue, R. J. Mitchell, C. Tyler-Smith, Deep Roots for Aboriginal Australian Y Chromosomes. *Curr. Biol.* **26**, 809–813 (2016). [doi:10.1016/j.cub.2016.01.028](https://doi.org/10.1016/j.cub.2016.01.028) Medline
156. M. Mondal, A. Bergström, Y. Xue, F. Calafell, H. Laayouni, F. Casals, P. P. Majumder, C. Tyler-Smith, J. Bertranpetti, Y-chromosomal sequences of diverse Indian populations and the ancestry of the Andamanese. *Hum. Genet.* **136**, 499–510 (2017). [doi:10.1007/s00439-017-1800-0](https://doi.org/10.1007/s00439-017-1800-0) Medline
157. G. Renaud, V. Slon, A. T. Duggan, J. Kelso, Schmutz: Estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol.* **16**, 224 (2015). [doi:10.1186/s13059-015-0776-0](https://doi.org/10.1186/s13059-015-0776-0) Medline
158. M. van Oven, M. Kayser, Updated comprehensive phylogenetic tree of global human mitochondrial DNA variation. *Hum. Mutat.* **30**, E386–E394 (2009). [doi:10.1002/humu.20921](https://doi.org/10.1002/humu.20921) Medline
159. C. Posth, G. Renaud, A. Mittnik, D. G. Drucker, H. Rougier, C. Cupillard, F. Valentin, C. Thevenet, A. Furtwängler, C. Wißing, M. Francken, M. Malina, M. Bolus, M. Lari, E. Gigli, G. Capecchi, I. Crevecoeur, C. Beauval, D. Flas, M. Germenpré, J. van der Plicht, R. Cottiaux, B. Gély, A. Ronchitelli, K. Wehrberger, D. Grigorescu, J. Svoboda, P. Semal, D. Caramelli, H. Bocherens, K. Harvati, N. J. Conard, W. Haak, A. Powell, J. Krause, Pleistocene Mitochondrial Genomes Suggest a Single Major Dispersal of Non-Africans and a Late Glacial Population Turnover in Europe. *Curr. Biol.* **26**, 827–833 (2016). [doi:10.1016/j.cub.2016.01.037](https://doi.org/10.1016/j.cub.2016.01.037) Medline
160. A. T. Duggan, A. J. T. Harris, S. Marciniak, I. Marshall, M. Kuch, A. Kitchen, G. Renaud, J. Southon, B. Fuller, J. Young, S. Fiedel, G. B. Golding, V. Grimes, H. Poinar, Genetic Discontinuity between the Maritime Archaic and Beothuk Populations in Newfoundland, Canada. *Curr. Biol.* **27**, 3149–3156.e11 (2017). [doi:10.1016/j.cub.2017.08.053](https://doi.org/10.1016/j.cub.2017.08.053) Medline
161. S. Kumar, G. Stecher, K. Tamura, MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874 (2016). [doi:10.1093/molbev/msw054](https://doi.org/10.1093/molbev/msw054) Medline
162. H. Weissensteiner, D. Pacher, A. Kloss-Brandstätter, L. Forer, G. Specht, H.-J. Bandelt, F. Kronenberg, A. Salas, S. Schönher, HaploGrep 2: Mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Res.* **44**, W58–W63 (2016). [doi:10.1093/nar/gkw233](https://doi.org/10.1093/nar/gkw233) Medline
163. D. Vianello, F. Sevini, G. Castellani, L. Lomartire, M. Capri, C. Franceschi, HAPLOFIND: A new method for high-throughput mtDNA haplogroup assignment. *Hum. Mutat.* **34**, 1189–1194 (2013). [doi:10.1002/humu.22356](https://doi.org/10.1002/humu.22356) Medline
164. P. Soares, L. Ermini, N. Thomson, M. Mormina, T. Rito, A. Röhl, A. Salas, S. Oppenheimer, V. Macaulay, M. B. Richards, Correcting for purifying selection: An improved human mitochondrial molecular clock. *Am. J. Hum. Genet.* **84**, 740–759 (2009). [doi:10.1016/j.ajhg.2009.05.001](https://doi.org/10.1016/j.ajhg.2009.05.001) Medline
165. R. Bouckaert, J. Heled, D. Kühnert, T. Vaughan, C.-H. Wu, D. Xie, M. A. Suchard, A. Rambaut, A. J. Drummond, BEAST 2: A software platform for Bayesian evolutionary analysis. *PLOS Comput. Biol.* **10**, e1003537 (2014). [doi:10.1371/journal.pcbi.1003537](https://doi.org/10.1371/journal.pcbi.1003537) Medline
166. T. G. Schurr, S. W. Ballinger, Y. Y. Gan, J. A. Hodge, D. A. Merriwether, D. N. Lawrence, W. C. Knowler, K. M. Weiss, D. C. Wallace, Amerindian mitochondrial DNAs have rare Asian mutations at high frequencies, suggesting they derived from four primary maternal lineages. *Am. J. Hum. Genet.* **46**, 613–623 (1990). [Medline](#)
167. A. Torroni, T. G. Schurr, M. F. Cabell, M. D. Brown, J. V. Neel, M. Larsen, D. G. Smith, C. M. Vullo, D. C. Wallace, Asian affinities and continental radiation of the four founding Native American mtDNAs. *Am. J. Hum. Genet.* **53**, 563–590 (1993). [Medline](#)
168. A. Achilli, U. A. Perego, C. M. Bravi, M. D. Coble, Q.-P. Kong, S. R. Woodward, A. Salas, A. Torroni, H.-J. Bandelt, The phylogeny of the four pan-American MtDNA haplogroups: Implications for evolutionary and disease studies. *PLOS ONE* **3**, e1764 (2008). [doi:10.1371/journal.pone.0001764](https://doi.org/10.1371/journal.pone.0001764) Medline
169. S. Brandini, P. Bergamaschi, M. F. Cerna, F. Gandini, F. Bastaroli, E. Bertolini, C. Cereda, L. Ferretti, A. Gómez-Carballa, V. Battaglia, A. Salas, O. Semino, A. Achilli, A. Olivieri, A. Torroni, The Paleo-Indian Entry into South America According to Mitogenomes. *Mol. Biol. Evol.* **35**, 299–311 (2018). [doi:10.1093/molbev/msx267](https://doi.org/10.1093/molbev/msx267) Medline
170. M. Bodner, U. A. Perego, G. Huber, L. Fendt, A. W. Röck, B. Zimmermann, A. Olivieri, A. Gómez-Carballa, H. Lancioni, N. Angerhofer, M. C. Bobillo, D. Corach, S. R. Woodward, A. Salas, A. Achilli, A. Torroni, H.-J. Bandelt, W. Parson, Rapid coastal spread of First Americans: Novel insights from South America's Southern Cone mitochondrial genomes. *Genome Res.* **22**, 811–820 (2012). [doi:10.1101/gr.131722.111](https://doi.org/10.1101/gr.131722.111) Medline
171. U. A. Perego, A. Achilli, N. Angerhofer, M. Accetturo, M. Pala, A. Olivieri, B. Hooshiar Kashani, K. H. Ritchie, R. Scozzari, Q.-P. Kong, N. M. Myres, A. Salas, O. Semino, H.-J. Bandelt, S. R. Woodward, A. Torroni, Distinctive Paleo-Indian migration routes from Beringia marked by two rare mtDNA haplogroups. *Curr. Biol.* **19**, 1–8 (2009). [doi:10.1016/j.cub.2008.11.058](https://doi.org/10.1016/j.cub.2008.11.058) Medline
172. Y. Cui, J. Lindo, C. E. Hughes, J. W. Johnson, A. G. Hernandez, B. M. Kemp, J. Ma, R. Cunningham, B. Petzelt, J. Mitchell, D. Archer, J. S. Cybulski, R. S. Malhi, Ancient DNA analysis of mid-holocene individuals from the Northwest Coast of North America reveals different evolutionary paths for mitogenomes. *PLOS ONE* **8**, e66948 (2013). [doi:10.1371/journal.pone.0066948](https://doi.org/10.1371/journal.pone.0066948) Medline
173. J. C. Chatters, D. J. Kennett, Y. Asmerom, B. M. Kemp, V. Polyak, A. N. Blank, P. A. Beddows, E. Reinhardt, J. Arroyo-Cabral, D. A. Bolnick, R. S. Malhi, B. J. Culleton, P. L. Erreguerena, D. Rissolo, S. Morell-Hart, T. W. Stafford Jr., Late Pleistocene human skeleton and mtDNA link Paleoamericans and modern Native Americans. *Science* **344**, 750–754 (2014). [doi:10.1126/science.1252619](https://doi.org/10.1126/science.1252619) Medline
174. K. Prüfer, M. Meyer, Comment on "Late Pleistocene human skeleton and mtDNA link Paleoamericans and modern Native Americans". *Science* **347**, 835–835 (2015). [doi:10.1126/science.1260617](https://doi.org/10.1126/science.1260617) Medline
175. B. M. Kemp, J. Lindo, D. A. Bolnick, R. S. Malhi, J. C. Chatters, Response to Comment on "Late Pleistocene human skeleton and mtDNA link Paleoamericans and modern Native Americans". *Science* **347**, 835–835 (2015). [doi:10.1126/science.1261188](https://doi.org/10.1126/science.1261188) Medline
176. C. de la Fuente, J. Galimany, B. M. Kemp, K. Judd, O. Reyes, M. Moraga, Ancient marine hunter-gatherers from Patagonia and Tierra Del Fuego: Diversity and differentiation using uniparentally inherited genetic markers. *Am. J. Phys. Anthropol.* **158**, 719–729 (2015). [doi:10.1002/ajpa.22815](https://doi.org/10.1002/ajpa.22815) Medline
177. M. de Saint Pierre, C. M. Bravi, J. M. B. Motti, N. Fuku, M. Tanaka, E. Llop, S. L. Bonatto, M. Moraga, An alternative model for the early peopling of southern South America revealed by analyses of three mitochondrial DNA haplogroups. *PLOS ONE* **7**, e43486 (2012). [doi:10.1371/journal.pone.0043486](https://doi.org/10.1371/journal.pone.0043486) Medline
178. B. A. Potter, J. D. Irish, J. D. Reuther, H. J. McKinney, New insights into Eastern Beringian mortuary behavior: A terminal Pleistocene double infant burial at Upward Sun River. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 17060–17065 (2014). [doi:10.1073/pnas.1413131111](https://doi.org/10.1073/pnas.1413131111) Medline
179. J. C. Tackney, B. A. Potter, J. Raff, M. Powers, W. S. Watkins, D. Warner, J. D. Reuther, J. D. Irish, D. H. O'Rourke, Two contemporaneous mitogenomes from terminal Pleistocene burials in eastern Beringia. *Proc. Natl. Acad. Sci. U.S.A.* **112**,

- 13833–13838 (2015). doi:10.1073/pnas.1511903112 Medline
180. F. A. Kaestle, D. G. Smith, Ancient mitochondrial DNA evidence for prehistoric population movement: The Numic expansion. *Am. J. Phys. Anthropol.* **115**, 1–12 (2001). doi:10.1002/ajpa.1051 Medline
181. G. S. Cabana, K. Hunley, F. A. Kaestle, Population continuity or replacement? A novel computer simulation approach and its application to the Numic expansion (Western Great Basin, USA). *Am. J. Phys. Anthropol.* **135**, 438–447 (2008). doi:10.1002/ajpa.20764 Medline
182. K. Thangaraj, G. Chaubey, T. Kivisild, A. G. Reddy, V. K. Singh, A. A. Rasalkar, L. Singh, Reconstructing the origin of Andaman Islanders. *Science* **308**, 996–996 (2005). doi:10.1126/science.1109987 Medline
183. S. S. Barik, R. Sahani, B. V. R. Prasad, P. Endicott, M. Metspalu, B. N. Sarkar, S. Bhattacharya, P. C. H. Annapoorna, J. Sreenath, D. Sun, J. J. Sanchez, S. Y. W. Ho, A. Chandrasekar, V. R. Rao, Detailed mtDNA genotypes permit a reassessment of the settlement and population structure of the Andaman Islands. *Am. J. Phys. Anthropol.* **136**, 19–27 (2008). doi:10.1002/ajpa.20773 Medline
184. M. G. Palanichamy, S. Agrawal, Y. G. Yao, Q. P. Kong, C. Sun, F. Khan, T. K. Chaudhuri, Y. P. Zhang, Comment on “Reconstructing the origin of Andaman islanders”. *Science* **311**, 470, author reply 470 (2006). doi:10.1126/science.1120176 Medline
185. P. Endicott, M. Metspalu, C. Stringer, V. Macaulay, A. Cooper, J. J. Sanchez, Multiplexed SNP typing of ancient DNA clarifies the origin of Andaman mtDNA haplogroups amongst South Asian tribal populations. *PLOS ONE* **1**, e81 (2006). doi:10.1371/journal.pone.0000081 Medline
186. H.-W. Wang, B. Mitra, T. K. Chaudhuri, M. G. Palanichamy, Q.-P. Kong, Y.-P. Zhang, Mitochondrial DNA evidence supports northeast Indian origin of the aboriginal Andamanese in the Late Paleolithic. *J. Genet. Genomics* **38**, 117–122 (2011). doi:10.1016/j.jgg.2011.02.005 Medline
187. P. Marrero, K. K. Abu-Amro, J. M. Larruga, V. M. Cabrera, Carriers of human mitochondrial DNA macrohaplogroup M colonized India from southeastern Asia. *BMC Evol. Biol.* **16**, 246 (2016). doi:10.1186/s12862-016-0816-8 Medline
188. M. Mondal, F. Casals, T. Xu, G. M. Dall’Olio, M. Pybus, M. G. Netea, D. Comas, H. Laayouni, Q. Li, P. P. Majumder, J. Bertranpetti, Genomic analysis of Andamanese provides insights into ancient human migration into Asia and adaptation. *Nat. Genet.* **48**, 1066–1070 (2016). doi:10.1038/ng.3621 Medline
189. M. Sikora et al., The population history of northeastern Siberia since the Pleistocene. *bioRxiv* 448829 [Preprint]. 22 October 2018. <https://doi.org/10.1101/448829>.
190. S. Mallick, H. Li, M. Lipson, I. Mathieson, M. Gymrek, F. Racimo, M. Zhao, N. Chennagiri, S. Nordenfelt, A. Tandon, P. Skoglund, I. Lazaridis, S. Sankararaman, Q. Fu, N. Rohland, G. Renaud, Y. Erlich, T. Willemse, C. Gallo, J. P. Spence, Y. S. Song, G. Poletti, F. Balloux, G. van Driem, P. de Knijff, I. G. Romero, A. R. Jha, D. M. Behar, C. M. Bravi, C. Capelli, T. Hervig, A. Moreno-Estrada, O. L. Posukh, E. Balanovskaya, O. Balanovsky, S. Karachanak-Yankova, H. Sahakyan, D. Toncheva, L. Yepiskoposyan, C. Tyler-Smith, Y. Xue, M. S. Abdullah, A. Ruiz-Linares, C. M. Beall, A. Di Rienzo, C. Jeong, E. B. Starikovskaya, E. Metspalu, J. Parik, R. Villem, B. M. Henn, U. Hodoglugil, R. Mahley, A. Sajantila, G. Stamatoyannopoulos, J. T. S. Wee, R. Khusainova, E. Khusnutdinova, S. Litvinov, G. Ayodo, D. Comas, M. F. Hammer, T. Kivisild, W. Klitz, C. A. Winkler, D. Labuda, M. Bamshad, L. B. Jorde, S. A. Tishkoff, W. S. Watkins, M. Metspalu, S. Dryomov, R. Sukernik, L. Singh, K. Thangaraj, S. Pääbo, J. Kelso, N. Patterson, D. Reich, The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* **538**, 201–206 (2016). doi:10.1038/nature18964 Medline
191. M. A. Yang, X. Gao, C. Theunert, H. Tong, A. Aximu-Petri, B. Nickel, M. Slatkin, M. Meyer, S. Pääbo, J. Kelso, Q. Fu, 40,000-Year-Old Individual from Asia Provides Insight into Early Population Structure in Eurasia. *Curr. Biol.* **27**, 3202–3208.e9 (2017). doi:10.1016/j.cub.2017.09.030 Medline
192. H. McColl, F. Racimo, L. Vinner, F. Demeter, T. Gakuhari, J. V. Moreno-Mayar, G. van Driem, U. Gram Wilken, A. Seguin-Orlando, C. de la Fuente Castro, S. Wasef, R. Shoocongdej, V. Souksavatdy, T. Sayavongkhamdy, M. M. Saidin, M. E. Allentoft, T. Sato, A.-S. Malaspina, F. A. Aghakhanian, T. Korneliussen, A. Prohaska, A. Margaryan, P. de Barros Damgaard, S. Kaewsutthi, P. Lertrit, T. M. H. Nguyen, H. C. Hung, T. Minh Tran, H. Nghia Truong, G. H. Nguyen, S. Shahidan, K. Wiradnyana, H. Matsumae, N. Shigehara, M. Yoneda, H. Ishida, T. Masuyama, Y. Yamada, A. Tajima, H. Shibata, A. Toyoda, T. Hanihara, S. Nakagome, T. Deviese, A.-M. Bacon, P. Duringer, J.-L. Ponche, L. Shackelford, E. Patole-Edoumba, A. T. Nguyen, B. Bellina-Pryce, J.-C. Galipaud, R. Kinaston, H. Buckley, C. Pottier, S. Rasmussen, T. Higham, R. A. Foley, M. M. Lahr, L. Orlando, M. Sikora, M. E. Phipps, H. Oota, C. Higham, D. M. Lambert, E. Willerslev, The prehistoric peopling of Southeast Asia. *Science* **361**, 88–92 (2018). doi:10.1126/science.aat3628 Medline
193. E. R. Jones, G. Gonzalez-Fortes, S. Connell, V. Siska, A. Eriksson, R. Martiniano, R. L. McLaughlin, M. Gallego Llorente, L. M. Cassidy, C. Gamba, T. Meshveliani, O. Bar-Yosef, W. Müller, A. Belfer-Cohen, Z. Matskevich, N. Jakeli, T. F. G. Higham, M. Currat, D. Lordkipanidze, M. Hofreiter, A. Manica, R. Pinhasi, D. G. Bradley, Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nat. Commun.* **6**, 8912 (2015). doi:10.1038/ncomms9912 Medline
194. C. Gamba, E. R. Jones, M. D. Teasdale, R. L. McLaughlin, G. Gonzalez-Fortes, V. Mattiangeli, L. Domboróczki, I. Kovári, I. Pap, A. Anders, A. Whittle, J. Dani, P. Raczyk, T. F. G. Higham, M. Hofreiter, D. G. Bradley, R. Pinhasi, Genome flux and stasis in a five millennium transect of European prehistory. *Nat. Commun.* **5**, 5257 (2014). doi:10.1038/ncomms6257 Medline
195. M. Sikora, A. Seguin-Orlando, V. C. Sousa, A. Albrechtsen, T. Korneliussen, A. Ko, S. Rasmussen, I. Duplanloup, P. R. Nigst, M. D. Bosch, G. Renaud, M. E. Allentoft, A. Margaryan, S. V. Vasilyev, E. V. Veselovskaya, S. B. Borutskaya, T. Deviese, D. Comeskey, T. Higham, A. Manica, R. Foley, D. J. Meltzer, R. Nielsen, L. Excoffier, M. Mirazon Lahr, L. Orlando, E. Willerslev, Ancient genomes show social and reproductive behavior of early Upper Paleolithic foragers. *Science* **358**, 659–662 (2017). doi:10.1126/science.aad1807 Medline
196. G. M. Kılıç, A. Omrak, F. Özer, T. Günther, A. M. Büyükkarakaya, E. Biçakçı, D. Baird, H. M. Dönertaş, A. Ghalichi, R. Yaka, D. Koptekin, S. C. Acan, P. Parviz, M. Krzewińska, E. A. Daskalaki, E. Yüncü, N. D. Dağtaş, A. Fairbaire, J. Pearson, G. Mustafaoğlu, Y. S. Erdal, Y. G. Çakan, İ. Togan, M. Somel, J. Storå, M. Jakobsson, A. Götherström, The Demographic Development of the First Farmers in Anatolia. *Curr. Biol.* **26**, 2659–2666 (2016). doi:10.1016/j.cub.2016.07.057 Medline
197. Z. Hofmanová, S. Kreutzer, G. Hellenthal, C. Sell, Y. Diekmann, D. Díez-Del-Molino, L. van Dorp, S. López, A. Kousathanas, V. Link, K. Kirsanow, L. M. Cassidy, R. Martiniano, M. Strobel, A. Scheu, K. Kotsakis, P. Halstead, S. Triantaphyllou, N. Kyparissi-Apostolika, D. Urem-Kotsou, C. Ziota, F. Adaktyiou, S. Gopalan, D. M. Bobo, L. Winkelbach, J. Blöcher, M. Unterländer, C. Leuenberger, Ç. Çilingiroğlu, B. Horejs, F. Gerritsen, S. J. Shennan, D. G. Bradley, M. Currat, K. R. Veeramah, D. Wegmann, M. G. Thomas, C. Papageorgopoulou, J. Burger, Early farmers from across Europe directly descended from Neolithic Aegeans. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 6886–6891 (2016). doi:10.1073/pnas.1523951113 Medline
198. F. Broushaki, M. G. Thomas, V. Link, S. López, L. van Dorp, K. Kirsanow, Z. Hofmanová, Y. Diekmann, L. M. Cassidy, D. Díez-Del-Molino, A. Kousathanas, C. Sell, H. H. Robson, R. Martiniano, J. Blöcher, A. Scheu, S. Kreutzer, R. Bollongino, D. Bobo, H. Davudi, O. Munoz, M. Currat, K. Abdi, F. Biglari, O. E. Craig, D. G. Bradley, S. Shennan, K. Veeramah, M. Mashkour, D. Wegmann, G. Hellenthal, J. Burger, Early Neolithic genomes from the eastern Fertile Crescent. *Science* **353**, 499–503 (2016). doi:10.1126/science.aaf7943 Medline
199. E. H. M. Wong, A. Khrunin, L. Nichols, D. Pushkarev, D. Khokhrin, D. Verbenko, O. Evgrafov, J. Knowles, J. Novembre, S. Limborska, A. Valouev, Reconstructing genetic history of Siberian and Northeastern European populations. *Genome Res.* **27**, 1–14 (2017). doi:10.1101/gr.202945.115 Medline
200. Q. Ayub, M. Mezzavilla, L. Pagani, M. Haber, A. Mohyuddin, S. Khaliq, S. Q. Mehdi, C. Tyler-Smith, The Kalash genetic isolate: Ancient divergence, drift, and selection. *Am. J. Hum. Genet.* **96**, 775–783 (2015). doi:10.1016/j.ajhg.2015.03.012 Medline
201. P. B. Damgaard, N. Marchi, S. Rasmussen, M. Peyrot, G. Renaud, T. Korneliussen, J. V. Moreno-Mayar, M. W. Pedersen, A. Goldberg, E. Usmanova, N. Baimukhanov, V. Loman, L. Hedeager, A. G. Pedersen, K. Nielsen, G. Afanasiev, K. Akmatov, A. Aldashev, A. Alpaslan, G. Baimbetov, V. I. Bazaliskii, A. Beisenov, B. Boldbaatar, B. Boldgiv, C. Dorzhu, S. Ellingvag, D. Erdenebaatar, R. Dajani, E. Dmitriev, V. Evdokimov, K. M. Frei, A. Gromov, A. Goryachev, H. Hakonarson, T. Hegay, Z. Khachatryan, R. Khaskhanov, E. Kitov, A. Kolbina, T. Kubatbek, A. Kukushkin, I. Kukushkin, N. Lau, A. Margaryan, I. Merkyte, I. V. Mertz, V. K. Mertz, E. Mijiddorj, V. Moiyesev, G. Mukhtarova, B. Nurmukhanbetov, Z. Orozbekova, I. Panyushkina, K. Pieta, V. Smrkova, I. Shevniina, A. Logvin, K.-G. Sjögren, T. Štolcová, A. M. Taravella, K. Tashbaeva, A. Tkachev, T. Tulegenov, D. Voyakin, L. Yepiskoposyan, S. Undrakhbold, V. Varfolomeev, A. Weber, M. A. Wilson Sayres, N. Kradin, M. E. Allentoft, L. Orlando, R. Nielsen, M. Sikora, E. Heyer, K. Kristiansen, E. Willerslev,

- 137 ancient human genomes from across the Eurasian steppes. *Nature* **557**, 369–374 (2018). doi:10.1038/s41586-018-0094-2 Medline
202. K. Prüfer, F. Racimo, N. Patterson, F. Jay, S. Sankararaman, S. Sawyer, A. Heinze, G. Renaud, P. H. Sudmant, C. de Filippo, H. Li, S. Mallick, M. Dannemann, Q. Fu, M. Kircher, M. Kuhlwilm, M. Lachmann, M. Meyer, M. Ongyerth, M. Siebauer, C. Theunert, A. Tandon, P. Moorjani, J. Pickrell, J. C. Mullikin, S. H. Vohr, R. E. Green, I. Hellmann, P. L. F. Johnson, H. Blanche, H. Cann, J. O. Kitzman, J. Shendure, E. E. Eichler, E. S. Lein, T. E. Bakken, L. V. Golovanova, V. B. Doronichev, M. V. Shunkov, A. P. Derevianko, B. Viola, M. Slatkin, D. Reich, J. Kelso, S. Pääbo, The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature* **505**, 43–49 (2014). doi:10.1038/nature12886 Medline
203. M. G. Llorente, E. R. Jones, A. Eriksson, V. Siska, K. W. Arthur, J. W. Arthur, M. C. Curtis, J. T. Stock, M. Coltorti, P. Pieruccini, S. Stretton, F. Brock, T. Higham, Y. Park, M. Hofreiter, D. G. Bradley, J. Bhak, R. Pinhasi, A. Manica, Ancient Ethiopian genome reveals extensive Eurasian admixture in Eastern Africa. *Science* **350**, 820–822 (2015). doi:10.1126/science.aad2879 Medline
204. I. Lazaridis, N. Patterson, A. Mittnik, G. Renaud, S. Mallick, K. Kirsanow, P. H. Sudmant, J. G. Schraiber, S. Castellano, M. Lipson, B. Berger, C. Economou, R. Bollongino, Q. Fu, K. I. Bos, S. Nordenfelt, H. Li, C. de Filippo, K. Prüfer, S. Sawyer, C. Posth, W. Haak, F. Hallgren, E. Fornander, N. Rohland, D. Delsate, M. Francken, J.-M. Guinet, J. Wahl, G. Ayodo, H. A. Babiker, G. Bailliet, E. Balanovska, O. Balanovsky, R. Barrantes, G. Bedoya, H. Ben-Ami, J. Bene, F. Berrada, C. M. Bravi, F. Brisighelli, G. B. J. Busby, F. Cali, M. Churnosov, D. E. C. Cole, D. Corach, L. Damba, G. van Driem, S. Dryomov, J.-M. Dugoujon, S. A. Fedorova, I. Gallego Romero, M. Gubina, M. Hammer, B. M. Henn, T. Hervig, U. Hodoglugil, A. R. Jha, S. Karachanak-Yankova, R. Khusainova, E. Khusnutdinova, R. Kittles, T. Kivisild, W. Klitz, V. Kučinskas, A. Kushniarevich, L. Laredj, S. Litvinov, T. Loukidis, R. W. Mahley, B. Melegh, E. Metspalu, J. Molina, J. Mountain, K. Näkkäläjärvi, D. Nesheva, T. Nyambo, L. Osipova, J. Parik, F. Platonov, O. Posukh, V. Romano, F. Rothhammer, I. Rudan, R. Ruizbakiev, H. Sahakyan, A. Sajantila, A. Salas, E. B. Starikovskaya, A. Tarekegn, D. Toncheva, S. Turdikulova, I. Uktvertveite, O. Utvenska, R. Vasquez, M. Villena, M. Voivedova, C. A. Winkler, L. Yepiskoposyan, P. Zalloua, T. Zemunik, A. Cooper, C. Capelli, M. G. Thomas, A. Ruiz-Linares, S. A. Tishkoff, L. Singh, K. Thangaraj, R. Villem, D. Comas, R. Sukernik, M. Metspalu, M. Meyer, E. E. Eichler, J. Burger, M. Slatkin, S. Pääbo, J. Kelso, D. Reich, J. Krause, Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature* **513**, 409–413 (2014). doi:10.1038/nature13673 Medline
205. S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham, PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007). doi:10.1086/519795 Medline
206. E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, Testing for ancient admixture between closely related populations. *Mol. Biol. Evol.* **28**, 2239–2252 (2011). doi:10.1093/molbev/msr048 Medline
207. B. M. Peter, Admixture, Population Structure, and F-Statistics. *Genetics* **202**, 1485–1501 (2016). doi:10.1534/genetics.115.183913 Medline
208. R. E. Green, J. Krause, A. W. Briggs, T. Maricic, U. Stenzel, M. Kircher, N. Patterson, H. Li, W. Zhai, M. H. Y. Fritz, N. F. Hansen, E. Y. Durand, A. S. Malaspina, J. D. Jensen, T. Marques-Bonet, C. Alkan, K. Prüfer, M. Meyer, H. A. Burbano, J. M. Good, R. Schultz, A. Aximu-Petri, A. Butthof, B. Höber, B. Höffner, M. Siegemund, A. Weihmann, C. Nusbaum, E. S. Lander, C. Russ, N. Novod, J. Affourtit, M. Egholm, C. Verna, P. Rudan, D. Brajkovic, Ž. Kucan, I. Gušić, V. B. Doronichev, L. V. Golovanova, C. Lalucea-Fox, M. de la Rasilla, J. Fortea, A. Rosas, R. W. Schmitz, P. L. F. Johnson, E. E. Eichler, D. Falush, E. Birney, J. C. Mullikin, M. Slatkin, R. Nielsen, J. Kelso, M. Lachmann, D. Reich, S. Pääbo, A draft sequence of the Neandertal genome. *Science* **328**, 710–722 (2010). doi:10.1126/science.1188021 Medline
209. J. Kelleher, A. M. Etheridge, G. McVean, Efficient coalescent simulation and genealogical analysis for large sample sizes. *PLOS Comput. Biol.* **12**, e1004842 (2016). doi:10.1371/journal.pcbi.1004842 Medline
210. J. V. Moreno-Mayar, S. Rasmussen, A. Seguin-Orlando, M. Rasmussen, M. Liang, S. T. Flåm, B. A. Lie, G. D. Gilfillan, R. Nielsen, E. Thorsby, E. Willerslev, A.-S. Malaspina, Genome-wide ancestry patterns in Rapanui suggest pre-European admixture with Native Americans. *Curr. Biol.* **24**, 2518–2525 (2014). doi:10.1016/j.cub.2014.09.057 Medline
211. A. Kong, M. L. Frigge, G. Masson, S. Besenbacher, P. Sulem, G. Magnusson, S. A. Gudjonsson, A. Sigurdsson, A. Jonasdottir, A. Jonasdottir, W. S. W. Wong, G. Sigurdsson, G. B. Walters, S. Steinberg, H. Helgason, G. Thorleifsson, D. F. Gudbjartsson, A. Helgason, O. T. Magnusson, U. Thorsteinsdottir, K. Stefansson, Rate of de novo mutations and the importance of father's age to disease risk. *Nature* **488**, 471–475 (2012). doi:10.1038/nature11396 Medline
212. H. Jónsson, P. Sulem, B. Kehr, S. Kristmundsdottir, F. Zink, E. Hjartarson, M. T. Hardarson, K. E. Hjorleifsson, H. P. Eggertsson, S. A. Gudjonsson, L. D. Ward, G. A. Arnadottir, E. A. Helgason, H. Helgason, A. Gylfason, A. Jonasdottir, A. Jonasdottir, T. Rafnar, M. Frigge, S. N. Stacey, O. Th Magnusson, U. Thorsteinsdottir, G. Masson, A. Kong, B. V. Halldorsson, A. Helgason, D. F. Gudbjartsson, K. Stefansson, Parental influence on human germline de novo mutations in 1,548 trios from Iceland. *Nature* **549**, 519–522 (2017). doi:10.1038/nature24018 Medline
213. T. B. Coplen, Reporting of stable hydrogen, carbon, and oxygen isotopic abundances (Technical Report). *Pure Appl. Chem.* **66**, 273–276 (1994). doi:10.1351/pac199466020273
214. M. Stuiver, H. A. Polach, Discussion Reporting of 14C Data. *Radiocarbon* **19**, 355–363 (1977). doi:10.1017/S003382200003672
215. F. Mizuno, J. Gojobori, L. Wang, K. Onishi, S. Sugiyama, J. Granados, C. Gomez-Trejo, V. Acuña-Alonso, S. Ueda, Complete mitogenome analysis of indigenous populations in Mexico: Its relevance for the origin of Mesoamericans. *J. Hum. Genet.* **59**, 359–367 (2014). doi:10.1038/jhg.2014.35 Medline
216. S. Kumar, C. Bellis, M. Zlotyutro, P. E. Melton, J. Blangero, J. E. Curran, Large scale mitochondrial sequencing in Mexican Americans suggests a reappraisal of Native American origins. *BMC Evol. Biol.* **11**, 293 (2011). doi:10.1186/1471-2148-11-293 Medline
217. C. M. Lewis Jr., B. Lizárraga, R. Y. Tito, P. W. López, G. C. Iannaccone, A. Medina, R. Martínez, S. I. Polo, A. F. De La Cruz, A. M. Cáceres, A. C. Stone, Mitochondrial DNA and the peopling of South America. *Hum. Biol.* **79**, 159–178 (2007). doi:10.1353/hub.2007.0031 Medline
218. E. J. Lee, D. A. Merriwether, Identification of Whole Mitochondrial Genomes from Venezuela and Implications on Regional Phylogenies in South America. *Hum. Biol.* **87**, 29–38 (2015). doi:10.13110/humanbiology.87.1.0029 Medline
219. L. Arias, C. Barbieri, G. Barreto, M. Stoneking, B. Pakendorf, High-resolution mitochondrial DNA analysis sheds light on human diversity, cultural interactions, and population mobility in Northwestern Amazonia. *Am. J. Phys. Anthropol.* **165**, 238–255 (2018). doi:10.1002/ajpa.23345 Medline
220. R. S. Just, T. M. Diegoli, J. L. Saunier, J. A. Irwin, T. J. Parsons, Complete mitochondrial genome sequences for 265 African American and U.S. "Hispanic" individuals. *Forensic Sci. Int. Genet.* **2**, e45–e48 (2008). doi:10.1016/j.fsigen.2007.12.001 Medline
221. N. J. R. Fagundes, R. Kanitz, R. Eckert, A. C. S. Valls, M. R. Bogo, F. M. Salzano, D. G. Smith, W. A. Silva Jr., M. A. Zago, A. K. Ribeiro-dos-Santos, S. E. B. Santos, M. L. Petzl-Erler, S. L. Bonatto, Mitochondrial population genomics supports a single pre-Clovis origin with a coastal route for the peopling of the Americas. *Am. J. Hum. Genet.* **82**, 583–592 (2008). doi:10.1016/j.ajhg.2007.11.013 Medline
222. A. Hartmann, M. Thieme, L. K. Nanduri, T. Stempfli, C. Moehle, T. Kivisild, P. J. Oefner, Validation of microarray-based resequencing of 93 worldwide mitochondrial genomes. *Hum. Mutat.* **30**, 115–122 (2009). doi:10.1002/humu.20816 Medline
223. D. M. Behar, M. van Oven, S. Rosset, M. Metspalu, E.-L. Loogväli, N. M. Silva, T. Kivisild, A. Torroni, R. Villems, A "Copernican" reassessment of the human mitochondrial DNA tree from its root. *Am. J. Hum. Genet.* **90**, 675–684 (2012). doi:10.1016/j.ajhg.2012.03.002 Medline

## ACKNOWLEDGMENTS

For permission and cooperation in providing human remains for this study, we thank the Native Village of Deering, Alaska; the Fallon Paiute-Shoshone Tribe; the Consejo de Monumentos Nacionales, Chile; and the Canadian Museum of History and the Stswecem'c Xgat'tem, Metlakatla, and Lax Kw'alaams First Nations. See (74, 75) for related analyses of ancient DNA samples from the Americas. We thank J. Ramos-Madrigal, L. Orlando, G. Renaud, F. Vieira, L. Benson, A. Bergström, T. Ferraz, and J. E. Santos-Júnior for discussions; M. Drummond, F. Zimmermann, and T. Tisler for image processing; B. Chestnut, B. Baldwin, and D. Cossette for

help and support; and the Danish National High-throughput Sequencing Centre for assistance with data generation. St. John's College, Cambridge University, provided E.W. and D.J.M. a congenial venue for scientific discussions. **Funding:** GeoGenetics members were supported by the Lundbeck Foundation and the Danish National Research Foundation (DNRF94) and KU2016. Study of the Lagoa Santa human remains was partly funded by the Augustinus Foundation. J.C. is supported in part by an NSF graduate research fellowship. J.C., J.P.S., and Y.S.S. are supported in part by the National Institutes of Health grant R01-GM094402, a Microsoft Azure research award, and a Packard Fellowship for Science and Engineering. A.S. was funded by FAPESP (2017/16451-2). Study of Patagonian samples by M.A.P.D. was sponsored by the National Geographic Society, Genographic Program, grant 14-01. H.S. was supported by HERA and the European Union's Horizon2020 Research and Innovation Programme (grant 649307). J.R.R. was supported by funding from the National Park Service Shared Beringian Heritage Program. A.S.M. thanks the Swiss National Science Foundation and the European Research Council (ERC starting grant) for funding. Y.S.S. is a Chan Zuckerberg Biohub Investigator. D.J.M. is supported by the Quest Archaeological Research Fund and the Potts and Sibley Foundation. **Author contributions:** The project was conceived by E.W., J.V.M.-M., and D.J.M. and headed by E.W. L.V., P.B.D., C.F., M.E.A., A.M., M.I.O., D.M., and H.S. processed ancient DNA. J.V.M.-M., S.R., and T.P. assembled datasets. J.V.M.-M., J.C., J.P.S., T.V., F.R., T.P., and Y.S.S. analyzed genetic data with input from P.B.D. and C.F. R.N., Y.S.S., M.Si., and A.-S.M. supervised bioinformatic and statistical analyses. Radiocarbon analyses were done by L.B.-V., D.Ch., D.Co., T.D., and T.H. C.V., F.M., M.V., S.W., K.G., N.L., M.M.L., K.K., and L.S. excavated, curated, sampled, or described human remains. D.J.M., D.K.G., V.A., C.P., J.E., C.R.-C., S.R., M.Q.R.B., J.C., M.W., C.B., K.L.H., M.M.L., A.St., A.Sa., F.M., M.A.-D., and J.T.R. conducted archaeological and bioanthropological analyses and provided contextualization. L.G. and H.H. partnered with R.M. and E.W. to engage with indigenous communities. All authors contributed to data interpretation. J.V.M.-M., D.J.M., Y.S.S., and E.W. wrote the manuscript, with input from J.C., J.P.S., M.A.-D., R.M., J.T.R., T.P., F.R.S., A.-S.M., and the other authors. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** Sequence data are available at the ENA under accession PRJEB29074.

## SUPPLEMENTARY MATERIALS

[www.sciencemag.org/cgi/content/full/science.aav2621/DC1](http://www.sciencemag.org/cgi/content/full/science.aav2621/DC1)

Materials and Methods

Supplementary Text

Figs. S1 to S80


Tables S1 to S18

References (76–223)


3 September 2018; accepted 30 October 2018

Published online 8 November 2018


10.1126/science.aav2621






**Fig. 1. Ancient genome overview and their broad genetic affinities.** (A) Sampling locations for ancient genomes (circles; new ones in bold) and present-day Native Americans [triangles colored by the grouping used in (2, 4)]. NNA and SNA were classified following (1). Present-day whole-genome data are labeled in dark blue. Broad geographic features mentioned in the text are shown in brown; the extent of glacial ice sheets ~15.5 ka (73) in light blue. (B) Calibrated radiocarbon ages for ancient genomes; previously published (open circles), this study (filled circles with depth of coverage). (C) Multidimensional scaling plot from the  $f_3$ -distance matrix computed from a subset of the SNP array dataset (~200,000 sites), including Siberian and Native American populations. (D and E) MDS plots similar to C, showing the first three dimensions for SNA groups only. (F) ADMIXTURE proportions assuming  $K = 16$  ancestral populations. Bars represent individuals; colors represent ancestral components. For clarity, we only show Native Americans, three individuals for populations with  $n > 3$ , and single genomes are represented as wider bars. Siberians and Native Americans are organized according to (4).




**Fig. 2. Admixture graphs modelling the ancestry of ancient North American genomes.** We enumerated all possible extensions of the seed graph (13) where we added Trail Creek (**A**), Big Bar (**B**), 939 (**C**), Kennewick (**D**), and ASO (**E**) as a “non-admixed” or an admixed population, and optimized the parameters for each topology using qpGraph. In each graph, the test population is shown in blue. We show the best-fitting model for each genome as inferred from the final fit score. Above each graph, we show: the four populations leading to the worst  $D$ -statistic residual; the observed value for such statistic, and the expected values under the fitted model, the residual, the standard error of the residuals and a Z-score for such a residual; and the model fit score. Numbers to the right of solid edges are proportional to optimized drift; percentages to the right of dashed edges represent admixture proportions. In each graph we highlight the test genome in blue. (**F**) Error-corrected  $D$ -statistics restricted to transversion polymorphisms testing the genetic affinity between ASO and different SNA pairs. Points represent  $D$ -statistics and error bars represent  $\sim 3.3$  standard errors ( $P \sim 0.001$ ). For each test, we show the absolute Z-score beside its corresponding  $D$  value. A pool of the five sequenced individuals represents the Lagoa Santa population.




**Fig. 3. *f*-statistics-based tests show a rapid dispersal into South America, followed by Mesoamerican-related admixture.** (A) Schematic representation of a model for SNA formation. This model represents a reasonable fit to most present-day populations (13). “UPopA”-, Mixe- and Australasian-related admixture edges are color-coded as in (C) to (F). (B) Fit score of the graph shown in (A) (excluding South Americans), as a function of “unsampled admixture” in Mixe. The point indicates the “unsampled admixture” proportion that yields the best fit score. (C) Error-corrected *D*-statistics showing that Lagoa Santa, Mixe and most SNA genomes cannot be modeled using a simple tree. In the top panel, we show the tested null hypothesis, together with an indication of the pair of populations with excess allele sharing, depending on the sign of *D*. SNA populations are organized according to their sampling location (right labels). Points represent *D*-statistics and error bars represent ~3.3 standard errors (which corresponds to a *P* value of ~0.001 in a Z-test). For each test, we show the absolute *Z*-score beside its corresponding *D* value. (D and E) Fit score surfaces for the “admixed” SNA model with fixed Mixe and Australasian admixture proportions. For Ayayema and Suruí, we explored the fit of the model shown in a., across a grid of values for the Mixe proportion into SNA {0,0.05,...,1} and the Australasian contribution into Lagoa Santa {0,0.01,...,0.1}. “X” indicates the parameter combination yielding the best score. Contour lines were drawn such that all parameter combinations contained within a given line, all yield a fit score lower than that indicated by the contour label. (F) A one-dimensional representation of (D) and (E) for all SNA populations. In this case, we fixed the Australasian contribution into Lagoa Santa to 3%. Each line is labeled at the value that yields the best fit score. We compare different models based on their fit scores, where a difference of ~3 corresponds to a *P* value of ~0.05 and a difference of ~4.6 corresponds to a *P* value of ~0.01. For (B) a pool of the five sequenced individuals represents the Lagoa Santa population. For (C) to (F), we considered the called-genotype dataset excluding transitions (13), and used the high-depth “Sumidouro5” individual as a representative of the Lagoa Santa population.



**Fig. 4. Allele sharing symmetry tests for pairs of Native Americans, relative to present-day Eurasian groups.** (A) We computed  $D$ -statistics of the form  $D(\text{Native American}, \text{Native American}; \text{Eurasian}, \text{Yoruba})$  to test whether a given Native American group carries excess “non-Native American” ancestry compared to other Native Americans. For each statistic, we obtained a Z-score (diamonds) based on a weighted block-jackknife procedure over 5-Mb blocks. Vertical lines represent  $\sim 3.3$  and  $\sim -3.3$  (which correspond to a  $P$  value of  $\sim 0.001$ ). Purple: Oceanians; pink: Southeast Asians; gray: non-Australasians. In this case we only show results for present-day Eurasian populations (13). (B) “Contamination-corrected”  $f_4$ -statistics of the form  $f_4(\text{Mixe}, \text{Lagoa Santa}; \text{Australasian}, \text{Yoruba})$ . For each statistic, we subtracted the value of  $f_4(\text{Mixe}, \text{French}; \text{Australasian}, \text{Yoruba})$ , weighted by an assumed contamination fraction  $c$  ranging between 0 and 10% (y-axis). Points represent  $f_4$ -statistics and error bars represent  $\sim 3.3$  standard errors. We observe that the apparent allele sharing between Lagoa Santa and Australasians increases as a function of the correction. As a reference, we show the value of  $f_4(\text{Mixe}, \text{Surui}; \text{Australasian}, \text{Yoruba})$  as solid vertical lines. All tests are from the whole-genome dataset described in (13) and excluding transition polymorphisms. (C) Approximate sampling location for Australasian groups highlighted in (A).



**Fig. 5. Demographic history of South Native Americans.** A schematic representation of the most likely model relating the ancient USR1, Anzick1, Spirit Cave and Lagoa Santa genomes and the present-day Mixe ( $n = 3$ ) and Karitiana ( $n = 5$ ). Demographic parameters were inferred using momi2 (13). This model features a quick north to south splitting pattern of South Native Americans over a period shorter than 2,000 years, with later admixture from an outgroup ("UPopA") into the Mesoamerican Mixe. In addition we find evidence of gene flow from the latter into present-day South Americans, represented in this case by the Karitiana. Admixture pulses from USR1 into the ancestors of other Native Americans follow the inference in (1).



**Fig. 6. Schematic depiction of the processes of human dispersal and divergence in the Americas, arranged chronologically.** (A) Initial entry into eastern Beringia then into unglaciated North America, ~25 ka to ~13 ka, during which there are multiple splits: first in Beringia (UPopA and AB from the NNA+SNA line), followed by the Big Bar ancestral population splitting from NNA+SNA, then lastly by the NNA/SNA split south of eastern Beringia. NNA groups remained in northern North America, while SNA groups began to disperse across the North American continent. (B) Period of dispersal hemisphere-wide, ~14 ka to ~6 ka, during which time SNA moved rapidly from North into South America, resulting in the close affinities of the nearly contemporaneous Spirit Cave and Lagoa Santa individuals. Early South American populations possibly carried an Australasian-related admixture, as seen in the Lagoa Santa individual, and diverged west and east of the Andes. There was also admixture in North America between NNA and SNA groups prior to 9 ka that formed the population of which Kennewick/Ancient One was a member. It is inferred that during this period but after 9 ka (the age of the Trail Creek AB individual) NNA groups moved north into Alaska. (C) Population expansion out of Mesoamerica sometime after ~8.7 ka. These groups moved north into the Great Basin, resulting in a population turnover after 2 ka, evidenced by the difference between the Lovelock 2 and Lovelock 3 individuals. In South America that expansion contributed to the ancestry of most South American groups, but did not reach Patagonia by 5.1 ka, the age of the Ayayema individual.

## Early human dispersals within the Americas

J. Víctor Moreno-Mayar, Lasse Vinner, Peter de Barros Damgaard, Constanza de la Fuente, Jeffrey Chan, Jeffrey P. Spence, Morten E. Allentoft, Tharsika Vimala, Fernando Racimo, Thomaz Pinotti, Simon Rasmussen, Ashot Margaryan, Miren Iraeta Orbegozo, Dorothea Mylopotamitaki, Matthew Wooller, Clement Bataille, Lorena Becerra-Valdivia, David Chivall, Daniel Comeskey, Thibaut Devèze, Donald K. Grayson, Len George, Harold Harry, Verner AlexanderSEN, Charlotte Primeau, Jon Erlandson, Claudia Rodrigues-Carvalho, Silvia Reis, Murilo Q. R. Bastos, Jerome Cybulski, Carlos Vullo, Flavia Morello, Miguel Vilar, Spencer Wells, Kristian Gregersen, Kasper Lykke Hansen, Niels Lynnerup, Marta Mirazón Lahr, Kurt Kjær, André Strauss, Marta Alfonso-Durruy, Antonio Salas, Hannes Schroeder, Thomas Higham, Ripan S. Malhi, Jeffrey T. Rasic, Luiz Souza, Fabricio R. Santos, Anna-Sapfo MalaspinaS, Martin Sikora, Rasmus Nielsen, Yun S. Song, David J. Meltzer and Eske Willerslev

published online November 8, 2018

### ARTICLE TOOLS

<http://science.scienmag.org/content/early/2018/11/07/science.aav2621>

### SUPPLEMENTARY MATERIALS

<http://science.scienmag.org/content/suppl/2018/11/07/science.aav2621.DC1>

### RELATED CONTENT

<http://science.scienmag.org/content/sci/362/6415/627.full>

### REFERENCES

This article cites 199 articles, 49 of which you can access for free

<http://science.scienmag.org/content/early/2018/11/07/science.aav2621#BIBL>

### PERMISSIONS

<http://www.scienmag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)