

# Anemone Coronaria

## MA Practicum

Perach Hen Elkayam

### Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Background</b>	<b>2</b>
<b>3</b>	<b>The Data</b>	<b>4</b>
<b>4</b>	<b>Methodology</b>	<b>5</b>
<b>5</b>	<b>Results</b>	<b>6</b>
5.1	Whether the herbivores chose to eat the leaves because of the color of the flower	6
5.2	Whether the florivors chose to eat the flower because of the color of the flower .	7
5.3	Whether the herbivory reduces the weight of seeds . . . . .	8
5.4	Whether the florivory reduces the number of seeds . . . . .	9
<b>6</b>	<b>Conclusions</b>	<b>10</b>
<b>7</b>	<b>Limitations</b>	<b>10</b>
<b>8</b>	<b>Appendix</b>	<b>11</b>
8.1	Sampling protocol: . . . . .	11
8.2	Variables Description . . . . .	12
8.3	Additional Models for the Rate of Leaf Damage (Herbivory) . . . . .	12
8.4	Additional Models for the Rate of Petals Damage (Florivory) . . . . .	13
8.5	Code and Data . . . . .	13
	<b>Reference</b>	<b>14</b>

# 1 Introduction

This project focus on an analysis of *Anemone coronaria* data that was collected by *Norah Saabnah* from the *School of Plant Sciences and Food Security*.

The following research questions will deal with the roles of florivory and herbivory in maintaining flower color variation in *Anemone coronaria*. The statistical questions are:

1. Whether the herbivores chose to eat the leaves because of the color of the flower
2. Whether the florivors chose to eat the flower because of the color of the flower
3. Whether the herbivory reduces the weight of seeds
4. Whether the florivory reduces the number of seeds

# 2 Background

*Anemone coronaria* is a winter flowering species that is dormant during the dry season and regenerates annually from subterranean buds on a corm. Each flower blooms for a few weeks in mid-winter. The flowers open and close according to the sunshine and other weather condition. *Anemone coronaria* displays a wide and heritable diversity of floral colors from red through purple and blue to white, with high variation within and among populations (EHRENDORFER et al. 2009). While red flowers are abundant throughout the whole distribution of the species, non-red flowers (white, pink, purple or blue) are present only in the Mediterranean climate region. We can also see that in the collected data we have mostly red *Anemone coronaria* (in Sarid and Shokeda all the flowers are red).




Figure 1: Flower color by site

Plants attract pollinators by presenting flowers in a distinct color (Kooi et al. 2016). Color vision is widespread among pollinators but varies among species and depends on the spectral sensitivities of the photoreceptors and their interplay (Kooi et al. 2021). Consequently, much research of flower color variation explored its consequences for interactions with pollinators. Balancing selection by pollinators and herbivores is thought to be an important mechanism maintaining variation in flower color. Flower color morphs that are preferred by pollinators and are also susceptible to herbivores, may undergo a conflicting selection pressure on flower color (Sapir, Gallagher, and Senden 2021). Herbivory, the consumption of plant tissues by an animal, can occur on leaves (herbivory) or flowers (florivory). From here on, we refer to leaf damage as *herbivory*, and flower damage as *florivory*.

Glaphyrid beetles are the major pollinators of *Anemone coronaria*, as well as one of the major florivores on *Anemone coronaria*, eating petals and pollen (Dafni et al. 1990). Previous studies showed that beetles see red flowers well, therefore visit red flower models more frequently than models of different colors (Martínez-Harms and Balvanera 2012). The distribution of *Anemone coronaria* in Israel is wide and it is highly abundant along the climate gradient.

A few characteristics make *Anemone coronaria* a good model plant: color is governed by a simple heredity system (Horovitz, Bullowa, and Negbi 1975), and the pollination system is relatively simple because the plant is self-incompatible and offers only pollen as reward for pollinators. For these reasons, and due to the prominent flower color variation, *Anemone coronaria* provides an excellent model system to understand the drivers of flower color variation and what maintains it.

### 3 The Data

The data consist of 60 field measurements in 4 different sites in Israel: Sarid, Megido, Beqoa, and Shokeda. The sites were chosen based on a preliminary survey of populations during 2020-2021 by Dr. Yuval Sapir and Prof. Tamar Keasar.

During the flowering season, the data was collected three times at each site: at the beginning of the flowering season (mid-January), in the middle of it (Early February), and at its end (late February/early March).

At each site, 30 flowers and their nearest neighbors were randomly chosen, and a total of 60 flowers were sampled at each site according to the [Protocol for Quantifying Variability in Plant-Herbivore Interactions](#) that is designed to deal with highly skewed distribution of plants damage (See Appendix for [detailed sampling protocol](#)). The flower color, weight and number of seeds, Flower sex, height, leaf damage, and petal damage were measured for each plant. The leaf damage and petal damage were measured by visually estimating the percentage of the area damaged according to [Herbvar online application](#).

In Figure 1 we can see that only in *Bequa* we have enough red and non-red flowers in order to make an analysis (in Sarid and Shokeda there are only red flowers, and in Megido there are only five red flowers). Including data from sites with imbalanced values of red and non-red flowers might lead to incorrect conclusions as the site variables play a role as confounders. Therefore, I will use only data from *Bequa* for the first two research questions. This data contains 60 observations.

See Appendix for [variable description](#).

#### 3.0.0.1 Preprocess

The data included some missing values. Observations with missing response variables were omitted from the analysis.

To apply Beta Regression I had to change some observations from 0 to 0.001 in the two response variables - the rate of the petal damage and the leaf damage:

Table 1: Number of values transformed from 0 to 0.001

Leaf	Petal
2/60	12/60

## 4 Methodology

I used Beta Regression (Cribari-Neto and Zeileis 2010) for the first two questions since the response variables (rate of damage of petals and leaves), are in the range of  $[0,1)$  and stand for rate. The main problem with this method is that there are some cases of 0s in the response variables. In order to overcome the problem, I manipulated the rate variables, as I mentioned in the [preprocess](#) section.

To ensure that the results are consistent I decided to try two additional methods: Linear Regression and Ordinal Regression that are more common and have better documentation than the Beta Regression. Both methods are not the ultimate solution for this kind of a problem since they are not designed for a rate outcome. Moreover the Ordinal Regression forced me to create an ordinal-categorical variable from the original outcome variable which leads to information loss.

For the last two question I used Linear Regression to model the weight and the number of seeds of the plant. In these research questions the choice was straight-forward since the outcome is continuous.

## 5 Results

### 5.1 Whether the herbivores chose to eat the leaves because of the color of the flower

The graph and the regression analysis show no evidence of the color's effect on the herbivory rate for every reasonable significance level. As can be seen in the appendix, the results from the [Linear Regression](#) and the [Categorical Regression](#) are consistent with the results of the beta regression (that is presented below).




Table 2: Beta Regression for Rate of Leaf Damage (Herbivory)

Coefficients	Estimate	P Value	
Intercept	-1.9004	0.0000	***
Color,Red	0.2049	0.4944	
Height	-0.0053	0.7383	
Flowersexual,Male	0.0476	0.8623	

## 5.2 Whether the florivores chose to eat the flower because of the color of the flower

According to the regression analysis, there is no evidence for the effect of the color over the florivory rate for every reasonable significance level. As can be seen in the appendix, the results from the [Linear Regression](#) and the [Categorical Regression](#) are consistent with the results of the beta regression (that is presented below).




Table 3: Beta Regression for Rate of Petals Damage (Florivory)

Coefficients	Estimate	P Value	
Intercept	-3.4081	0.0000	***
Color,Red	-0.0927	0.7390	
Height	0.0046	0.7540	
Flowersexual,Male	0.4927	0.0566	.

### 5.3 Whether the herbivory reduces the weight of seeds

According to the regression analysis, there is a significant negative effect of the rate of leaf damage (Herbivory) over the weight of the seeds for a significance level of 0.05 but not for a significance level of 0.01.




Table 4: Linear Model for Seed Weight (Fitness)

Coefficients	Estimate	P Value	
Intercept	0.1257	0.0000	***
Rate of Leaf Damage (Herbivory)	-0.2433	0.0131	*
Color,Red	0.0355	0.1048	
Height	0.0022	0.0222	*
Flowersexual,Male	-0.0297	0.0375	*
Site,Megido	-0.0788	0.0000	***
Site,Sarid	-0.0445	0.0710	.
Site,Shokeda	-0.0883	0.0030	**



## 5.4 Whether the florivory reduces the number of seeds

The regression analysis shows that there is a significant negative effect of the rate of petals damage (Florivory) over the number of seeds for a significance level of 0.1 but not for a significance level of 0.05.




Table 5: Linear Model for the Number of Seeds (Fitness)

Coefficients	Estimate	P Value	
Intercept	323.7180	0.0000	***
Rate of Petals Damage (Florivory)	-386.9033	0.0819	.
Color,Red	119.2225	0.0255	*
Height	4.5871	0.0445	*
Flowersexual,Male	-70.0309	0.0466	*
Site,Megido	-178.0583	0.0001	***
Site,Sarid	-249.0064	0.0000	***
Site,Shokeda	-205.3261	0.0043	**

## 6 Conclusions

In this report, I fitted several models to four research questions. For the first two questions - whether the herbivores or florivores chose to eat the leaves because of the color of the flower, I fitted Beta, Linear, and Ordinal Regression to observe whether I got consistent results. I found no evidence in all the models that the color of the flower affects the herbivory or the florivory rate.

For the two other questions - whether the herbivory reduces the weight of seeds and whether the florivory reduces the number of seeds, I fitted Linear Regression. I found that there is a moderate evidence that the rate of leaf damage (Herbivory) reduces the weight of the seeds and that there is a weak evidence that the rate of petals damage (Florivory) reduces the number of seeds.


## 7 Limitations

- The leaf and petal damage were measured by visually estimating the percentage of the area damaged - a subjective measure.
- Two first research questions are analyzed only with data from Bequa site. Recommendation for future research - collect a sufficient number of red flowers in Megido to use data from another site besides Bequa for the two first research questions.

## 8 Appendix

### 8.1 Sampling protocol:

- (1) Record plant density in 10 randomly located 1-m<sup>2</sup> areas to estimate average plant density, which is used to calculate quadrat radius.
- (2) A quadrat with one focal plant and its nearest neighbor (outside quadrat).
- (3) A quadrat with no focal plants.
- (4) A quadrat with 5 focal plants; plant 3 is randomly selected for data collection, and its nearest neighbor is plant 4. Diagram by Moria Robinson.



## 8.2 Variables Description

Name	Description
Site	Site name
ht_cm	Measured from ground to highest plant tissue
Flowercolor	Red, Purple, Pink, Pink-white and White
color	The flower color categorized as red or non-red
Flowersexual	Female or Male
weight_of_seeds	Weight of seeds
number_of_seeds	Number of seeds
rate_herb	The rate of leaf damage - measured by visually estimating
rate_petals	The rate of petal damage - measured by visually estimating

## 8.3 Additional Models for the Rate of Leaf Damage (Herbivory)

### 8.3.0.1 Categorical Regression

Table 7: Categorical Regression for Rate of Leaf Damage (Herbivory)

Coefficients	Estimate	P Value
Color,Red	0.3000	0.5883
Height	-0.0055	0.8595
Flowersexual,Male	-0.0087	0.9866
[0,0.3%) (0.3%,2.5%)	-0.2157	0.7814
(0.3%,2.5%) (2.5%, 100%]	1.2235	0.1242

### 8.3.0.2 Linear Regression

Table 8: Linear Regression for Rate of Leaf Damage (Herbivory)

Coefficients	Estimate	P Value
Intercept	0.0212	0.5359
Color,Red	0.0022	0.9314
Height	0.0005	0.6952
Flowersexual,Male	0.0343	0.1452

## 8.4 Additional Models for the Rate of Petals Damage (Florivory)

### 8.4.0.1 Categorical Regression

Table 9: Categorical Regression for Rate of Petals Damage (Florivory)

Coefficients	Estimate	P Value	
Color,Red	-0.2661	0.6587	
Height	0.0152	0.6334	
Flowersexual,Male	1.2528	0.0383	*
0 (0,1%)	-0.4891	0.5582	
(0,1%) [1%, 100%]	2.7118	0.0037	**

### 8.4.0.2 Linear Regression

Table 10: Linear Regression for Rate of Petals Damage (Florivory)

Coefficients	Estimate	P Value
Intercept	0.0301	0.3088
Color,Red	-0.0121	0.5834
Height	0.0002	0.8790
Flowersexual,Male	0.0261	0.1967

## 8.5 Code and Data

## Reference

- Cribari-Neto, Francisco, and Achim Zeileis. 2010. “Beta Regression in *R*.” *Journal of Statistical Software* 34 (2). <https://doi.org/10.18637/jss.v034.i02>.
- Dafni, A., P. Bernhardt, A. Shmida, Y. Ivri, S. Greenbaum, Ch. O’Toole, and L. Losito. 1990. “RED BOWL-SHAPED FLOWERS: CONVERGENCE FOR BEETLE POLLINATION IN THE MEDITERRANEAN REGION.” *Israel Journal of Botany* 39 (1-2): 81–92. <https://doi.org/10.1080/0021213X.1990.10677134>.
- EHRENDORFER, FRIEDRICH, SVETLANA N. ZIMAN, CHRISTIANE KÖNIG, CARL. S. KEENER, BRYAN E. DUTTON, OLGA N. TSARENKO, ELENA V. BULAKH, MONICA BOŞCAIU, FRÉDÉRIC MÉDAIL, and ARNDT KÄSTNER. 2009. “Taxonomic Revision, Phylogenetics and Transcontinental Distribution of *Anemones* section *Anemone* (Ranunculaceae).” *Botanical Journal of the Linnean Society* 160 (3): 312–54. <https://doi.org/10.1111/j.1095-8339.2009.00861.x>.
- Horovitz, Auguste, Susan Bullock, and Moshe Negbi. 1975. “Germination Characters in Wild and Cultivated *Anemone Coronaria* L.” *Euphytica* 24 (1): 213–20. <https://doi.org/10.1007/bf00147188>.
- Kooi, Casper J. van der, J. Theo M. Elzenga, Marten Staal, and Doekele G. Stavenga. 2016. “How to Colour a Flower: On the Optical Principles of Flower Coloration.” *Proceedings of the Royal Society B: Biological Sciences* 283 (1830): 20160429. <https://doi.org/10.1098/rspb.2016.0429>.
- Kooi, Casper J. van der, Doekele G. Stavenga, Kentaro Arikawa, Gregor Belušič, and Almut Kelber. 2021. “Evolution of Insect Color Vision: From Spectral Sensitivity to Visual Ecology.” *Annual Review of Entomology* 66 (1): 435–61. <https://doi.org/10.1146/annurev-ento-061720-071644>.
- Martínez-Harms, María José, and Patricia Balvanera. 2012. “Methods for Mapping Ecosystem Service Supply: A Review.” *International Journal of Biodiversity Science, Ecosystem Services & Management* 8 (1-2): 17–25. <https://doi.org/10.1080/21513732.2012.663792>.
- Sapir, Yuval, M. Kate Gallagher, and Esther Senden. 2021. “What Maintains Flower Colour Variation Within Populations?” *Trends in Ecology & Evolution* 36 (6): 507–19. <https://doi.org/10.1016/j.tree.2021.01.011>.