

Asymptotic Properties of the Hill estimator

Jaakko Pere

School of Science

Bachelor's thesis
Espoo xx.8.2018

Supervisor

Ph.D Pauliina Ilmonen

Advisor

M.Sc Matias Heikkilä

Copyright © 2018 Jaakko Pere

Author Jaakko Pere

Title Asymptotic Properties of the Hill estimator

Degree programme Technical Physics and Mathematics

Major Mathematics and Systems Analysis

Code of major SCI3025

Supervisor Ph.D Pauliina Ilmonen

Advisor M.Sc Matias Heikkilä

Date xx.8.2018

Number of pages 11+1

Language English

Abstract

Your abstract in English. Keep the abstract short. The abstract explains your research topic, the methods you have used, and the results you obtained.

The abstract text of this thesis is written on the readable abstract page as well as into the pdf file's metadata via the \thesisabstract macro (see above). Write here the text that goes onto the readable abstract page. You can have special characters, linebreaks, and paragraphs here. Otherwise, this abstract text must be identical to the metadata abstract text.

If your abstract does not contain special characters and it does not require paragraphs, you may take advantage of the abstracttext macro (see the comment below).

Keywords For keywords choose, concepts that are, central to your, thesis

Tekijä Jaakko Pere

Työn nimi Opinnäytteen otsikko

Koulutusohjelma Elektroniikka ja sähkötekniikka

Pääaine Sopiva pääaine

Pääaineen koodi SCI3025

Vastuuopettaja Ph.D Pauliina Ilmonen

Työn ohjaaja TkT Alan Advisor

Päivämäärä xx.8.2018

Sivumäärä 11+1

Kieli Englanti

Tiivistelmä

Tiivistelmässä on lyhyt selvitys kirjoituksen tärkeimmästä sisällöstä: mitä ja miten on tutkittu, sekä mitä tuloksia on saatu.

Avainsanat Vastus, resistanssi, lämpötila

Preface

I want to thank Professor Pirjo Professori and my instructor Dr Alan Advisor for their good and poor guidance.

Otaniemi, 24.4.2018

Eddie E. A. Engineer

Contents

Abstract	3
Abstract (in Finnish)	4
Preface	5
Contents	6
Symbols and abbreviations	7
1 Introduction	8
2 Background	9
2.1 Extreme Value Theory	9
2.2 Problem	9
3 Consistency of the Hill estimator	10
4 Simulations	10
References	11

Symbols and abbreviations

Symbols

$x^* = \sup\{x : F(x) < 1\}$ right endpoint of the distribution
 γ extreme value index

Operators

Abbreviations

cdf cumulative distribution function
 i.d.d independent and identically distributed

1 Introduction

2 Background

2.1 Extreme Value Theory

First approach to study behavior of extreme events would be to find limiting distribution of the sample maxima $M_n = \max(X_1, X_2, \dots, X_n)$. Here X_1, X_2, \dots, X_n are i.i.d random variables from cdf F . Function for the cdf of M_n can be derived, because X_1, X_2, \dots, X_n are i.i.d.

$$\begin{aligned} P(\max(X_1, X_2, \dots, X_n) \leq x) &= P(X_1 \leq x, X_2 \leq x, \dots, X_n \leq x) = \\ &P(X_1 \leq x)P(X_2 \leq x) \dots P(X_n \leq x) = F^n(x). \end{aligned}$$

Now it can be shown that this approach is not very fruitful since

$$\lim_{n \rightarrow \infty} F^n(x) = \begin{cases} 0, & x < x^* \\ 1, & x \geq x^*. \end{cases}$$

To achieve a nondegenerate distribution it is necessary to normalize the sample maxima M_n . After normalization a nondegenerate distribution is gained as stated in the Fisher-Tippett-Gnedenko Theorem [1].

Theorem 2.1 *There exists real constants $a_n > 0$ and $b_n \in \mathbb{R}$ such that*

$$\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = G_\gamma(x) = \begin{cases} \exp(-(1 + \gamma x)^{-\frac{1}{\gamma}}), & \gamma \neq 0 \\ \exp(-e^{-x}), & \gamma = 0, \end{cases} \quad (1)$$

for all x with $1 + \gamma x > 0$ where $\gamma \in \mathbb{R}$.

If cdf F satisfies the equation 1 for some $\gamma \in \mathbb{R}$ it is said that F is in the maximum domain of attraction of G_γ i.e $F \in D(G_\gamma)$. Concerning the Hill estimator special interest lies in the case $F \in D(G_{\gamma>0})$.

2.2 Problem

3 Consistency of the Hill estimator

4 Simulations

References

- [1] A. F. Laurens De Haan. *Extreme Value Theory*. Springer, 2009.

Appendix