



Aalto University

Moment Estimator-Based Extreme Quantile Estimation with Erroneous Observations

Joint work with Pauliina Ilmonen and Lauri Viitasaari

<https://doi.org/10.48550/arXiv.2502.08510>

Jaakko Pere

Thursday 28th May, 2026

Agenda of the Presentation

Motivation

Preliminaries of Extreme Value Theory

Extreme Value Theory with Approximations

Extreme Quantile Region Estimation under Ellipticity

Table of Contents

Motivation

Preliminaries of Extreme Value Theory

Extreme Value Theory with Approximations

Extreme Quantile Region Estimation under Ellipticity

Extreme Value Theory: Tools for Extrapolation

- Extreme value theory is concerned with inference of rare events.

Extreme Value Theory: Tools for Extrapolation

- Extreme value theory is concerned with inference of rare events.
- Consider estimating 0.999-quantile (red cross) with a reasonable sample (black circles).

Extreme Value Theory: Tools for Extrapolation

- Extreme value theory is concerned with inference of rare events.
- Consider estimating 0.999-quantile (red cross) with a reasonable sample (black circles).




Figure: Insurance company keeps records of fire insurance claims per windstorm. In total, the figure shows 700 observations.

Noisy Observations

What if instead of the original observations only approximations are available?

⇒ Extreme value inference with the noisy sample.

Noisy Observations

What if instead of the original observations only approximations are available?

⇒ Extreme value inference with the noisy sample.

Why?

Noisy Observations

What if instead of the original observations only approximations are available?

⇒ Extreme value inference with the noisy sample.

Why?

- Extreme value index estimation for projections of functional data (Kokoszka and Xiong 2018).
- Extreme quantile/expectile regression (Girard et al. 2021).
- Latent model extreme value index estimation (Virta et al. 2024).

Noisy Observations

What if instead of the original observations only approximations are available?

⇒ Extreme value inference with the noisy sample.

Why?

- Extreme value index estimation for projections of functional data (Kokoszka and Xiong 2018).
- Extreme quantile/expectile regression (Girard et al. 2021).
- Latent model extreme value index estimation (Virta et al. 2024).

Much more work has been done especially in the heavy-tailed case (few or no moments exist)...

Approaches from the Literature (for the Heavy Tail)

1. (Kim and Kokoszka 2020; Kim and Kokoszka 2023): We have

$\hat{Y}_i = Y_i + \varepsilon_i$, where

- the errors ε_i are i.i.d.,
- $\{Y_i\} \perp\!\!\!\perp \{\varepsilon_i\}$, and
- $\lim_{t \rightarrow \infty} \mathbb{P}(|\varepsilon| > t) / \mathbb{P}(X_1 > t) = 0$

Approaches from the Literature (for the Heavy Tail)

1. (Kim and Kokoszka 2020; Kim and Kokoszka 2023): We have

$\hat{Y}_i = Y_i + \varepsilon_i$, where

- the errors ε_i are i.i.d.,
- $\{Y_i\} \perp\!\!\!\perp \{\varepsilon_i\}$, and
- $\lim_{t \rightarrow \infty} \mathbb{P}(|\varepsilon| > t) / \mathbb{P}(X_1 > t) = 0$

2. (Girard et al. 2021): We have that the errors $|\hat{Y}_i - Y_i|$

- can be dependent,
- they do not need to be identically distributed, and
- $\{Y_i\} \perp\!\!\!\perp \{\varepsilon_i\}$ can fail.
- $\max_{i \in \{1, \dots, n\}} |\hat{Y}_i - Y_i| / Z_n \xrightarrow{\mathbb{P}} 0$, as $n \rightarrow \infty$.

Scope of the Presentation

1. We quantify the effect of approximation errors for a certain *extreme quantile estimator*.
 - Short-, light-, and heavy-tailed distributions are considered.

Scope of the Presentation

1. We quantify the effect of approximation errors for a certain *extreme quantile estimator*.
 - Short-, light-, and heavy-tailed distributions are considered.
2. Application to multivariate extreme quantile region estimation under ellipticity.

Table of Contents

Motivation

Preliminaries of Extreme Value Theory

Extreme Value Theory with Approximations

Extreme Quantile Region Estimation under Ellipticity

Maximum Domain of Attraction ($Y \in \text{MDA}(G)$)

Let $Y_1, \dots, Y_n$ be i.i.d. observations of a random variable Y . We have $Y \in \text{MDA}(G)$ if there exist sequences $a_n > 0$ and $b_n \in \mathbb{R}$, and a random variable G with a nondegenerate distribution such that

$$\frac{\max(Y_1, \dots, Y_n) - b_n}{a_n} \xrightarrow{\mathcal{D}} G, \quad n \rightarrow \infty.$$

Extreme Value Index (Fisher and Tippett 1928; Gnedenko 1943)

Up to location and scale, the distribution of $G = G_\gamma$ is characterized by the parameter γ , called the *extreme value index*. That is, the distribution of G_γ is of the type

$$F_{G_\gamma}(x) = \begin{cases} \exp\left(-(1 + \gamma x)^{-1/\gamma}\right), & 1 + \gamma x > 0 \quad \text{if } \gamma \neq 0, \\ \exp(-e^{-x}), & x \in \mathbb{R} \quad \text{if } \gamma = 0. \end{cases}$$

Maximum domain of attraction $\text{MDA}(G_\gamma)$ for different values of the extreme value index γ .

γ	Example of $F \in \text{MDA}(G_\gamma)$	Properties of F
> 0	Pareto distribution	For a larger γ fewer moments exist*
$= 0$	Normal distribution	All moments exist*
< 0	Uniform distribution	Finite endpoint

*The claims about the moments hold if the left tail of a distribution is not considered.

Tail Quantile Function

Define the tail quantile function corresponding to a distribution F by

$$U(t) = F^{\leftarrow} \left(1 - \frac{1}{t} \right), \quad t > 1,$$

where we denote the left-continuous inverse of a nondecreasing function by $f^{\leftarrow}(y) = \inf \{x \in \mathbb{R} : f(x) \geq y\}$.

Tail Quantile Function

Define the tail quantile function corresponding to a distribution F by

$$U(t) = F^{\leftarrow} \left(1 - \frac{1}{t} \right), \quad t > 1,$$

where we denote the left-continuous inverse of a nondecreasing function by $f^{\leftarrow}(y) = \inf \{x \in \mathbb{R} : f(x) \geq y\}$.

That is, $U(1/p)$ is the $(1 - p)$ -quantile.

MDA Condition in Terms of Quantiles (de Haan 1984)

Let U be the tail quantile function corresponding to a random variable Y . Then, $Y \in \text{MDA}(G_\gamma)$ if and only if there exists a positive function $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for all $x > 0$,

$$\lim_{t \rightarrow \infty} \frac{U(tx) - U(t)}{a(t)} = \frac{x^\gamma - 1}{\gamma}.$$

MDA Condition in Terms of Quantiles (de Haan 1984)

Let U be the tail quantile function corresponding to a random variable Y . Then, $Y \in \text{MDA}(G_\gamma)$ if and only if there exists a positive function $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for all $x > 0$,

$$\lim_{t \rightarrow \infty} \frac{U(tx) - U(t)}{a(t)} = \frac{x^\gamma - 1}{\gamma}.$$

Choose $t = n/k$ and $x = k/(np)$ to get the approximation

$$U\left(\frac{1}{p}\right) \approx U\left(\frac{n}{k}\right) + a\left(\frac{n}{k}\right) \frac{\left(\frac{k}{np}\right)^\gamma - 1}{\gamma}.$$

Extreme Quantile Estimation

Suppose $\mathbf{Y} = (Y_1, \dots, Y_n)$ is an i.i.d. sample of $Y \in \text{MDA}(G_\gamma)$, $\gamma \in \mathbb{R}$. Denote order statistics corresponding to the sample $\mathbf{Y}$ by $\mathbf{Y}_{1,n} \leq \dots \leq \mathbf{Y}_{n,n}$. Then an estimator for the extreme $(1 - p)$ -quantile $x_p = U(1/p)$ can be given as

$$\hat{x}_p(\mathbf{Y}) = \mathbf{Y}_{n-k,n} + \hat{a}\left(\frac{n}{k}\right) \frac{\left(\frac{k}{np}\right)^{\hat{\gamma}(\mathbf{Y})} - 1}{\hat{\gamma}(\mathbf{Y})},$$

where $\hat{\gamma}$ is an estimator for the extreme value index γ .

The Moment Estimator (Dekkers et al. 1989)

Set, for $\ell \in \{1, 2\}$,

$$M_n^{(\ell)}(\mathbf{Y}) = \frac{1}{k} \sum_{j=0}^{k-1} (\ln \mathbf{Y}_{n-j,n} - \ln \mathbf{Y}_{n-k,n})^\ell,$$

and

$$\hat{\gamma}_+(\mathbf{Y}) = M_n^{(1)}(\mathbf{Y}) \quad \text{and} \quad \hat{\gamma}_-(\mathbf{Y}) = 1 - \frac{1}{2} \left(1 - \frac{(M_n^{(1)}(\mathbf{Y}))^2}{M_n^{(2)}(\mathbf{Y})} \right)^{-1}.$$

Then the moment estimator is defined as

$$\hat{\gamma}_M(\mathbf{Y}) = \hat{\gamma}_+(\mathbf{Y}) + \hat{\gamma}_-(\mathbf{Y}),$$

and the corresponding estimator for $a(n/k)$ is defined as

$$\hat{\sigma}_M(\mathbf{Y}) = \mathbf{Y}_{n-k,n} M_n^{(1)}(\mathbf{Y}) (1 - \hat{\gamma}_-(\mathbf{Y})).$$

Table of Contents

Motivation

Preliminaries of Extreme Value Theory

Extreme Value Theory with Approximations

Extreme Quantile Region Estimation under Ellipticity

Back to Approximations

- What if instead of the sample $\mathbf{Y}$, only approximations $\hat{\mathbf{Y}} = (\hat{Y}_1, \dots, \hat{Y}_n)$ are available?

Back to Approximations

- What if instead of the sample $\mathbf{Y}$, only approximations $\hat{\mathbf{Y}} = (\hat{Y}_1, \dots, \hat{Y}_n)$ are available?
- How the approximation error affects the asymptotics?

Back to Approximations

- What if instead of the sample $\mathbf{Y}$, only approximations $\hat{\mathbf{Y}} = (\hat{Y}_1, \dots, \hat{Y}_n)$ are available?
- How the approximation error affects the asymptotics?

That is, next we quantify how the error

$$\max_{0 \leq j \leq k} \left| \frac{\hat{Y}_{n-j,n}}{Y_{n-j,n}} - 1 \right|$$

has to behave in order to have asymptotic normality for $\hat{\chi}_\rho(\hat{\mathbf{Y}})$.

Assumptions for the Main Result

1. Y is an almost surely positive random variable with $F \in \mathcal{D}(G_\gamma)$, $\gamma \in \mathbb{R}$ (and U satisfies certain second-order conditions).
2. $\mathbf{Y} = (Y_1, \dots, Y_n)$ are i.i.d. copies of Y and let $\hat{\mathbf{Y}} = (\hat{Y}_1, \dots, \hat{Y}_n)$ be an approximated sample.

Assumptions for the Main Result

1. Y is an almost surely positive random variable with $F \in \mathcal{D}(G_\gamma)$, $\gamma \in \mathbb{R}$ (and U satisfies certain second-order conditions).
2. $\mathbf{Y} = (Y_1, \dots, Y_n)$ are i.i.d. copies of Y and let $\hat{\mathbf{Y}} = (\hat{Y}_1, \dots, \hat{Y}_n)$ be an approximated sample.
3. Suppose $k = k_n \rightarrow \infty$, $k_n/n \rightarrow 0$, as $n \rightarrow \infty$.
4. $p = p_n$, $np_n = o(k_n)$ and $\ln(np_n) = o(\sqrt{k_n})$.

Sketch of the Main Result

Suppose that assumptions of the previous slide hold and that

$$\max_{0 \leq j \leq k_n} \left| \frac{\hat{\mathbf{Y}}_{n-j,n}}{\mathbf{Y}_{n-j,n}} - 1 \right| = O_{\mathbb{P}}(h_n)$$

for some sequence h_n such that $\sqrt{k_n} \frac{h_n U\left(\frac{n}{k_n}\right)}{a\left(\frac{n}{k_n}\right)} = o(1)$. Denote

$q_{\gamma}(t) = \int_1^t s^{\gamma-1} \ln s \, ds$. Then

$$\sqrt{k_n} \frac{\hat{x}_{p_n}(\hat{\mathbf{Y}}) - U(1/p_n)}{a\left(\frac{n}{k_n}\right) q_{\gamma}\left(\frac{k_n}{np_n}\right)}$$

is asymptotically normal.

Condition for the Error

Suppose $k = k_n \rightarrow \infty$, $k_n/n \rightarrow 0$, as $n \rightarrow \infty$. Assume also that U satisfies a certain second-order condition and that

$$\max_{0 \leq j \leq k_n} \left| \frac{\hat{\mathbf{Y}}_{n-j,n}}{\mathbf{Y}_{n-j,n}} - 1 \right| = O_{\mathbb{P}}(h_n)$$

for some sequence h_n . Then $\sqrt{k_n} \frac{h_n U\left(\frac{n}{k_n}\right)}{a\left(\frac{n}{k_n}\right)} \rightarrow 0$ if

1. $\gamma > 0$ and $\sqrt{k_n} h_n \rightarrow 0$,
2. $\gamma = 0$ and $\sqrt{k_n} h_n = O\left(\left(\frac{n}{k_n}\right)^{-\delta}\right)$ for some $\delta > 0$,
3. $\gamma < 0$ and $\sqrt{k_n} h_n = o\left(\left(\frac{n}{k_n}\right)^{\gamma}\right)$.

Example

If

$$\max_{0 \leq j \leq k_n} \left| \frac{\hat{\mathbf{y}}_{n-j,n}}{\mathbf{y}_{n-j,n}} - 1 \right| = o_{\mathbb{P}} \left(\frac{1}{\sqrt{n}} \right),$$

then the sufficient condition for the error does not hold if $\gamma \leq -1/2$!

Table of Contents


Motivation

Preliminaries of Extreme Value Theory

Extreme Value Theory with Approximations

Extreme Quantile Region Estimation under Ellipticity

t-distribution, $n = 1000$, $k = 100$, $\gamma = 0.25$, $p = 1/5000$, $\alpha_{\text{MCD}} = 0.5$



Extreme Quantile Region Estimation

Let $X \in \mathbb{R}^d$ be a random vector with density f . We wish to estimate

$$Q_p = \{x \in \mathbb{R}^d : f(x) \leq \beta_p\},$$

where β_p is chosen such that $\mathbb{P}(X \in Q_p) = p$, and p is small.

Extreme Quantile Region Estimation

Let $X \in \mathbb{R}^d$ be a random vector with density f . We wish to estimate

$$Q_p = \{x \in \mathbb{R}^d : f(x) \leq \beta_p\},$$

where β_p is chosen such that $\mathbb{P}(X \in Q_p) = p$, and p is small.

The problem has been considered under multivariate regular variation by

- Cai et al. 2011 in the context of density contours, and
- He and Einmahl 2017 in the context of halfspace depth contours under multivariate regular variation.

Elliptical Distributions (Cambanis et al. 1981)

We say that a d -variate random vector X is elliptically distributed if

$$X \stackrel{\mathcal{D}}{=} \mu + \mathcal{R}\Lambda S,$$

where

- $\mu \in \mathbb{R}^d$ is called the *location vector*,
- $\mathcal{R}$ is a nonnegative random variable called the *generating variate*,
- $\Lambda \in \mathbb{R}^{d \times d}$ is a matrix such that $\Sigma = \Lambda\Lambda^\top$ is a symmetric positive definite matrix (matrix Σ is called *scatter matrix*),
- S is an d -dimensional random vector uniformly distributed over the unit-sphere $\{x \in \mathbb{R}^d : x^\top x = 1\}$ and
- Random variables $\mathcal{R}$ and S are independent.

Role of the Generating Variate $\mathcal{R}$

- We suppose $\mathcal{R} \in \text{MDA}(G_\gamma)$ for $\gamma \in \mathbb{R}$

Role of the Generating Variate $\mathcal{R}$

- We suppose $\mathcal{R} \in \text{MDA}(G_\gamma)$ for $\gamma \in \mathbb{R}$
- $\mathcal{R} \stackrel{\mathcal{D}}{=} \sqrt{(X - \mu)^\top \Sigma^{-1} (X - \mu)}$

Role of the Generating Variate $\mathcal{R}$

- We suppose $\mathcal{R} \in \text{MDA}(G_\gamma)$ for $\gamma \in \mathbb{R}$
- $\mathcal{R} \stackrel{\mathcal{D}}{=} \sqrt{(X - \mu)^\top \Sigma^{-1} (X - \mu)}$
- $\hat{R}_i = \sqrt{(X_i - \hat{\mu}(\mathbf{X}))^\top \hat{\Sigma}^{-1}(\mathbf{X}) (X_i - \hat{\mu}(\mathbf{X}))}$
- $\hat{\mathbf{R}} = (\hat{R}_1, \dots, \hat{R}_n)$

The Estimator

- We wish to estimate

$$Q_p = \left\{ x \in \mathbb{R}^d : \sqrt{(x - \mu)^\top \Sigma^{-1} (x - \mu)} \geq r_p \right\},$$

for a very small p , where r_p is the $(1 - p)$ -quantile of the generating variate $\mathcal{R}$.

The Estimator

- We wish to estimate

$$Q_p = \left\{ x \in \mathbb{R}^d : \sqrt{(x - \mu)^\top \Sigma^{-1} (x - \mu)} \geq r_p \right\},$$

for a very small p , where r_p is the $(1 - p)$ -quantile of the generating variate $\mathcal{R}$.

- Define estimator $\hat{Q}_p$ as

$$\hat{Q}_p = \left\{ x \in \mathbb{R}^d : \sqrt{(x - \hat{\mu}(\mathbf{X}))^\top \hat{\Sigma}^{-1}(\mathbf{X}) (x - \hat{\mu}(\mathbf{X}))} \geq \hat{x}_p(\hat{\mathbf{R}}) \right\},$$

where $\hat{x}_p(\hat{\mathbf{R}})$ is the extreme quantile estimator computed with approximations $\hat{\mathbf{R}}$.

Approximation Error for the Generating Variate

Assume $k = k_n \rightarrow \infty$, $k_n/n \rightarrow 0$, as $n \rightarrow \infty$ and that estimators of the location and scatter satisfy $\sqrt{n}(\hat{\mu}_n - \mu) = O_{\mathbb{P}}(1)$ and $\sqrt{n}(\hat{\Sigma}_n - \Sigma) = O_{\mathbb{P}}(1)$. Then

$$\max_{0 \leq j \leq k_n} \left| \frac{\hat{\mathbf{R}}_{n-j,n}}{\mathbf{R}_{n-j,n}} - 1 \right| = O_{\mathbb{P}}\left(\frac{1}{\sqrt{n}}\right).$$

Consistency and Affine Equivariance

- **Consistency:** if estimators of the location and scatter are $\sqrt{n}$ -consistent, $\mathcal{R} \in \text{MDA}(\mathbf{G}_\gamma)$ for $\gamma > -1/2$, $p_n \rightarrow 0$ with a suitable rate, as $n \rightarrow \infty$ (and other technical conditions),

$$\frac{\mathbb{P}\left(X \in \hat{Q}_{p_n} \triangle Q_{p_n}\right)}{p_n} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

where $B_1 \triangle B_2 = (B_1 \setminus B_2) \cup (B_2 \setminus B_1)$, $B_1, B_2 \in \mathbb{R}^d$.


Consistency and Affine Equivariance

- **Consistency:** if estimators of the location and scatter are $\sqrt{n}$ -consistent, $\mathcal{R} \in \text{MDA}(\mathbf{G}_\gamma)$ for $\gamma > -1/2$, $p_n \rightarrow 0$ with a suitable rate, as $n \rightarrow \infty$ (and other technical conditions),

$$\frac{\mathbb{P}\left(X \in \hat{Q}_{p_n} \triangle Q_{p_n}\right)}{p_n} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

where $B_1 \triangle B_2 = (B_1 \setminus B_2) \cup (B_2 \setminus B_1)$, $B_1, B_2 \in \mathbb{R}^d$.

- **Equivariance under affine transformations** $Ax + b$, $A \in \mathbb{R}^{d \times d}$ invertible.



— Theoretical

--- Estimate

Link to slides: [https://github.com/perej1/presentations/
tree/main/light-error](https://github.com/perej1/presentations/tree/main/light-error)

References

- Cai, Juan-Juan, John H. J. Einmahl, and Laurens de Haan (2011). “Estimation of extreme risk regions under multivariate regular variation”. In: *The Annals of Statistics* 39.3, pp. 1803–1826. DOI: 10.1214/11-AOS891.
- Cambanis, S., S. Huang, and G. Simons (1981). “On the theory of elliptically contoured distributions”. In: *Journal of Multivariate Analysis* 11.3, pp. 368–385.
- de Haan, Laurens (1984). “Slow Variation and Characterization of Domains of Attraction”. In: *Statistical Extremes and Applications*. Ed. by J. Tiago de Oliveira. Dordrecht: Springer, pp. 31–48. DOI: 10.1007/978-94-017-3069-3_4.

References II

- Dekkers, A. L. M., J. H. J. Einmahl, and L. de Haan (1989). “A moment estimator for the index of an extreme-value distribution”. In: *The Annals of Statistics* 17.4, pp. 1833–1855. URL: <https://www.jstor.org/stable/2241667>.
- Fisher, Ronald Aylmer and Leonard Henry Caleb Tippett (1928). “Limiting forms of the frequency distribution of the largest or smallest member of a sample”. In: *Mathematical Proceedings of the Cambridge Philosophical Society* 24.2, pp. 180–190. DOI: 10.1017/S0305004100015681.
- Girard, Stéphane, Gilles Stupfler, and Antoine Usseglio-Carleve (2021). “Extreme conditional expectile estimation in heavy-tailed heteroscedastic regression models”. In: *The Annals of Statistics* 49.6, pp. 3358–3382. DOI: 10.1214/21-AOS2087.

References III

- Gnedenko, Boris (1943). “Sur La Distribution Limite Du Terme Maximum D’Une Série Aléatoire”. In: *Annals of Mathematics* 44.3, pp. 423–453. DOI: 10.2307/1968974.
- He, Yi and John H. J. Einmahl (2017). “Estimation of extreme depth-based quantile regions”. In: *Journal of the Royal Statistical Society. Series B (Statistical Methodology)* 79.2, pp. 449–461. URL: <http://www.jstor.org/stable/44682520>.
- Kim, Mihyun and Piotr Kokoszka (2020). “Consistency of the Hill estimator for time series observed with measurement errors”. In: *Journal of Time Series Analysis* 41.3, pp. 421–435. DOI: 10.1111/jtsa.12515.
- (2023). “Asymptotic and finite sample properties of Hill-type estimators in the presence of errors in observations”. In: *Journal of Nonparametric Statistics* 35.1, pp. 1–18. DOI: 10.1080/10485252.2022.2136662.

References IV

- Kokoszka, Piotr and Qian Xiong (2018). “Extremes of projections of functional time series on data-driven basis systems”. In: *Extremes* 21, pp. 177–204. DOI: 10.1007/s10687-017-0302-8.
- Virta, Joni et al. (2024). “Latent model extreme value index estimation”. In: *Journal of Multivariate Analysis* 202, p. 105300. DOI: 10.1016/j.jmva.2024.105300.