

# PGRE: Studying for Death?

Gray Ezequiel G.B. Perez<sup>1</sup> and Schlack-Dog<sup>1</sup>

<sup>1</sup>*Reed College, 3203 SE Woodstock Blvd, Portland, OR 97202, USA*

Attempting to avoid death by making a big study guide!

## CONTENTS

I. Variable definitions	1
II. Classical mechanics (20%)	1
III. Electromagnetism (18%)	1
A. Maxwell's Equations	1
B. Magnetic and Electric Fields in Matter	1
IV. Quantum Mechanics (12%)	2
V. Thermodynamics and Statistical Mechanics (10%)	2
VI. Atomic Physics (10%)	2
VII. Optics and Wave Phenomena (9%)	2
A. Doppler Effect	2
VIII. Specialized Topics (9%)	2
IX. Special Relativity (6%)	2
X. Laboratory Methods (6%)	2
A. Error Analysis	2
B. Instrumentation	3

## I. VARIABLE DEFINITIONS

$a \rightarrow$  acceleration

$c \rightarrow$  speed of light:

standard  $\approx 3 * 10^8 \frac{m}{s}$

## II. CLASSICAL MECHANICS (20%)

Such as kinematics, Newton's laws, work and energy, oscillatory motion, rotational motion about a fixed axis, dynamics of systems of particles, central forces and celestial mechanics, three-dimensional particle dynamics, Lagrangian and Hamiltonian formalism, noninertial reference frames, elementary topics in fluid dynamics.

## III. ELECTROMAGNETISM (18%)

Such as electrostatics, currents and DC circuits, magnetic fields in free space, Lorentz force, induction,



FIG. 1. Do figures work? Hell if I know. Here's a salamander.

Maxwell's equations and their applications, electromagnetic waves, AC circuits, magnetic and electric fields in matter.

### A. Maxwell's Equations

- **Speed of light:** The definition (in vacuum) can be found by taking the curl of Maxwell's equations and looking for something in the form of the wave equation. It is

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}. \quad (1)$$

For light in matter, replace  $\mu_0$  and  $\epsilon_0$  with  $\mu$  and  $\epsilon$ , respectively.

### B. Magnetic and Electric Fields in Matter

- **Permittivity:** In matter, the permittivity is given by

$$\epsilon = \kappa_E \epsilon_0, \quad (2)$$

where  $\epsilon$  is the absolute permittivity,  $\kappa_E$  is the dielectric constant, and  $\epsilon_0$  is vacuum permittivity.

If the applied field is not constant, then  $\epsilon$  becomes frequency-dependent because the material's polarization does not change instantly. In this case

$$\epsilon(\omega) = \kappa_E \epsilon_0 - i \frac{\sigma}{\omega}, \quad (3)$$

where  $\sigma$  is the conductivity of the material and  $\omega$  is the frequency of the applied field.

Note that electric susceptibility is related to the dielectric constant with the simple relation

$$\chi_E = \kappa_E - 1. \quad (4)$$

- **Permeability:** The permeability is given by

$$\mu = \kappa_B \mu_0, \quad (5)$$

where  $\mu$  is the absolute permeability,  $\kappa_B$  is a constant, and  $\mu_0$  is vacuum permeability.  $\kappa_B$  is related to the magnetic susceptibility by

$$\chi_B = \kappa_B - 1. \quad (6)$$

Similarly to electric fields in matter,  $\mu$  does have a frequency dependence. However, this dependence is negligible in non-magnetic materials.

Fun fact: a material with  $\mu < \mu_0$  is **diamagnetic**, and a material with  $\mu > \mu_0$  is **paramagnetic**.

#### IV. QUANTUM MECHANICS (12%)

Such as fundamental concepts, solutions of the Schrödinger equation (including square wells, harmonic oscillators, and hydrogenic atoms), spin, angular momentum, wave function symmetry, elementary perturbation theory.

#### V. THERMODYNAMICS AND STATISTICAL MECHANICS (10%)

Such as the laws of thermodynamics, thermodynamic processes, equations of state, ideal gases, kinetic theory, ensembles, statistical concepts and calculation of thermodynamic quantities, thermal expansion and heat transfer.

#### VI. ATOMIC PHYSICS (10%)

Such as properties of electrons, Bohr model, energy quantization, atomic structure, atomic spectra, selection rules, black-body radiation, x-rays, atoms in electric and magnetic fields.

#### VII. OPTICS AND WAVE PHENOMENA (9%)

Such as wave properties, superposition, interference, diffraction, geometrical optics, polarization, Doppler effect.

#### A. Doppler Effect

The general form of the (classical) Doppler effect is

$$f = \frac{c \pm v_r}{c \pm v_s} f_0, \quad (7)$$

where  $f$  is the observed frequency,  $f_0$  is the source frequency,  $c$  is the velocity of the waves in medium,  $v_r$  is the velocity of the receiver relative to the medium, and  $v_s$  is the velocity of the source relative to the medium.  $v_r$  is positive if the receiver is moving towards the source and negative if it is moving away from the source.  $v_s$  is positive if the source is moving away from the receiver and negative if it is towards the receiver.

In the relativistic case, the velocities above should be added using the velocity addition formula. This gives

$$f = \sqrt{\frac{1 - v/c}{1 + v/c}} f_0, \quad (8)$$

where  $v$  is the velocity of the source relative to the observer.  $v$  is positive if the source and observer are receding from each other, and negative if the source and observer are approaching each other.

#### VIII. SPECIALIZED TOPICS (9%)

Nuclear and Particle physics (e.g., nuclear properties, radioactive decay, fission and fusion, reactions, fundamental properties of elementary particles), Condensed Matter (e.g., crystal structure, x-ray diffraction, thermal properties, electron theory of metals, semiconductors, superconductors), Miscellaneous (e.g., astrophysics, mathematical methods, computer applications)

#### IX. SPECIAL RELATIVITY (6%)

Such as introductory concepts, time dilation, length contraction, simultaneity, energy and momentum, four-vectors and Lorentz transformation, velocity addition.

#### X. LABORATORY METHODS (6%)

Such as data and error analysis, electronics, instrumentation, radiation detection, counting statistics, interaction of charged particles with matter, lasers and optical interferometers, dimensional analysis, fundamental applications of probability and statistics.

#### A. Error Analysis

- **Counting error:** the error in counting a sample of size  $N$  is given by

$$\Delta N = \sqrt{N}. \quad (9)$$

## B. Instrumentation

### • Lasers:

- *Dye Laser*: Dye lasers use organic dye as a gain medium. Since the gain medium is a liquid, these lasers can be constructed in many different ways. They are very versatile, as they have high wavelength agility and the ability for high-energy pulses and high output power. This gives them many applications in science—they are used in spectroscopy and astronomy. According to Lucas, they can even be edible.
- *Helium-Neon Laser*: The helium-neon (HeNe) laser uses a mixture of helium and neon gas as its gain medium. It is most commonly used for its 632.8 nm (red) output, although it can be tuned to other wavelengths. They are primarily used for research in optics labs due to their relatively low cost, ease of operation, and laser quality.
- *Excimer Laser*: Excimer lasers (sometimes called exciplex lasers) are ultraviolet lasers. They are used in medical applications like eye surgery, as well as microelectronic devices and integrated circuits.
- *Ruby Laser*: As the name suggests, this is a solid-state laser that uses a synthetic ruby crystal as its gain medium. They produce deep red pulses (694.3 nm) that are on the order of milliseconds. They are used to pump dye lasers and in research when short, red pulses are needed. They have other applications such as military rangefinding and hair removal, but in recent years they have been being replaced with more efficient Nd:YAG lasers.
- *Nd:YAG Laser*: Neodymium-doped Yttrium Aluminum Garnet (Nd:YAG) lasers are solid-state lasers that use Nd:YAG as a gain medium. They typically emit infrared light at 1064 nm. They are one of the most commonly used lasers today, with applications in medicine, science, and industry. They are a popular choice for optical tweezers due to their infrared light. Since they are capable of producing fast, high-energy pulses, they are also used for laser-induced breakdown spectroscopy.