

Path-Coloring Algorithms for Planar Graphs

Aven Bross

Department of Computer Science

University of Alaska

Fairbanks, AK 99775-6670

dabross@alaska.edu

Glenn G. Chappell

Department of Computer Science

University of Alaska

Fairbanks, AK 99775-6670

chappellg@member.ams.org

Chris Hartman

Department of Computer Science

University of Alaska

Fairbanks, AK 99775-6670

cmhartman@alaska.edu

May 29, 2017

2010 Mathematics Subject Classification. Primary 05C38; Secondary 05C10, 05C15.

Key words and phrases. Path coloring, list coloring, algorithm.

Abstract

A path coloring of a graph G is a vertex coloring of G so that each color class induces a disjoint union of paths. We present two efficient algorithms to construct a path coloring of a planar graph.

The first algorithm, based on a proof of Poh [6], is given a planar graph; it produces a path coloring of the given graph using three colors.

The second algorithm, based on similar proofs by Hartman [5] and Škrekovski [7], performs a list-coloring generalization of the above. The algorithm is given a planar graph and an assignment of lists of three colors to each vertex; it produces a path coloring of the given graph in which each vertex receives a color from its list.

Implementations of both algorithms are available.

1 Introduction

All graphs will be finite, simple, and undirected.

A *path coloring* of a graph G is a vertex coloring (not necessarily proper) of G so that each color class induces a disjoint union of paths. A graph G is *path k -colorable* if G admits a path coloring using k colors.

Broere & Mynhardt conjectured [1, Conj. 16] that every planar graph is path 3-colorable. This was proven independently by Poh [6, Thm. 2] and by Goddard [4, Thm. 1].

Theorem 1.1 (Poh 1990, Goddard 1991). *Let G be a planar graph. Then G is path 3-colorable.* \square

Hartman [5, Thm. 4.1] (see also Chappell & Hartman [3, Thm. 2.1]) proved a list-coloring generalization of Theorem 1.1. A graph G is *path k -choosable* if, whenever each vertex of G is assigned a list of k colors, there exists a path coloring of G in which each vertex receives a color from its list.

Theorem 1.2 (Hartman 1997). *Let G be a planar graph. Then G is path 3-choosable.* \square

Essentially the same technique was used by Škrekovski [7, Thm. 2.2b] to prove a result slightly weaker than Theorem 1.2.

We discuss two efficient path-coloring algorithms based on proofs of the above theorems.

Section 2 covers an algorithm based on Poh’s proof of Theorem 1.1. The algorithm is given a planar graph; it produces a path coloring of the given graph using three colors.

Section 3 covers an algorithm based Hartman’s proof of Theorem 1.2, along with the proof of Škrekovski mentioned above. The algorithm is given a planar graph and an assignment of a list of three colors to each vertex; it produces a path coloring of the given graph in which each vertex receives a color from its list.

Implementations of both algorithms are available; see Bross [2].

2 Path Coloring: the Poh Algorithm

ZZZ

3 Path List Coloring: the Hartman-Škrekovski Algorithm

ZZZ

References

- [1] I. Broere and C. M. Mynhardt, Generalized colorings of outerplanar and planar graphs, *Graph theory with applications to algorithms and computer science* (Kalamazoo, Mich., 1984), 151–161, Wiley-Intersci. Publ., Wiley, New York, 1985.
- [2] A. Bross, *Path coloring algorithms on plane graphs*, Masters Project, University of Alaska, 2017, available from http://github.com/permutationlock/path_coloring_bgl.

- [3] G. G. Chappell and C. Hartman, Path choosability of planar graphs, in preparation.
- [4] W. Goddard, Acyclic colorings of planar graphs, *Discrete Math.* **91** (1991), no. 1, 91–94.
- [5] C. M. Hartman, *Extremal Problems in Graph Theory*, Ph.D. Thesis, University of Illinois, 1997.
- [6] K. S. Poh, On the linear vertex-arboricity of a planar graph, *J. Graph Theory* **14** (1990), no. 1, 73–75.
- [7] R. Škrekovski, List improper colourings of planar graphs, *Combin. Probab. Comput.* **8** (1999), no. 3, 293–299.