Path-Coloring Algorithms for Planar Graphs

Aven Bross

Department of Computer Science
University of Alaska
Fairbanks, AK 99775-6670
dabross@alaska.edu

Glenn G. Chappell

Department of Computer Science
University of Alaska
Fairbanks, AK 99775-6670
chappellg@member.ams.org

Chris Hartman

Department of Computer Science
University of Alaska
Fairbanks, AK 99775-6670
cmhartman@alaska.edu

May 29, 2017

2010 Mathematics Subject Classification. Primary 05C38; Secondary 05C10, 05C15. Key words and phrases. Path coloring, list coloring, algorithm.

Abstract

A path coloring of a graph G is a vertex coloring of G so that each color class induces a disjoint union of paths. We discuss implementations of two algorithms that construct path colorings of planar graphs.

The first algorithm, based on a proof by Poh [6], is given a planar graph; it produces a path coloring of the given graph using three colors.

The second algorithm performs a list-coloring generalization of the above. It is based on two similar proofs: one by Hartman [5], the other by Škrekovski [7]. The algorithm is given a planar graph and an assignment of lists of three colors to each vertex; it produces a path coloring of the given graph in which each vertex receives a color from its list.

Links to implementations of the two algorithms are provided.

1 Introduction

All graphs will be finite, simple, and undirected.

A path coloring of a graph G is a vertex coloring (not necessarily proper) of G so that each color class induces a linear forest, that is, a forest of maximum degree at most 2, or,

equivalently, a disjoint union of paths. A graph G is path k-colorable if G admits a path coloring using k colors.

Broere & Mynhardt conjectured [1, Conj. 16] that every planar graph is path 3-colorable. This was proven independently by Poh [6, Thm. 2] and by Goddard [4, Thm. 1].

Theorem 1.1 (Poh 1990, Goddard 1991). Let G be a planar graph. Then G is path 3-colorable. \square

Hartman [5, Thm. 4.1] (see also Chappell & Hartman [3, Thm. 2.1]) proved a list-coloring generalization of Theorem 1.1. A graph G is $path \ k$ -choosable if, whenever each vertex of G is assigned a list of k colors, there exists a path coloring of G in which each vertex receives a color from its list.

Theorem 1.2 (Hartman 1997). Let G be a planar graph. Then G is path 3-choosable. \square

A slightly weaker result was proven by Škrekovski [7, Thm. 2.2b] using essentially the same technique.

We discuss two efficient path-coloring algorithms based on proofs of the above results. Section 2 covers an algorithm based on Poh's proof of Theorem 1.1. The algorithm is given a planar graph; it produces a path coloring of the given graph using three colors.

Section 3 covers an algorithm based Hartman's proof of Theorem 1.2, along with a similar proof of a weaker result by Škrekovski. The algorithm is given a planar graph and an assignment of a list of three colors to each vertex; it produces a path coloring of the given graph in which each vertex receives a color from its list.

Implementations of both algorithms are available; see Bross [2].

2 Path Coloring: the Poh Algorithm

ZZZ

3 Path List Coloring: the Hartman-Škrekovski Algorithm

ZZZ

References

[1] I. Broere and C. M. Mynhardt, Generalized colorings of outerplanar and planar graphs, *Graph theory with applications to algorithms and computer science* (Kalamazoo, Mich., 1984), 151–161, Wiley-Intersci. Publ., Wiley, New York, 1985.

- [2] A. Bross, *Path coloring algorithms on plane graphs*, Masters Project, University of Alaska, 2017, available from http://github.com/permutationlock/path_coloring_bgl.
- [3] G. G. Chappell and C. Hartman, Path choosabiility of planar graphs, in preparation.
- [4] W. Goddard, Acyclic colorings of planar graphs, *Discrete Math.* **91** (1991), no. 1, 91–94.
- [5] C. M. Hartman, Extremal Problems in Graph Theory, Ph.D. Thesis, University of Illinois, 1997.
- [6] K. S. Poh, On the linear vertex-arboricity of a planar graph, *J. Graph Theory* **14** (1990), no. 1, 73–75.
- [7] R. Škrekovski, List improper colourings of planar graphs, *Combin. Probab. Comput.* 8 (1999), no. 3, 293–299.