ChatGPT

SMAD4 (DPC4/MADH4) Mutations in Germline and Cancer: A Comprehensive Review

Abstract

Background: **SMAD4** (also known as **DPC4** or **MADH4**) encodes a common mediator of transforming growth factor-β (TGF-β) signaling and has dual importance in hereditary syndromes and somatic tumorigenesis. Germline pathogenic variants in SMAD4 cause **juvenile polyposis syndrome (JPS)** often co-occurring with hereditary hemorrhagic telangiectasia (HHT), whereas somatic inactivation of SMAD4 is frequent in several cancers (notably pancreatic, colorectal, and gastric) [1] [2]. This review synthesizes the evidence on SMAD4 across these contexts, including mechanistic roles, epidemiology, inheritance patterns, clinical phenotypes, diagnostic approaches, management strategies, and outcomes.

Methods: A thorough literature search (1996–2025) was conducted, incorporating original studies, meta-analyses, clinical guidelines, and database analyses (ClinVar, ClinGen, cBioPortal, OncoKB, COSMIC, etc.). Key outcomes extracted include overall and disease-free survival (OS/DFS) hazard ratios, treatment response, metastatic patterns, and quality-of-life metrics. Evidence strength was graded based on study size and design (with preference to cohorts n≥50, meta-analyses, and guidelines). Conflicting findings were explicitly noted and unresolved issues highlighted. (**Figure 1** illustrates literature selection from a representative meta-analysis [3].)

Results: **Germline SMAD4** variants cause autosomal-dominant JPS, characterized by multiple gastrointestinal hamartomatous polyps and ~15% lifetime cancer risk (primarily colorectal and gastric) [4] [5]. SMAD4 carriers often manifest an overlapping HHT phenotype (telangiectasia and arteriovenous malformations), absent in BMPR1A-linked JPS [6] [7]. Clinical practice guidelines recommend early genetic testing and intensive endoscopic surveillance from adolescence [8] [9], as well as screening for AVMs (e.g. echocardiography for pulmonary shunts) [10]. **Somatic SMAD4** alterations (frequently loss-of-function truncating mutations, deletions, or loss of protein expression) occur in ~50% of pancreatic ductal adenocarcinomas (PDAC) [1], ~10–15% of colorectal cancers (CRC) [2] [11], ~5–10% of gastric cancers, ~4–6% of non-small cell lung cancers (NSCLC) [12], and a substantial subset of cholangiocarcinomas [13]. In PDAC and CRC, SMAD4 loss correlates with more aggressive disease and worse survival on univariate analysis [3] [14]. For example, a meta-analysis in PDAC found SMAD4 inactivation conferred a nearly 1.9-fold higher hazard of death after multivariate adjustment [3]. SMAD4 status also stratifies patterns of failure: PDAC tumors retaining SMAD4 tend toward local recurrence, whereas SMAD4-null tumors more often disseminate metastatically [15] [16]. In CRC, SMAD4-mutant cases have ~1.5-fold increased hazard of death and shorter progression-free survival on chemotherapy [11] [14]; SMAD4 mutations co-occur with KRAS/NRAS and predict inferior response to EGFR-targeted therapy in RAS-wildtype metastatic CRC [17] [18]. No targeted therapy exists for SMAD4 loss, though TGF-β pathway inhibitors (e.g. **galunisertib**) have shown modest survival benefit in trials [19].

Conclusions: SMAD4 plays a context-dependent tumor suppressor role: germline SMAD4 mutations drive hereditary polyposis and vascular anomalies, while somatic loss promotes tumor progression and

metastasis in several cancers. SMAD4 status has emerged as a prognostic biomarker and potential predictive marker of therapy response (e.g. in CRC). Management of germline SMAD4 mutation carriers involves multidisciplinary surveillance (endoscopy and HHT screenings) and prophylactic interventions (polypectomy or resection) to mitigate cancer risk. Future research is needed to clarify genotype–phenotype modifiers in JPS, to exploit synthetic lethal vulnerabilities of SMAD4-deficient tumors, and to test TGF-β/SMAD4-targeted strategies. **This review provides an up-to-date synthesis, highlighting consensus findings, dissenting reports, and gaps for further investigation.**

Introduction to SMAD4

SMAD4 (Mothers Against Decapentaplegic homolog 4) is a critical intracellular mediator in the canonical **TGF-β** signaling pathway. Originally named **DPC4** ("Deleted in Pancreatic Carcinoma, locus 4") upon its discovery as a pancreatic tumor suppressor in 1996 [20], SMAD4 is also known as **MADH4**. The SMAD4 protein contains an N-terminal MH1 domain (which binds DNA) and a C-terminal MH2 domain (which mediates transcriptional activation and protein interactions) [21]. A flexible linker region connects MH1 and MH2, containing a **SMAD activation domain (SAD)** important for transcriptional activity [22]. SMAD4 shuttles between cytoplasm and nucleus, aided by nuclear localization (NLS) and export signals (NES) (see **Figure 2A**). Unlike receptor-regulated SMADs (R-SMADs such as SMAD2/3 for TGF-β, and SMAD1/5/8 for bone morphogenetic protein signaling) which must be phosphorylated by TGF-β/BMP receptors, SMAD4 serves as the sole **common-mediator SMAD (Co-SMAD)** that partners with phosphorylated R-SMADs [23] [24]. Upon TGF-β ligand binding and receptor activation, SMAD2/3 are phosphorylated and release from the receptor, then form heteromeric complexes with SMAD4 and translocate to the nucleus [25]. In the nucleus, the SMAD2/3/4 complex binds SMAD-binding elements (SBE) in DNA and recruits co-factors to regulate gene transcription [25]. Through this pathway, SMAD4 mediates TGF-β's canonical growth-inhibitory signals, inducing **cell cycle arrest** (upregulating CDK inhibitors p15^INK4B^, p21^CIP1^, p27^KIP1^) and promoting apoptosis via target genes like **TIEG**, **DAPK**, **GADD45β**, and **BIM** [26] [27]. In normal cells and early tumors, intact TGF-β/SMAD4 signaling enforces anti-proliferative effects. Indeed, restoring SMAD4 expression in SMAD4-null pancreatic cancer cells reactivates p21 and growth arrest [28], whereas TGF-β cannot induce p21 in SMAD4-deficient cells [28].

However, TGF-β signaling is paradoxical in cancer: at advanced stages, tumor cells often lose responsiveness to TGF-β's suppressive effects, and TGF-β can then drive epithelial–mesenchymal transition, invasion, immunosuppression, and angiogenesis via SMAD-independent pathways [29]. Loss of SMAD4 is one mechanism by which tumors evade TGF-β's early brake on the cell cycle. Conversely, the absence of SMAD4 may also modulate late-stage behaviors – for example, altering patterns of metastasis (discussed below). SMAD4 itself is subject to regulation by other signaling pathways ("crosstalk"). Mitogen-activated protein kinase (MAPK) signaling (RAS/RAF/ERK) can phosphorylate SMAD2/3 at sites that prevent nuclear accumulation, promote SMAD4 degradation, and increase secretion of TGF-β1, thereby dampening or re-routing the pathway [30]. PI3K/AKT signaling can block TGF-β-induced cytostatic signals by inactivating downstream effectors like FOXO and mTOR [31] [32]. The Wnt/β-catenin pathway also intersects: SMAD7 (an inhibitory SMAD) can bind the AXIN–APC complex, linking SMAD signaling to β-catenin stabilization [32]. These interactions create a complex network where SMAD4 is a central node (Figure 2B).

Figure 2: SMAD4 structure and role in TGF-β/BMP signaling. (A) Domain schematic of SMAD4 protein: MH1 (DNA-binding) and MH2 (partner-binding) domains, with NLS (nuclear localization signal), NES (nuclear export signal), and linker region containing the SMAD Activation Domain (SAD) [21]. **(B)** SMAD4 as the common mediator in canonical TGF-β signaling: TGF-β ligands trigger type II and I receptor

phosphorylation of SMAD2/3, which then complex with SMAD4 and translocate to the nucleus to regulate target genes (e.g. cyclin-dependent kinase inhibitors for growth arrest) [33] [26]. Key points of crosstalk are shown: RAS/ERK can inhibit nuclear translocation of SMADs and promote SMAD4 degradation; PI3K/AKT/mTOR signaling can blunt TGF-β cytostatic effects; Wnt pathway components (AXIN/APC complex) interface with SMAD7 to influence β-catenin stability [30] [32]. (Adapted from Zhao *et al.*, 2018 [34] [25]).

In summary, SMAD4 is pivotal for canonical TGF-β/BMP signal transduction. Its loss-of-function can abrogate TGF-β's tumor-suppressive arm, contributing to uncontrolled proliferation. Yet, the consequences of SMAD4 inactivation are context-dependent and can be modulated by other pathways – a theme that recurs when comparing different tissues and stages of cancer. The sections below detail the historical discovery of SMAD4, its germline mutation syndromes, and its somatic roles in malignancies.

Historical Evolution of Knowledge

SMAD4's significance first came to light in the mid-1990s through studies of pancreatic cancer. In **1996**, Hahn *et al.* identified **DPC4** on chromosome 18q21.1 as a frequently deleted or mutated **tumor suppressor gene** in pancreatic ductal adenocarcinoma (PDAC) [20]. They reported biallelic inactivation of DPC4 in ~50% of PDAC tumors, coining the DPC ("Deleted in Pancreatic Carcinoma") nomenclature. This placed SMAD4 alongside KRAS, TP53, and CDKN2A as one of the core genetically altered drivers of pancreatic tumorigenesis. Shortly thereafter, researchers uncovered SMAD4's broader role in TGF-β signaling – it was found to be the human homolog of **Drosophila Mad** proteins, explaining its name "SMAD." In these early years, **dissenting evidence** emerged on whether SMAD4 loss alone could initiate cancer. For instance, mouse models with pancreas-specific Smad4 deletion did **not** develop spontaneous tumors, suggesting that SMAD4 loss promotes progression only in the presence of other oncogenic events [35] [36]. This aligns with the current view that SMAD4 acts as a **progression factor** (enabling malignant behavior when early drivers like KRAS are present) rather than an initiating oncogene.

By the late 1990s, attention turned to germline SMAD4 mutations. In **1998**, Howe *et al.* described families with **Juvenile Polyposis Syndrome (JPS)** caused by inherited SMAD4 mutations [37]. This was a breakthrough in understanding a rare hereditary GI polyposis condition. Almost simultaneously, another gene (BMPR1A, a BMP receptor) was also found mutated in JPS, indicating genetic heterogeneity. Over the next decade, clinicians noticed that a subset of SMAD4-mutant JPS patients also showed features of **Hereditary Hemorrhagic Telangiectasia (HHT)** – such as recurrent nosebleeds (epistaxis) and arteriovenous malformations. In **2004–2006**, several case reports and small series solidified this **JPS-HHT overlap syndrome**, unique to SMAD4 carriers [6]. Wain *et al.* (2014) systematically showed that SMAD4-mutant JPS patients had HHT symptoms (e.g. epistaxis, mucocutaneous telangiectases, AVMs) with high frequency, whereas BMPR1A patients did not [6] [7]. This expanded SMAD4's clinical significance beyond polyposis to include vascular pathology, prompting updated management guidelines (e.g. screening SMAD4 carriers for AVMs).

In the 2000s and 2010s, extensive work detailed SMAD4's role in **cancer biology**. Notably, **Iacobuzio-Donahue et al. (2009)** demonstrated that PDAC tumors stratified by DPC4/SMAD4 status exhibit distinct failure patterns after therapy [15]. Through rapid autopsies of PDAC patients, they showed **SMAD4-intact** tumors often pursued a "locally destructive" course (killing the patient via local invasion/complications), whereas **SMAD4-deficient** tumors more often caused widespread **metastatic** disease [15]. This finding – that SMAD4 genotype correlates with metastasis vs. local recurrence – was a milestone linking molecular genetics to clinical behavior. It also introduced the concept of using SMAD4 status as a **prognostic**

biomarker to inform treatment (e.g. consideration of more aggressive systemic therapy if SMAD4 is lost) [38]. Over the same period, multiple groups evaluated SMAD4 as a prognostic factor in other cancers, especially CRC. Early immunohistochemistry (IHC) studies in colorectal cancer in 2000–2010 suggested that loss of SMAD4 (or 18q LOH encompassing SMAD4) portended worse outcomes [39]. However, some inconsistencies appeared – for example, different studies yielded varying hazard ratios, likely due to differences in patient cohorts and co-mutations.

By the late 2010s, **meta-analyses** helped resolve some debates. A 2016 meta-analysis (Xing *et al.*) of 14 studies in PDAC confirmed that SMAD4 loss is associated with significantly shorter survival (pooled HR≈1.8 for OS) [40]. In 2021, another meta-analysis (Fang *et al.*) in CRC (10 studies, n=4394) found SMAD4 mutations conferred higher risk of death (HR~1.46) and recurrence (HR~1.59), and correlated with adverse features like nodal metastases and mucinous histology [41]. Meanwhile, large genomic consortia (TCGA, etc.) firmly established SMAD4 among the top mutated tumor suppressors in pancreatic and colorectal cancers, and documented its involvement (at lower frequency) in gastric, esophageal, lung, cholangiocarcinoma, and other cancers [42]. **Figure 1** below illustrates a representative PRISMA flow diagram from a recent meta-analysis of SMAD4 in PDAC, reflecting the growing body of literature synthesized in such analyses.

Figure 1: PRISMA flow diagram from a meta-analysis of SMAD4 in pancreatic cancer [3]. This illustrates the selection of 14 studies (n=1762 patients) from an initial 564 articles, after applying inclusion/exclusion criteria [40]. Such meta-analyses in PDAC [40] and CRC [41] have solidified the evidence that SMAD4 inactivation is an adverse prognostic factor. (Adapted from Xing *et al.*, 2016.)

Current Consensus and Open Questions: Today, SMAD4 is recognized as a **bona fide tumor suppressor** in multiple contexts. Germline testing for SMAD4 is standard in JPS and sequencing panels for hereditary GI cancer syndromes [43] [44]. In oncology practice, SMAD4 IHC or sequencing is often performed in PDAC and CRC; its loss may influence prognostic discussions and research trial stratification (though it does not yet dictate a specific therapy). OncoKB and other knowledge bases classify SMAD4 inactivating mutations as likely oncogenic (Tier 2) but **not directly targetable**. There remain **dissenting findings**: for instance, one study in resected PDAC paradoxically found no OS difference by SMAD4 status and higher local recurrence with SMAD4-intact tumors [16] (consistent with the local vs. distant failure pattern), underscoring that SMAD4's effect on survival can be modulated by therapy and competing risks. Another nuanced finding is that certain SMAD4 missense mutations might carry **neomorphic properties** – e.g. **R361** hotspot mutants (common in CRC) disrupt SMAD4's binding with SMAD2/3 (abrogating TGF-β signaling) but can still bind alternate partners like transcription factor LEF1, potentially **enhancing Wnt signaling** in those tumors [45] [46]. This raises the possibility that not all SMAD4 mutations are equivalent; most are loss-of-function, but some may endow tumor cells with alternative pro-tumorigenic pathways (an area of active research).

In summary, over 25 years the understanding of SMAD4 has evolved from a single-gene discovery in pancreatic cancer to a broad appreciation of its role in hereditary disease and solid tumor progression. The following sections detail germline SMAD4 syndromes and somatic SMAD4 alterations in cancer, integrating the historical and current evidence, and noting areas of consensus and controversy.

Germline SMAD4: Juvenile Polyposis and JPS-HHT Syndrome

Genetic Inheritance and Penetrance: Germline pathogenic variants in SMAD4 cause **Juvenile Polyposis Syndrome (JPS)**, an autosomal dominant disorder [47]. Each child of an affected individual has a 50% chance of inheriting the mutation [48]. Notably, ~50% of JPS cases are **de novo** (the proband has no family

history) [49], reflecting either new mutations or unrecognized mild disease in a parent. Penetrance is high but **variable** – nearly all mutation carriers develop some polyps by adulthood, but the number and timing vary widely [50] [51]. Even within a family, one person with SMAD4 mutation may have only a few polyps in their 20s, while another has hundreds by their teens [4] [52]. This variable expressivity suggests other genetic or environmental modifiers at play. There are no strong genotype–phenotype correlations; for instance, family members with the identical SMAD4 variant can have vastly different polyp burdens. One general observation is that **SMAD4 mutations** (as opposed to BMPR1A) are more likely to cause **upper GI polyposis** (stomach, duodenum) in addition to colonic polyps. Consequently, SMAD4-JPS tends to be more severe in the stomach (often requiring more interventions), whereas BMPR1A-JPS is usually confined to the colon/rectum. In terms of inheritance patterns, SMAD4-JPS shows classic autosomal dominance with high penetrance, but rare instances of **germline mosaicism** have been reported (parents phenotypically normal with an affected child) [53] [54]. Thus, genetic counseling for "sporadic" JPS should include the possibility of mosaic parental mutations and recommendations to screen siblings.

Clinical Phenotype – Juvenile Polyposis: JPS is characterized by development of multiple **juvenile polyps** (benign hamartomatous polyps with abundant stroma and cystic glands) in the gastrointestinal tract. "Juvenile" refers to the polyp histology, not necessarily the age of onset, though polyps often appear by the **teenage years** [50]. Diagnostic criteria for JPS include [55] any of: (a) ≥5 juvenile polyps in the colon/rectum; (b) multiple juvenile polyps throughout GI tract; or (c) any juvenile polyps with a family history of JPS. SMAD4 mutations account for roughly **27%** of JPS cases, and BMPR1A mutations ~28%, while ~45% have no mutation identified [56] [57]. SMAD4-JPS typically involves both **upper and lower GI polyps**. Polyps can occur from the stomach to the rectum; the colon is most commonly affected, but up to 50% of SMAD4 carriers have polyps in the stomach or duodenum as well. Patients often present with chronic GI bleeding (painless hematochezia) or anemia due to polyp ulceration [58]. Other symptoms can include abdominal pain, diarrhea, or intussusception if polyps are large. The polyp count ranges widely: some individuals have only a handful over their life, while others can develop hundreds (sometimes termed "JP of infancy" when polyps present in infancy with >100 polyps and protein-losing enteropathy) [50]. If untreated, polyps can cause complications such as bleeding, anemia, prolapse (if rectal), and rarely bowel obstruction. There is also a risk of **malignant transformation**: juvenile polyps themselves are benign, but JPS confers a predisposition to GI cancers arising from these polyposis areas.

Cancer Risk in JPS: Individuals with JPS have a markedly elevated risk of gastrointestinal malignancies. Estimates of the **lifetime GI cancer risk** in JPS have varied. Early reports from high-risk families showed alarming figures (up to 50–>80%), but these were likely overestimates due to ascertainment bias [4] [59]. More recent cohort studies suggest that with modern surveillance, the risk is more moderate. A 2020 analysis of the largest series of JPS patients (n= allelic carriers from literature and registries) found an overall cancer incidence of **15%** [60] [61]. This aligns with other contemporary studies after the implementation of endoscopic surveillance [4]. The **colon** bears the highest risk: historically, CRC risk was ~50% by age 60 in unscreened patients [62]. In one large family with SMAD4-JPS, the lifetime CRC risk was ~40% [63]. GeneReviews reports incidence of colorectal cancer ~17–22% by age 35, rising to ~68% by age 60 without intervention [62]. However, with proactive polyp removal, these numbers are likely attenuated. **Gastric cancer** is also a concern, particularly in SMAD4 carriers who often have heavy gastric polyposis. In patients with gastric involvement, the gastric cancer incidence is ~21% [62]. Strikingly, in SMAD4-JPS cohorts, up to 27% of all cancers were gastric, whereas BMPR1A-JPS patients rarely if ever developed gastric cancer (0% in one series) [7]. This reinforces the need to surveil the upper GI tract in SMAD4 carriers. Pancreatic cancer has been reported in a few JPS cases but appears to be a much lower risk than colon/stomach (no clear percentage, possibly slightly above population risk) [64]. The median age of cancer diagnosis in JPS is

around **40–45 years** (early for colon cancer, which in sporadic cases peaks in 60s) [62] [65]. In summary, SMAD4-JPS confers a substantial risk of CRC and gastric cancer by mid-life if no intervention; with surveillance, many of these cancers can be prevented (polyps removed before malignant change).

JPS–HHT Overlap (SMAD4-related HHT): A unique aspect of SMAD4 germline mutations is the combined **Juvenile Polyposis/Hereditary Hemorrhagic Telangiectasia (JPS/HHT)** syndrome. Approximately **20–30%** of patients with SMAD4 mutations exhibit signs of HHT [66]. HHT is a vascular dysplasia syndrome characterized by mucocutaneous telangiectasias and arteriovenous malformations (AVMs) in organs. In SMAD4 carriers, the HHT features can include **recurrent epistaxis** (nosebleeds from nasal telangiectases), which occurred in ~60–70% of SMAD4-JPS patients in one series [67]. **Telangiectases** on the skin, oral mucosa, or GI tract appear in ~57% [67], often after age 30 [67]. More ominously, large AVMs occur: **pulmonary AVMs** are found in over half of SMAD4 patients (53–81%) [68], which can cause right-to-left shunting of blood, leading to clubbing, low oxygen levels, or paradoxical emboli (stroke risk). **Hepatic AVMs** occur in ~38% [69] and can lead to high-output cardiac failure or portal hypertension in severe cases. **Cerebral AVMs** are less common (~4%) but can be life-threatening if they bleed [69]. Another vascular finding in SMAD4 carriers is **aortopathy**: up to 38% had aortic aneurysm or dilatation (especially root aneurysms) in one study [70] [71]. The median age for detecting aortopathy was 24 years, indicating it can manifest in young adults [70]. These features underscore that SMAD4 mutations affect TGF-β/BMP signaling in vasculature, not just the gut. Indeed, SMAD4 is involved in **angiogenesis and vascular integrity** (likely through BMP pathways), explaining the HHT phenotype overlap. It is noteworthy that HHT due to SMAD4 is clinically similar to HHT caused by ENG or ACVRL1 mutations, except that telangiectases of the **digits/nails** are less prominent in SMAD4-HHT [72] and the combination with polyposis is unique. From a clinical management perspective, any patient diagnosed with JPS due to SMAD4 should be evaluated for HHT signs (and vice versa), as this has implications for screening and interventions (e.g. treating AVMs to prevent hemorrhagic stroke or brain abscess).

Surveillance and Management in SMAD4 Carriers: Given the substantial cancer risks and HHT complications, aggressive surveillance is recommended for SMAD4 mutation carriers. Current guidelines (e.g. **GeneReviews** and GI society recommendations) suggest starting surveillance by the **teenage years**. Specifically, **colonoscopy and upper endoscopy** should begin by age **15** (or earlier if symptoms) and be repeated **every 3 years** if no polyps are found [73] [9]. If polyps are detected and removed, endoscopies are done **yearly** until the patient is polyp-free, then can revert to 3-year intervals [9]. For individuals with unknown mutation but a family history, a somewhat relaxed interval (every 5 years in adulthood) may be used if no polyps on initial exam [74]. During these endoscopies, polyps larger than a few millimeters are typically removed (polypectomy) to reduce polyp burden and cancer risk. If polyps are too numerous or histologically concerning (high-grade dysplasia), consideration is given to prophylactic surgery. **Colectomy** (partial or total) is indicated if the colon polyp burden becomes unmanageable endoscopically (e.g. hundreds of polyps or significant dysplasia) [75]. Often a **subtotal colectomy with ileorectal anastomosis** is favored to preserve continence, unless rectal involvement is severe requiring proctocolectomy [75]. Similarly, a **partial or total gastrectomy** might be recommended if there are diffuse, recurrent gastric polyps with dysplasia that cannot be adequately controlled endoscopically [76]. These major surgeries are life-altering and not undertaken lightly; the goal is to avoid them with diligent endoscopic management. In practice, a significant minority of SMAD4-JPS patients do end up needing colorectal or gastric resection in mid-adulthood, but exact frequencies vary (recent series report perhaps 20–30% eventually require some GI surgery) [77]. It has been observed that SMAD4 carriers, because of aggressive gastric polyposis, more often require gastric surgery than BMPR1A carriers. One study noted that of 27 SMAD4 patients under surveillance, 4 still developed cancer (despite some having had surgery) [52] – highlighting that even with

interventions, vigilance is required. Beyond endoscopy, **HHT-related surveillance** is crucial: guidelines advise that SMAD4 mutation carriers get a **transthoracic echocardiogram with bubble study** in childhood to screen for pulmonary AVMs (which manifest as an intrapulmonary shunt) [10]. If the bubble echo is positive or if clinical suspicion is high, confirmatory imaging (CT or MRI of chest) is done, and treatable AVMs (>2–3 mm feeding artery) can be **embolized** prophylactically to prevent paradoxical emboli or hemoptysis. Screening MRI of the brain in adolescence is often recommended to detect cerebral AVMs that could be neurosurgically treated to prevent hemorrhagic stroke [78] [79]. Routine **epistaxis management** is also part of care – patients are counseled on humidification, topical hemostatic therapies, and to avoid NSAIDs that exacerbate bleeding [80] [81]. If nosebleeds are severe, laser ablation of nasal telangiectases can be performed (though repeated procedures are often needed). Caution is urged against certain interventions: for example, **liver biopsy is contraindicated** in SMAD4-HHT patients due to risk of catastrophic bleeding from hepatic AVMs [82].

Quality of Life in SMAD4-JPS/HHT: Living with SMAD4 mutation can impose significant burdens. Starting in adolescence, patients undergo frequent invasive procedures (endoscopies every 1–3 years, numerous polypectomies). While these are done under sedation, the sheer frequency can cause anxiety and discomfort. Many patients have chronic iron-deficiency **anemia** from polyps or epistaxis, requiring iron supplements or even transfusions. In SMAD4 carriers, anemia was twice as prevalent as in BMPR1A patients (58% vs 26%) [7], reflecting the heavier bleeding from GI polyps and HHT. Chronic anemia and repeated bleeding episodes (e.g. weekly nosebleeds) can reduce energy levels and QoL. If major surgeries like colectomy or gastrectomy are needed, patients must adjust to altered gastrointestinal function (increased stool frequency, dietary modifications, risk of short gut if extensive). Those with gastrectomy may have lifelong impacts on nutrition and require vitamin B12 injections. Psychologically, knowing one's high cancer risk can be stressful, though successful surveillance often prevents cancer and provides some reassurance. It is worth noting that with current management, many SMAD4 mutation carriers **can lead long lives** – unlike untreated historical cases that might succumb to cancer by their 40s. For example, one report mentioned that median survival in a cohort of treated JPS patients had improved substantially compared to older data (no precise number given, but clearly improved with surveillance) [37]. Nonetheless, robust quantitative QoL studies in JPS are lacking. Patients often report **procedure-related anxiety**, **dietary limitations** (to avoid polyp-provoked symptoms), and the psychosocial impact of living with a hereditary syndrome (including guilt about passing it to children). Support from genetic counselors and connecting with others via patient networks can be beneficial.

In summary, germline SMAD4 mutations cause a **multi-system syndrome** requiring coordinated care: gastroenterologists for polyposis, interventional radiology or pulmonology for AVMs, cardiology for high-output states, ENT for epistaxis, and genetics for family screening. The evidence base for management is strong (based on decades of case series and evolving guidelines), but some uncertainties remain – for example, the true residual cancer risk under optimal surveillance, or the best age to consider prophylactic surgery. Ongoing collection of data in registries (such as the EU-funded JPS registries [83] [84]) will hopefully refine these estimates. There is also interest in **chemoprevention** for JPS (as exists for adenomatous polyposis with NSAIDs), but so far no agents have proven effective [85]. An intriguing development is the use of **mTOR inhibitors (sirolimus)** in the rare infantile-onset JPS (caused by PTEN/BMPR1A double deletions) to reduce polyp burden [86]. It's speculative, but similar pathway-based therapies might one day help in SMAD4-JPS if, for instance, a drug could enhance TGF-β signaling or target downstream effects. For now, **early detection and intervention** remain the cornerstones that have significantly improved outcomes for SMAD4 mutation carriers.

Somatic SMAD4 in Cancer

Somatic (acquired) alterations of SMAD4 occur across a spectrum of malignancies, most prominently in gastrointestinal cancers. As a **tumor suppressor gene**, SMAD4 is typically inactivated via **loss-of-function** point mutations (nonsense or frameshift mutations causing truncation), **deletions** (partial or whole gene), or loss of protein expression (often reflecting biallelic genetic loss). **Missense** mutations also occur, especially clustering in the MH2 domain which is critical for SMAD4's binding to partner SMADs [87] [88]. Many missense mutations in MH2 disrupt heterodimerization with SMAD2/3, effectively abolishing canonical signaling [89] [46]. In general, a single SMAD4 mutation (with the other allele lost by deletion or a second hit) leads to loss of SMAD4 function in the tumor cell. Below we review SMAD4's role by cancer type, focusing on prevalence, co-mutation patterns, prognostic impact, and treatment correlations. We prioritize **pancreatic, colorectal, gastric, and lung** cancers (where SMAD4 has been most studied), and then briefly note other tumor types with significant SMAD4 involvement (e.g. cholangiocarcinoma).

Pancreatic Ductal Adenocarcinoma (PDAC)

Prevalence and Types of Alterations: SMAD4 is inactivated in roughly **50%** of PDACs by the time of diagnosis [1]. It is one of the "big 4" mutated genes in pancreatic cancer (after KRAS (~90%), TP53 (~70%), and CDKN2A (~50%)). The inactivation typically involves **biallelic loss**: one allele may have a point mutation (often a truncating mutation), while the other is lost via deletion or loss of heterozygosity. In some cases, there is **homozygous deletion** of the SMAD4 locus. Immunohistochemical loss of SMAD4 protein is highly concordant with underlying genetic SMAD4 loss [90] – pathologists often use DPC4 IHC as a surrogate marker. Notably, SMAD4 mutations are usually **late** events in PDAC tumorigenesis: in PanIN (pancreatic intraepithelial neoplasia) precursor lesions, SMAD4 is typically intact until the high-grade stage, whereas KRAS and CDKN2A mutate earlier. This timing aligns with SMAD4's role in progression rather than initiation. The types of SMAD4 mutations in PDAC include nonsense mutations (e.g. **p.Arg361** *is a recurrent truncation), frameshifts, and missense changes in MH2 domain (e.g. at residues like Asp351, Pro356, Arg361). The MH2 domain is the mutational hotspot region* [87] [88]*, as it mediates SMAD4-SMAD2/3 interaction – mutations here (whether truncating or missense) often abrogate complex formation. Figure 3A shows a lollipop plot of SMAD4 mutations from a PDAC cohort (MD Anderson Cancer Center series), illustrating the clustering of alterations in the MH2 domain (exons 8–11)* [91] [92]*. Co-occurring genetic events in SMAD4-deficient PDAC frequently include KRAS activation and TP53 mutation (nearly all SMAD4-mutant PDACs have KRAS/TP53 mutations as well* [93]*). Interestingly, there is an inverse correlation between SMAD4 and SMAD3/SMAD2**** mutations – since SMAD4 is downstream of SMAD2/3, usually a tumor inactivates one or the other, but not both (a form of synthetic lethality; SMAD2/3 mutations are much rarer than SMAD4 in PDAC). Copy-number loss of chromosome 18q (SMAD4 locus) is common; often the entire 18q arm is lost, sometimes affecting other tumor suppressors too. About half of PDACs therefore have complete absence of SMAD4 protein, whereas the others retain SMAD4. This dichotomy has been studied for clinical correlations.

Figure 3: SMAD4 mutation spectrum in gastrointestinal cancers. (A) Distribution of SMAD4 mutations in a pancreatic cancer cohort (MD Anderson, n=49 with SMAD4 sequencing) [94] [91]. Each lollipop indicates a mutation; note clustering in the MH2 domain (right side), including recurrent truncating mutations (e.g. at Arg361, asterisk) and missense mutations (e.g. p.P356 and p.R361 variants – common "hotspots" in PDAC and CRC) [91] [92]. **(B)** SMAD4 mutations in colorectal cancer (TCGA dataset, n=220) [95] show a similar MH2 clustering. Missense mutations (blue) often affect the SMAD4–SMAD2/3 interface, while nonsense/frameshift (red) cause loss of function. (Data from Sarshekeh *et al.*, 2017 [91] [92], with permission under CC-BY license.)

Prognostic Impact in PDAC: SMAD4 status is one of the most studied prognostic biomarkers in pancreatic cancer. The **preponderance of evidence** indicates that SMAD4 inactivation is associated with more aggressive disease and worse survival, although not all studies agree on the extent. The seminal Iacobuzio-Donahue study (2009) revealed the link between SMAD4 and **metastatic propensity**: in their autopsy series, 75% of SMAD4-negative PDACs had widespread metastases at death versus only 14% of SMAD4-positive tumors [15]. SMAD4 loss was significantly correlated with the **metastatic failure** pattern (P=0.007) [15]. This translates clinically to the observation that SMAD4-negative patients often develop liver and lung metastases early, whereas SMAD4-intact patients might recur locally (e.g. anastomotic recurrence after surgery) without immediate distant spread [15] [16]. Such locally confined recurrences can still be lethal in PDAC (due to pain, biliary obstruction, etc.), but they might have slightly longer post-progression survival than widely metastatic cases. Large retrospective series have consistently found that **absence of SMAD4 (by IHC)** is associated with shorter median survival in both resected and advanced PDAC. For example, a Johns Hopkins cohort found median survival ~9 months in SMAD4-null vs 16.5 months in SMAD4-positive resected periampullary cancers [96] [97]. A meta-analysis by Zhang *et al.* (2018) encompassing 17 studies confirmed SMAD4 loss correlates with poorer overall survival (pooled HR ~1.74) and also poorer disease-free survival [98] [99]. Another meta-analysis (Xing *et al.*, 2016) including 1762 patients calculated a combined HR for OS of **1.20** (95%CI 1.03–1.40) on univariate analysis and **1.88** (1.31–2.70) on multivariate analysis [3] [40]. The strengthening of effect after adjustment suggests SMAD4 loss is an independent prognostic factor, not just a surrogate for other high-risk features. **Figure 4** shows the forest plot of overall survival from that meta-analysis, highlighting the elevated hazard associated with SMAD4-negative tumors. Notably, some heterogeneity was observed: subsets of studies (e.g. smaller ones, or those with different follow-up durations) had varying results [100]. But there was no publication bias detected, reinforcing confidence in the finding [101].

Figure 4: SMAD4 loss and overall survival in pancreatic cancer (meta-analysis). Pooled hazard ratios (HR) for OS comparing SMAD4-negative versus SMAD4-positive PDAC, from Xing *et al.* (2016) [3] [102]. **(A)** Univariate analysis across 11 studies (n=1401) shows HR ~1.20 (95% CI 1.03–1.40) [3]. **(B)** Multivariate analysis (8 studies, n=927) shows adjusted HR ~1.88 (95% CI 1.31–2.70) [102]. This indicates patients with SMAD4-inactivated tumors have significantly worse survival, even after accounting for stage and other factors. Diamond shapes = summary HR, boxes = individual study HR, horizontal lines = CI. (Data from [3])

It must be noted that **not all studies have found SMAD4 to be an independent prognosticator**. For instance, in some cohorts of resected PDAC, once stage, margin status, and therapy are accounted for, SMAD4's impact on OS diminishes (HR ~1.2, p = 0.1 in one multivariate model) [103] [104]. The Mayo Clinic study by Gits *et al.* (2021) even found **no difference in overall survival** by SMAD4 status among patients who had upfront surgery and adjuvant therapy [16]. In that study, 92% of patients were SMAD4-negative (only 8% retained SMAD4, which is lower than typical, possibly reflecting selection bias for advanced tumors), and survival was uniformly poor (~2.1 years median for both groups) [105] [16]. However, they did find **locoregional recurrence** was 7–9 times higher in SMAD4-positive patients [16] [106], consistent with the "local vs distant" paradigm. These nuances suggest that while SMAD4 loss generally portends a worse prognosis, its effect can be overshadowed by other variables in certain contexts (for example, if everyone has metastatic disease, then having or not having SMAD4 may not further stratify survival because outcomes are uniformly poor). Nonetheless, the broad consensus is that **SMAD4 status provides prognostic information**: SMAD4-intact PDAC might follow a more indolent or local trajectory, whereas SMAD4-null PDAC indicates a higher likelihood of rapid systemic spread. This has practical implications: some have proposed using SMAD4 to guide therapy – e.g., if SMAD4 is intact (and thus risk of isolated local recurrence high), one might more strongly consider radiotherapy to address the primary tumor bed [107].

Conversely, if SMAD4 is lost, emphasis might shift to systemic therapy early, given micrometastases are likely. This concept is still being evaluated; it's not yet standard of care to alter treatment solely on SMAD4, but it is a subject of prospective studies.

Predictive/Treatment Response in PDAC: Unlike some molecular markers (e.g. BRCA1/2 in pancreatic cancer predicting platinum response), SMAD4 has no direct targeted therapy. There is no "SMAD4 restoration" drug, nor a way to directly kill SMAD4-deficient cells currently. However, research indicates SMAD4 loss might influence response to certain treatments. For instance, one study found that SMAD4-negative PDACs had higher rates of **distant progression** even after adjuvant chemoradiation, implying that systemic therapy should be prioritized [108]. Another study suggested patients with SMAD4-negative tumors may not benefit as much from 5-FU-based adjuvant therapy as SMAD4-positive patients [109] [110] – specifically, a cohort of 202 patients showed those with intact SMAD4 got a 3-fold greater survival benefit from 5-FU chemo than those with SMAD4 loss [39] [110]. This hints that SMAD4-null tumors (which often have mesenchymal, metastatic phenotype) might be more chemoresistant. Indeed, SMAD4 loss has been linked to chemoresistance mechanisms (e.g. through promoting epithelial-mesenchymal transition and therapy-resistant cell subpopulations) [111]. On the other hand, a tantalizing preclinical finding is that SMAD4-deficient pancreatic cancers may be more susceptible to **PARP inhibitors or DNA-damage agents** due to a role of SMAD4 in DNA repair: knockdown of SMAD4 limited DNA repair in PDAC cells, increasing sensitivity to PARP inhibition in a recent study [112] [113]. This is early research but suggests synthetic lethal approaches could be developed. Additionally, **TGF-β pathway inhibitors** (like **galunisertib**, a TGF-β receptor I kinase inhibitor) have been tested in PDAC. A Phase II trial of galunisertib + gemcitabine showed a modest improvement in survival (median 8.9 vs 7.1 months vs gemcitabine alone) [19]. While not a game-changer, it indicates that blunting TGF-β signaling might help, especially in patients with intact SMAD4 where TGF-β's pro-tumor effects remain active. In SMAD4-null tumors, TGF-β signals cannot signal through SMAD4, but they can still signal via non-canonical pathways; whether TGF-β inhibitors work preferentially in SMAD4-null or intact contexts is unclear. Trials did not select by SMAD4 status.

Co-mutations and Metastatic Patterns: In PDAC, SMAD4 loss often co-occurs with **TP53 mutations** and **KRAS**. There is some evidence that the **metastatic pattern** differs: SMAD4-null PDAC more frequently metastasizes to distant organs like lung and liver [15], whereas SMAD4-retained tumors might remain confined to the pancreas/locoregional area (though they eventually prove fatal by local invasion). One autopsy series described two phenotypes: "local spreaders" (DPC4 intact) and "distant spreaders" (DPC4 lost) [15]. Clinically, this could be seen in the difference between a patient who recurs with isolated local mass vs. one who recurs with multiple liver lesions shortly after surgery. This has been partly validated in contemporary series. A 2015 study found that SMAD4 expression was associated with higher locoregional recurrence (HR 7.0) and a trend toward lower distant metastasis (P=0.06) in resected PDAC [16]. Thus, SMAD4 status might predict **where** the disease will come back, even if it doesn't vastly change **if** or **when** it will come back (since PDAC overall has high recurrence rates regardless). Another interesting co-mutation pattern is with **BRCA2** and **PALB2**: some PDAC with DNA repair mutations also have SMAD4 loss, but there's no clear functional interaction known there, other than both being common alterations in advanced PDAC.

In summary for PDAC: loss of SMAD4 is very common and generally indicates a **worse prognosis**, especially due to early metastasis. It is a robust prognostic marker confirmed by meta-analyses [102]. Its use in clinical decision-making is evolving, with ongoing research into whether SMAD4 can guide local vs systemic therapy focus. No therapies directly target SMAD4 loss yet, but pathway approaches (TGF-β inhibitors) and synthetic lethal strategies are under exploration. Importantly, about half of PDAC patients still have intact SMAD4 – understanding why their tumors did not inactivate SMAD4 (and whether those tumors rely on TGF-β

signaling for invasion) is an area of current investigation, as it could open up therapeutic vulnerabilities in SMAD4-wildtype PDAC (perhaps using TGF-β inhibitors in that subset).

Colorectal Cancer (CRC)

Prevalence and Genetic Context: SMAD4 is mutated in approximately **10%** of all colorectal adenocarcinomas, with reported frequencies ranging from ~5% to 20% in different studies [114] [2]. TCGA data showed ~10% mutation rate [2], while some single-institution series using hotspot panels found ~12% in metastatic CRC [115]. The range (5–20%) can be explained by stage (advanced cancers have higher frequency of SMAD4 loss), site (colon vs rectum differences), and assay sensitivity. In addition to point mutations, **18q deletions** (loss of heterozygosity at SMAD4 locus) are common in CRC – historically, 18q LOH was a marker of poor prognosis in stage II/III colon cancer, likely due to SMAD4 and nearby genes being lost [116]. Indeed, reduced SMAD4 protein expression is seen in a higher fraction of CRC (~30%) than mutation frequency, indicating some cases have loss via deletion or epigenetic silencing [117]. A significant proportion of **microsatellite stable (MSS)**, CIN (chromosomal instability) colorectal cancers undergo 18q loss late in adenoma-carcinoma sequence, contributing to invasion and metastasis. By contrast, **MSI-H (microsatellite unstable)** CRC rarely have SMAD4 mutations; they more often disrupt TGF-β signaling by frameshifting the TGF-β type II receptor (TGFBR2) and typically retain 18q (since MSI tumors are diploid). Thus, SMAD4 mutations are enriched in **MSS** tumors. There is also an association with **mucinous histology** – the meta-analysis by Fang *et al.* found CRC with SMAD4 mutations had OR ~2.23 for being mucinous type [118]. Mucinous CRC often have concurrent KRAS mutations and 18q loss, which fits this profile. Another correlation is tumor location: SMAD4 mutations were modestly more common in **colon** (especially proximal colon) than in rectal cancers [118] [115]. One study showed an OR ~2.85 for SMAD4 mutation in colon vs rectum [119] [120]. This might reflect different molecular pathways in rectal cancer (which less frequently go through the TGF-β pathway inactivation, or perhaps selection biases in studies). SMAD4-mutant CRC frequently co-occur with **KRAS** or **BRAF** mutations. Fang *et al.* reported an OR ~2.13 for RAS mutation in SMAD4-mutants [17]. Similarly, Sarshekeh *et al.* found SMAD4 mutations clustered in the **CMS3 "metabolic" subtype** of CRC, which is known to have high KRAS mutation rates [121] [122]. Mechanistically, this makes sense: KRAS-driven tumors benefit from evading TGF-β suppression via SMAD4 loss. Also, **mucinous** histology CRC (which SMAD4 mutations predispose to) is often KRAS-mutated and CIMP-high. Interestingly, SMAD4 mutation was *not* associated with **MSI** status [17] (consistent with MSI using a different route to disrupt TGF-β signals). In terms of specific mutations, CRC shares hotspots with PDAC: e.g. **p.Pro356Thr** and **p.Arg361His/Cys** in MH2 domain are recurrent missense changes [123] [91]. These can act in dominant-negative fashion or partially alter SMAD4 function. Truncating mutations (nonsense or frameshift) are also common, often leading to loss of the C-terminal MH2 (thereby abrogating complex formation). Homozygous deletions of SMAD4 are rare in CRC (less common than in PDAC), likely because copy-neutral LOH plus point mutation suffices.

Prognostic Impact in CRC: Multiple studies have identified SMAD4 loss as a **poor prognostic factor** in colorectal cancer, especially in the metastatic setting. The 2021 systematic review (Fang et al.) concluded SMAD4 mutation correlates with significantly worse **overall survival** (HR 1.46) and **progression-free survival** (HR 1.59) [41]. In practical terms, one large single-center study (MD Anderson, n=734 metastatic CRC patients) found median OS 29 months in SMAD4-mutants versus 56 months in SMAD4-wildtype (HR ~2.08, p<0.001) [115] [14]. That is a substantial difference, on par with or greater than the impact of many traditional prognostic factors. However, after adjusting for other variables, the independent effect can attenuate. In the MD Anderson cohort, multivariate analysis showed SMAD4 mutation with HR 1.39 and a p-value of 0.10 [103] [104], suggesting some of the effect overlapped with other factors (e.g. poor differentiation,

which was also more common in SMAD4-mutants). Still, numerous independent cohorts (including population-based ones) have corroborated the adverse prognosis. The **Association of SMAD4 with metastasis** is of note: several studies found SMAD4 loss is linked to higher risk of **liver metastasis** and generally stage IV disease [118]. One older concept was that **18q LOH** (surrogate for SMAD4 loss) was a marker of poor outcome in **stage II colon cancer**, suggesting those patients might benefit from adjuvant chemo. This was borne out in some analyses: stage II patients whose tumors had lost SMAD4 expression fared worse than those retaining it [124]. That hints that even at earlier stages, SMAD4 status might inform prognosis. It's worth noting a potential **dissenting finding**: at least one study (O'Connell et al., 2015) did not find SMAD4 IHC to be prognostic in a multivariate model for stage III colon cancer, whereas 18q LOH was (the discrepancy possibly due to IHC technical issues) [125]. But the weight of evidence favors SMAD4 loss as unfavorable. The mechanism is intuitive: without SMAD4, the TGF-β pathway cannot induce growth arrest or immune modulation in the tumor, potentially leading to more rapid growth and immune evasion. Indeed, a recent analysis showed **SMAD4-deficient CRC had an immune-evasive gene signature** (upregulating CXCL1, etc., to exclude T cells) [126], consistent with more aggressive behavior.

Therapeutic and Predictive Relevance in CRC: Perhaps the most clinically actionable aspect of SMAD4 in CRC is its role as a **predictor of resistance to EGFR inhibitor therapy**. In metastatic CRC, anti-EGFR monoclonal antibodies (cetuximab, panitumumab) are effective only in RAS/BRAF wild-type tumors. However, even among those, a significant subset doesn't respond. Several studies have identified SMAD4 loss as a marker of poor response to anti-EGFR. Sarshekeh et al. looked at a subset of 44 patients wild-type for KRAS/NRAS/BRAF who were treated with anti-EGFR; those with SMAD4 mutations had a **shorter progression-free survival** (median 111 days) compared to those with intact SMAD4 (180 days) [127] [128]. This was a significant difference (p=0.003). The implied hazard ratio was ~3 for progression on anti-EGFR if SMAD4 was mutated [128] [129]. Similarly, other studies (e.g. an analysis by Loupakis et al.) showed loss of SMAD4 (by IHC or mutation) predicted lack of benefit from EGFR antibodies in RAS/BRAF WT patients [130]. The biological rationale: SMAD4 loss often coexists with KRAS mutation, but even independent of RAS, SMAD4 deficiency might activate parallel pathways (like Wnt or MAPK via feedback) that make the tumor **intrinsically resistant** to EGFR blockade. Some guidelines or expert opinions now suggest that **SMAD4 status could be used as a negative predictive marker** – meaning if a tumor is RAS/BRAF WT but SMAD4-null, one might anticipate less benefit from cetuximab/panitumumab and consider other options. This is not yet standard (unlike RAS testing, which is mandatory, SMAD4 is not routinely tested for therapy selection), but it's a consideration in difficult cases. There is also emerging evidence that SMAD4 loss may confer relative **resistance to fluoropyrimidine chemo**. As mentioned, 5-FU adjuvant benefit seemed attenuated in SMAD4-lost tumors [109]. Preclinically, SMAD4 inactivation in colon cancer cells can induce chemo resistance by enabling EMT and changes in drug metabolism. Conversely, one could speculate SMAD4-deficient tumors might respond better to **immune checkpoint inhibitors** since they often have immune evasion patterns – however, since most SMAD4-deficient CRC are MSS (immune cold), checkpoint inhibitors are generally ineffective there.

Co-mutations and Patterns: In CRC, SMAD4 mutation is part of the canonical progression in a subset of tumors: APC mutation (adenoma) → KRAS (intermediate) → SMAD4/TP53 (late, invasive carcinoma). Thus SMAD4 alterations often come with KRAS and sometimes TP53. The meta-analysis found OR ~1.42 for lymph node metastasis in SMAD4-mutants [118], indicating those tumors were more often node-positive at diagnosis. They were also more often Stage IV (OR ~1.28) [118]. There isn't a known specific metastatic pattern unique to SMAD4 in CRC (unlike PDAC's local vs distant dichotomy). But given their aggressive nature, SMAD4-mutant CRC probably metastasize earlier and possibly to multiple sites. One study did suggest that **SMAD4/TP53 co-loss** defines a particularly bad subset – patients whose tumors lost both had

much shorter survival than those losing only one or neither [131]. Also, interestingly, patients with **SMAD4 wild-type + KRAS wild-type** had about double the survival of those with **SMAD4 loss + KRAS mutant** (83.8 vs 36.7 months) in a combined analysis [132]. This underscores how combination of pathways (RAS and TGF-β) yields especially aggressive disease.

In summary for CRC: SMAD4 is mutated in a minority of cases but tends to denote a **high-risk phenotype** – often right-sided, mucinous, RAS-mutant tumors with a tendency for metastasis and poor response to certain therapies. It is a potential biomarker for treatment stratification (pending further validation). The evidence is bolstered by meta-analyses and large cohorts, although one must be cautious that some of the prognostic impact may correlate with other high-risk features. Nonetheless, SMAD4 status is frequently reported in molecular pathology of CRC and discussed in tumor boards for its prognostic implications.

Gastric Cancer

Prevalence: SMAD4 is less frequently altered in gastric adenocarcinoma than in PDAC or CRC, but still significant. Reported mutation rates range from ~5% to 15% in gastric cancer, depending on the cohort (higher in high-stage tumors). The TCGA stomach cancer analysis (2014) found SMAD4 mutations in ~8% of cases. An Asian cohort study reported ~16% frequency of SMAD4 genetic alterations (including mutations or homozygous deletions) [133]. More commonly, **loss of SMAD4 protein** expression is observed in gastric cancers (via IHC) due to a combination of mutations and deletions. For example, one study noted **LOH at 18q/SMAD4** in ~30% of gastric cancers and correlated it with advanced stage [133]. SMAD4 inactivation may be enriched in certain histologic subtypes: the **diffuse-type** gastric cancers (including signet ring carcinoma) often have CDH1 mutations and less frequently SMAD4 loss, whereas the **intestinal-type** (with gland formation) coming from the adenoma pathway might more often inactivate SMAD4 as they progress. Additionally, chronic inflammation settings (like gastritis leading to carcinoma) can involve TGF-β pathway changes. But data on subtype specificity is limited.

Prognosis: Several studies indicate that loss of SMAD4 in gastric cancer is an adverse prognostic marker. Kim *et al.* (2015) found that reduced SMAD4 expression by IHC was associated with poorer survival and was an **independent prognostic factor** on multivariate analysis [134] [135]. Similarly, a Chinese study observed that SMAD4 gene loss (via deletion or mutation) correlated with deeper invasion and nodal metastases, and patients with SMAD4-negative tumors had significantly worse 5-year survival [133]. For instance, 5-year survival was ~34% in SMAD4-negative vs ~62% in SMAD4-positive tumors ($p<0.01$) in one series [136]. On multivariate analysis, SMAD4 loss remained an independent predictor of mortality (often alongside p53 status). There is also evidence linking SMAD4 loss to **recurrence patterns** in gastric cancer. An interesting finding by Zou *et al.* was that SMAD4-negative gastric cancers more frequently had **peritoneal metastasis** and recurrence, whereas SMAD4-retained tumors had more hematogenous metastasis [137]. If validated, that could parallel the local vs distant theme seen in PDAC, albeit in a different way (peritoneal spread vs bloodborne). Mechanistically, SMAD4 deficiency in gastric cancer cells was shown to promote an immune-evasive phenotype (via CXCL1 upregulation attracting immunosuppressive cells) [126], which might contribute to metastasis. Another mechanistic angle: in Helicobacter pylori-associated gastric carcinogenesis, TGF-β/SMAD signaling seems to act as a tumor suppressor restraining inflammation-associated proliferation, so SMAD4 loss might accelerate the inflammation-cancer transition.

Therapy Considerations: There is no specific therapy targeting SMAD4 in gastric cancer either. However, given that TGF-β is often overactive in advanced gastric cancer (promoting fibrosis and immune evasion), some trials have looked at TGF-β inhibitors in this disease too. One translational insight is that SMAD4 loss

might synergize with **FGFR2 pathway** (since SMAD4-null gastric lines showed upregulation of FGF/FGFR autocrine signaling) – thus FGFR inhibitors could be a strategy in SMAD4-deficient gastric tumors (still hypothetical). Clinically, SMAD4 status is not routinely tested in gastric cancer patients, but it could be in the future for risk stratification. For example, if a resected gastric cancer is SMAD4-negative, one might predict a higher chance of relapse and consider more aggressive adjuvant therapy or trials. Conversely, SMAD4-intact might have a slightly better prognosis with standard therapy. One study suggested SMAD4 expression could predict benefit from 5-FU chemo in gastric cancer, analogous to CRC, but data are sparse.

In summary, while not as extensively studied as in PDAC/CRC, SMAD4 appears to be a **tumor suppressor in gastric cancer** as well, with its loss denoting a worse outcome. The evidence is moderate (mostly single-institution studies) but consistently points that way. Gastric cancers often have concurrent alterations (TP53, etc.), so dissecting independent effect requires larger datasets. As comprehensive molecular profiling of gastric cancer continues, SMAD4 might join the panel of markers considered for personalized approaches.

Non-Small Cell Lung Cancer (NSCLC)

Prevalence: SMAD4 mutations occur in approximately **4–8%** of NSCLC, predominantly in **lung adenocarcinoma** subtype [138] [139]. They are relatively rare in lung squamous cell carcinoma. A 2021 study reported an alteration frequency of ~4% in NSCLC (with point mutations making up over half of those, and the rest deletions) [140]. While this is a small fraction, given the incidence of lung cancer, it's not negligible. SMAD4 is not among the top tier of mutated genes in lung adenocarcinoma (which are TP53, KRAS, KEAP1, STK11, EGFR, etc.), but it is a known event in a subset. Co-occurring mutations often include KRAS (in smokers) or sometimes EGFR (in which case SMAD4 mutation might emerge as a resistance or progression event). Copy-number loss of chromosome 18q can also hit SMAD4 in lung cancers. Beyond mutations, **reduced SMAD4 protein expression** is seen in a higher proportion of NSCLCs – up to ~30% in some IHC studies [117]. This suggests other mechanisms (transcriptional downregulation or protein degradation pathways) might reduce SMAD4 function in lung tumors.

Clinical Impact: Recent evidence suggests SMAD4 mutation/deficiency is a **negative prognostic factor in NSCLC**. Wang *et al.* (Lab Invest 2021) performed an integrated analysis and found NSCLC patients with SMAD4 mutations had significantly worse survival [141]. SMAD4 mutation was an independent risk factor for shorter OS, even after controlling for stage and other gene mutations [141]. In their data, SMAD4-mutant cases tended to present at higher stage and had features of aggressive biology. Moreover, they discovered that **SMAD4 expression was inversely associated with chemo-resistance**: patients whose tumors had low SMAD4 expression showed poorer disease-free survival and were less responsive to platinum-based chemotherapy [12]. This implies SMAD4-intact tumors might respond better to standard chemo, whereas SMAD4-deficient tumors might be more chemo-refractory, perhaps due to enhanced DNA repair or other escape mechanisms in the absence of SMAD4. Supporting this, SMAD4 knockdown in lung cancer cell lines increased their invasiveness and altered gene profiles toward a metastatic phenotype [142]. There is also an interesting interplay with the **TGF-β pathway in lung**: lung adenocarcinomas often experience TGF-β-driven EMT which promotes metastasis; if SMAD4 is intact, TGF-β can induce a full EMT program, but if SMAD4 is lost, cells might partially escape TGF-β growth inhibition but also not fully undergo EMT – this dichotomy is still being studied. Some mouse models (Kras^G12D^; Smad4^−/−^) developed lung tumors that paradoxically had **higher incidence of metastasis** compared to Smad4^+/+ tumors in one report [143], although Smad4 loss alone did not initiate lung cancer.

Therapeutic Context: There is no targeted therapy for SMAD4 in lung cancer either, but one notable angle is the relationship with the **DNA damage response**. A Nature 2022 study found SMAD4-deficient lung cancer cells had impaired NHEJ DNA repair and were more sensitive to PARP inhibitors and platinum [112]. If validated, SMAD4 loss could serve as a biomarker to use PARP inhibitors beyond BRCA mutations. In immunotherapy context, SMAD4's role is unclear – one might hypothesize that SMAD4-deficient tumors, being more mesenchymal, could be "cold" tumors less infiltrated by T cells, hence potentially less responsive to PD-1 inhibitors. Some preliminary data from Wang et al. showed SMAD4 mutation was associated with an **immune-suppressive tumor microenvironment** in NSCLC (via gene set enrichment) [141] [144], which could mean poorer immunotherapy response. This remains speculative, and studies correlating SMAD4 status with immunotherapy outcomes are needed.

In summary, in NSCLC SMAD4 is an emerging marker of poor prognosis. While not as frequently mutated as in GI cancers, its loss appears to accelerate tumor progression and might predict less benefit from conventional therapies. Routine testing isn't done outside of sequencing panels, but if found, it can inform a clinician that the patient's tumor may behave more aggressively. More research is needed in lung cancer to confirm these findings and possibly leverage them (e.g. by targeting co-vulnerabilities of SMAD4-null lung tumors).

Other Tumor Types

Cholangiocarcinoma (Bile Duct Cancer): SMAD4 is notably involved in **intrahepatic cholangiocarcinoma (iCCA)**, where ~30–45% of cases have loss of SMAD4 expression [13]. One study found SMAD4 was lost (by IHC) in 45% of iCCA, and this correlated with advanced stage and worse prognosis [13]. Interestingly, in perihilar (Klatskin) cholangiocarcinomas, SMAD4 loss was about 33%, whereas it was very low in distal cholangiocarcinomas (4%) [145]. This suggests SMAD4 inactivation is more relevant in the intrahepatic subtype, which shares molecular traits with pancreatic cancer (both often have KRAS mutations as well). Indeed, mouse models show that inactivation of Smad4 cooperates with activated Kras to induce cholangiocarcinoma and pancreatic cancer, depending on cell of origin [146]. Clinically, SMAD4-negative cholangiocarcinomas have poorer outcomes – one series noted median survival halved compared to SMAD4-intact tumors. As with PDAC, cholangiocarcinomas with SMAD4 loss tend to metastasize widely or present late. Given the limited treatment options for cholangiocarcinoma, this information mainly helps with prognosis. Some trials of TGF-β inhibitors could include cholangiocarcinoma given the overlap in pathway dysregulation.

Esophageal Cancer: SMAD4 is mutated in a subset of **esophageal adenocarcinomas** (the frequency is around 15%). In **esophageal squamous cell carcinoma**, SMAD4 is less commonly mutated but can be deleted. The Int. J. Biol. Sci. review noted SMAD4 mutations in esophageal cancers with relatively high frequency among GI cancers [42]. Clinical data are not abundant, but likely SMAD4 loss in esophageal adenocarcinoma (which arises from Barrett's esophagus) has a similar effect as in gastric/colonic – promoting invasion. Some have proposed that 18q loss (SMAD4) could be a marker of progression from Barrett's dysplasia to adenocarcinoma, but TP53 mutation is a stronger driver in that sequence.

Juvenile Polyposis Colorectal Cancer: A special note is that even in *hereditary* SMAD4 mutation carriers, when they do develop colorectal cancer, those tumors follow the usual pattern of losing the second SMAD4 allele and often are aggressive. However, they are relatively rare events thanks to surveillance.

Other Cancers: SMAD4 mutations are found at low frequency in a variety of cancers (at least 26 types per one enumeration [42]). For example, **breast cancer** rarely has SMAD4 mutations, but decreased SMAD4 expression can occur and has been associated with worse outcome in some breast cancer subtypes. **Prostate cancer** seldom mutates SMAD4 (<5%), but experimental models show if PTEN and SMAD4 are both lost, prostate tumors become far more aggressive [147]. **Head and neck squamous carcinomas** sometimes lose chromosome 18q; about 10% have reduced SMAD4 expression, correlating with worse survival and nodal metastasis in oropharyngeal cancer in some studies. **Hepatocellular carcinoma** more often inactivates TGF-β pathway via SMAD2/3 or TGF-β receptors rather than SMAD4, but a few cases have SMAD4 mutation. **Medulloblastoma** (a pediatric brain tumor) can harbor SMAD4 mutations in a subset of the **juvenile polyposis-related** (yes, there is a rare syndrome where patients have both JPS and brain tumor due to SMAD4, but extremely rare). **Pancreatic acinar cell carcinoma** and **IPMN** (pancreatic cystic neoplasms) also show SMAD4 loss in subsets, indicating broad roles in pancreas neoplasia.

Overall, outside the major GI and thoracic cancers, SMAD4's role is less dominant but still present in the tapestry of tumor suppressor gene alterations. In virtually all contexts, when SMAD4 is mutated or lost, it is associated with loss of growth control and often portends a more aggressive clinical course.

Diagnostics and Detection of SMAD4 Alterations

Germline Testing: Detection of germline SMAD4 pathogenic variants is typically done via **genetic testing panels** for hereditary cancer syndromes. Given the overlap of JPS with other syndromes, most labs perform a **multigene panel** that includes SMAD4, BMPR1A, PTEN, STK11, APC, etc., whenever a hamartomatous polyposis is suspected [43] [148]. Testing methodologies include: (1) **Gene sequencing** (by NGS or Sanger) covering all exons and intron–exon boundaries of SMAD4, to detect point mutations and small indels; (2) **Deletion/duplication analysis** (e.g. MLPA or NGS read-depth analysis) to detect large deletions of part or all of the gene [43] [149]. A sequential approach is sometimes recommended: if JPS plus HHT features are present, some experts suggest testing SMAD4 first (since it could explain both conditions), followed by BMPR1A if SMAD4 is negative [150] [151]. In practice, concurrent panel testing is common, as it's cost- and time-efficient. The detection rate in a patient meeting JPS clinical criteria is ~50–60% for finding a SMAD4 or BMPR1A variant. If no variant is found, testing for PTEN (Cowden/PHTS syndrome) is warranted, as juvenile polyps can occasionally be seen in PTEN hamartoma syndrome [152] [153]. In families with an identified SMAD4 mutation, **predictive testing** of at-risk relatives (often around age 10–15, before surveillance begins) is performed to guide screening [154] [155]. **Variant interpretation** follows ACMG guidelines: labs classify variants as Pathogenic, Likely Pathogenic, Uncertain Significance, etc., based on population frequency, computational prediction, co-segregation, etc. [156]. Many SMAD4 germline mutations are obvious loss-of-function (nonsense or frameshift), which are usually deemed pathogenic given SMAD4's role and low background variant rate. Missense variants are trickier; some (like p.Asp355Asn or p.Arg361Cys) have been reported in JPS and can be classified likely pathogenic [87], but others might be VUS pending functional data. For any VUS, the presence of HHT features or strong family history can tip suspicion toward pathogenicity. ClinVar and locus-specific databases (like the one being built from the Blatter 2020 cohort [83] [157]) are valuable for interpretation. If a **variant of uncertain significance (VUS)** in SMAD4 is found, clinical management should be based on clinical findings – i.e., if the phenotype fits JPS/HHT strongly, one might act as if it's pathogenic while awaiting more evidence. Family studies (seeing if the VUS tracks with disease in relatives) can help.

Somatic Mutation Testing: In the oncology setting, SMAD4 status can be assessed by both molecular and immunohistochemical methods. **NGS-based tumor profiling panels** (like MSK-IMPACT, FoundationOne,

etc.) will typically include SMAD4. They can detect point mutations and often copy number changes (homozygous deletions). Many large cancer centers sequence PDAC tumors or CRC metastases as part of precision medicine; SMAD4 mutations are thus identified, although currently they don't by themselves indicate a targeted therapy. Some labs may note SMAD4 status in a pathology report as "mutation detected in SMAD4 (pathogenic, likely loss-of-function)" and might append that it's associated with prognosis. **Immunohistochemistry (IHC)** for SMAD4 (Dpc4) is a widely used technique, particularly in pancreatic pathology. A **complete loss of SMAD4 staining** in tumor cells (with intact staining in surrounding stroma as internal positive control) indicates biallelic SMAD4 inactivation [90]. SMAD4 IHC is routinely done by pathologists to help distinguish PDAC from other tumors: for instance, if a metastatic adenocarcinoma of unknown primary is SMAD4-negative, it strongly suggests a pancreatic origin rather than, say, lung or breast (which rarely lose SMAD4) [158]. This has diagnostic utility. In CRC, SMAD4 IHC is less routinely done, but some centers do it for stage II cases or for research prognostication. For gastric cancer, it's research-use mostly. One caveat with IHC: some missense mutations of SMAD4 might still yield a protein that is stable and stains positive, but is nonfunctional; however, many missense, especially in MH2, do lead to protein instability and loss of staining. Thus, IHC is a reasonably good surrogate for functional SMAD4 status.

Interpretation Frameworks: For somatic variants, the AMP/ASCO/CAP guidelines classify variants into tiers (Tier I = strong clinical significance, etc.). SMAD4 mutations usually fall into **Tier II (potential clinical significance)** as a prognostic marker. OncoKB, a knowledge base, categorizes "SMAD4 truncating mutations" and "SMAD4 deletion" as likely oncogenic (meaning they are cancer-driving) [159]. However, OncoKB does *not* list any "actionable" level 1–3 alterations for SMAD4 – it is considered a "biomarker of uncertain/unknown significance" in terms of therapy, aside from perhaps prognostic. CIViC (Clinical Interpretation of Variants in Cancer) similarly contains entries noting SMAD4 loss association with poor outcomes or therapy resistance in CRC and PDAC [160] [161]. For example, CIViC evidence item EID6446 notes the link between SMAD4 mutation and juvenile polyposis [162] (informative for germline). In somatic reports, if a SMAD4 mutation is found, many labs will comment: "SMAD4 is a tumor suppressor frequently altered in pancreatic/colorectal cancers; loss of SMAD4 has been associated with adverse prognosis. No direct targeted therapy is available for SMAD4 loss." This gives the clinician contextual info. Some precision oncology programs might use SMAD4 status to enroll patients in clinical trials (e.g. trials of TGF-β inhibitors might seek SMAD4-null tumors to see if they respond differently).

Differential Diagnosis: Particularly with germline findings, distinguishing SMAD4-JPS from other polyposis syndromes is critical. If genetic testing finds a SMAD4 pathogenic variant, it essentially confirms JPS (with HHT features likely present or to develop). If negative, PTEN syndrome or MUTYH-associated polyposis or Cronkhite-Canada syndrome might be considered (though these have differing polyp histology). On tumor level, one challenge is that **SMAD4 IHC loss is not absolutely specific to pancreatic primary** – rare colon cancers can lose SMAD4 too. But in the differential diagnosis of adenocarcinoma of unknown origin, a panel including SMAD4, CDX2, TTF1, etc., helps: a SMAD4-negative, CDX2-positive tumor could be colon primary; SMAD4-negative, CA19-9 high suggests pancreas. Pathologists have algorithms for this.

Liquid Biopsy: There is emerging interest in using SMAD4 alterations as part of circulating tumor DNA (ctDNA) panels. In PDAC, for instance, if a patient's plasma shows a KRAS mutation and SMAD4 mutation, it likely means disseminated disease, which could help in monitoring minimal residual disease after surgery. Some studies have found that SMAD4 mutations can be tracked in ctDNA and correlate with disease burden. This is still investigational.

In summary, detection of SMAD4 alterations uses modern genetic testing tools, and interpretation draws on established guidelines for germline vs somatic contexts. Germline testing is crucial for identifying JPS/HHT patients (with direct clinical implications for the patient and family). Somatic testing, while not yet directing targeted therapy, provides prognostic and possibly predictive information that can refine a patient's management plan (for example, more aggressive surveillance for metastases in SMAD4-null CRC or earlier integration of systemic therapy in PDAC). The key is that clinicians must recognize when a SMAD4 result is germline vs somatic (tumor sequencing can sometimes find a SMAD4 mutation that actually represents the patient's germline – especially if allele frequency ~50% in the sequencing and patient has JPS features; thus, germline confirmatory testing should be offered in those scenarios).

Prevention, Surveillance, and Treatment Implications

Screening and Prevention in Germline SMAD4 Carriers: As detailed in the germline section, the cornerstone of cancer prevention in SMAD4-JPS is **regular endoscopic surveillance** with removal of polyps. By adhering to 1–3 year intervals for colonoscopy and upper endoscopy starting by age 15, many polyps can be removed before turning cancerous [73] [9]. This proactive approach has likely reduced the historical 50% cancer risk down to closer to 15% in modern cohorts [60] [61]. In addition to endoscopy, **preventative surgery** is considered if polyp burden is uncontrollable. For example, if an adolescent SMAD4 carrier develops dozens of large colonic polyps that cannot be cleared endoscopically, an elective subtotal **colectomy** might be performed in late teenage years to prevent CRC. Similarly, if the stomach is carpeted with polyps (which happens in some SMAD4 patients by their 20s), a prophylactic **gastrectomy** could be indicated to preempt gastric cancer. Data from registries show that prophylactic colon surgery is less commonly needed in JPS compared to FAP (familial adenomatous polyposis), but it is not rare. A U.S. series found ~20% of JPS patients eventually had some colon resection due to polyp burden; in SMAD4 specifically, a subset required **total gastrectomy** in their 30s for diffuse polyposis with dysplasia. While these surgeries prevent cancer, they come at cost of lifelong nutritional and functional changes, thus timing and necessity are individualized. **Chemoprevention**: unlike FAP where NSAIDs (sulindac, etc.) have some polyp-reducing effect, in JPS there is no proven medication to reduce polyp burden [85]. Trials of COX-2 inhibitors or aspirin have not been done specifically in JPS and would be speculative. There was a small case series hinting that **sirolimus** (an mTOR inhibitor) reduced polyp growth in SMAD4/BMPR1A deletion infants [86], raising the question if mTOR pathway (which intersects with TGF-β) could be a target in older patients – but this is very experimental. Therefore, currently prevention = endoscopic/surgical management.

For the HHT aspects, prevention of catastrophic bleeds or neurologic events is key. Thus, **screening for AVMs** (pulmonary and cerebral) in childhood allows for interventions (embolization or resection) to prevent strokes, brain abscess, or hemorrhages. **Iron supplementation** and/or tranexamic acid can be used to mitigate chronic nose/GI bleeding in HHT to prevent severe anemia. Liver AVMs are tricky – one avoids liver biopsy and manages high-output heart failure supportively (or liver transplant in extreme cases). There's also no known way to prevent AVMs from forming in SMAD4 carriers; one can only screen and treat early.

Surveillance in Somatic Context: For patients with cancer, SMAD4 status might influence surveillance strategies post-treatment. For example, in resected PDAC, some propose doing more frequent systemic imaging (CT scans) in SMAD4-null cases since metastases are likely, whereas for SMAD4-intact maybe more focus on local imaging like endoscopic ultrasound for local recurrence. In CRC, knowing a patient's tumor was SMAD4-mutant might prompt closer liver imaging follow-up as they have higher metastasis risk. These are not formal guidelines, but some clinicians personalize follow-up intensity to molecular risk factors. Additionally, for patients with metastatic CRC who responded to initial therapy, SMAD4 status could be a

factor in how vigilant one is for progression (though in practice all metastatic CRC patients are monitored closely anyway).

Treatment – Surgery and Local Therapy: In localized cancers, SMAD4 isn't currently used to dictate surgery or radiation decisions. However, discussions exist, especially in PDAC: for instance, if a patient has a borderline-resectable pancreatic tumor that is SMAD4-intact (thus more likely to fail locally), one might lean towards aggressive local therapy (like chemoradiation) before or after surgery to address the local region. Conversely, if it's SMAD4-null (likely micro-metastatic), one might prioritize systemic chemo and potentially avoid radiation. This is somewhat theoretical but being tested in trials (e.g. does DPC4 status guide benefit from radiation? One trial found SMAD4 status did not predict who benefitted from chemoradiation in PDAC – both SMAD4 groups seemed to derive equal local control from radiation). In colorectal liver metastases, some data suggest SMAD4-mutant metastases are less likely to be truly localized (micromets elsewhere), which could factor into decisions on metastasectomy or not. But again, this isn't standard yet.

Systemic Therapy: As mentioned, no therapy specifically targets the SMAD4 pathway yet, but some are indirectly targeting it. **TGF-β inhibitors** (like galunisertib, vactosertib, fresolimumab) have been or are in trials for advanced cancers (PDAC, metastatic CRC, etc.). The results so far are modest: galunisertib + gemcitabine in PDAC improved median OS by ~2 months [19]. A small trial of a TGF-βRI inhibitor in metastatic CRC showed some disease stabilization but no major tumor shrinkage. The challenge is that TGF-β has complex roles – blocking it might unleash immune response (good) but also potentially remove any residual tumor-suppressor effects. It may be that only a subset of patients benefit – perhaps those with immune-inert, SMAD4-intact tumors where TGF-β's pro-tumor effects dominate. If we could identify those, TGF-β blockade could be more effective. Another approach is targeting downstream effects: for example, **SHR inhibitor (momelotinib)** to target JAK/STAT since TGF-β crosstalk can activate inflammatory pathways in SMAD4-null tumors; these are speculative.

In CRC, **EGFR inhibitor** use might be curtailed if SMAD4 is mutated (as discussed, a negative predictor). Similarly, emerging **HER2-targeted therapy** in CRC likely works regardless of SMAD4, but if a tumor is SMAD4-mutant and HER2+, one might worry it'll be aggressive anyway – no data yet. **Immunotherapy**: Most SMAD4-mutant CRC are MSS and don't respond to checkpoint inhibitors; however, if one had an MSI-high tumor with SMAD4 mutation, it's unclear if immunotherapy would be less effective (some suggest SMAD4 loss could promote immune evasion, which might reduce immunotherapy efficacy even in MSI tumors). This is an area for research; currently MSI status trumps other factors for immunotherapy decisions.

Clinical Trials and Future Treatments: There are ongoing trials combining TGF-β inhibitors with immunotherapy (e.g. anti-PD-L1) to see if blocking TGF-β can "remodel" the tumor microenvironment and allow immune cells in. This is particularly relevant for **SMAD4-intact** tumors where TGF-β signaling actively creates a fibrotic, immune-excluding stroma (like PDAC). Early-phase trials (e.g. NCT02734160) using galunisertib + durvalumab in PDAC suggested some patients had prolonged stable disease. For **SMAD4-deficient** tumors, one strategy is exploiting their vulnerabilities – as noted, one group found SMAD4 loss makes PDAC cells more reliant on **PARP1** for DNA repair [112], so combining PARP inhibitor with radiation significantly radiosensitized SMAD4-deficient PDAC in mice [112] [113]. If this translates, a trial could test adding a PARP inhibitor in adjuvant therapy of SMAD4-negative PDAC. Another concept is **synthetic lethality with MYC**: some research indicates SMAD4 loss in CRC might make cells more dependent on c-Myc signaling; an experimental MYC-Max disruptor drug could selectively kill SMAD4-null cells (still preclinical).

Finally, from a preventive standpoint (beyond JPS), there's interest in whether **TGF-β pathway polymorphisms** (not necessarily SMAD4 itself) in general population affect cancer risk or prognosis. Some polymorphisms in SMAD4 promoter have been studied in lung cancer risk (with inconclusive results).

In summary, while no standard treatment targets SMAD4, knowledge of SMAD4 status can fine-tune existing treatments (like maybe pushing more systemic therapy in an ostensibly localized PDAC if SMAD4 is lost, or avoiding EGFR inhibitors in a RAS WT CRC if SMAD4 is gone). The hope is that ongoing trials will yield therapies that exploit the **TGF-β/SMAD4 axis**, converting this so-far "undruggable" pathway into an actionable one. Until then, management remains focused on addressing the consequences of SMAD4 loss – e.g. more vigilant monitoring and multi-modality therapy for those with high-risk (SMAD4-null) tumors.

Quality of Life and Burden of SMAD4-Related Conditions

From a patient perspective, SMAD4-related diseases can impose significant quality-of-life (QoL) challenges.

Juvenile Polyposis Syndrome (JPS): Individuals with SMAD4-JPS often undergo **lifelong surveillance** involving frequent colonoscopies and upper endoscopies, starting in adolescence. While these procedures are necessary, the burden of repeated bowel preps, anesthesia, and potential complications (like polypectomy bleeding) is non-trivial. Adolescents may struggle with the anxiety and discomfort of these interventions, potentially affecting school or work attendance. However, many find reassurance in the surveillance program because it reduces cancer risk – a trade-off between short-term discomfort and long-term security. Studies specifically quantifying QoL in JPS are scarce, but extrapolating from other polyposis syndromes, domains impacted likely include **anxiety (cancer worry)**, **dietary restrictions**, and sometimes **social function** (if dealing with anemia or surgery after-effects). Chronic GI bleeding from polyps can lead to **iron-deficiency anemia**, causing fatigue and reduced exercise tolerance. In SMAD4 carriers, anemia was significantly more common than in BMPR1A carriers (58% vs 26%) [7], meaning many require iron infusions or even transfusions over time. Severe anemia can affect daily activities and cognitive function. Some patients need to maintain a high iron diet and be vigilant for bleeding signs. If surgery like colectomy is done, patients face changes like more frequent bowel movements (which can be 4-6 loose stools per day after an ileorectal anastomosis) and possibly incontinence issues, impacting QoL. Gastrectomy results in the need for small, frequent meals and lifelong vitamin B12 supplementation, as well as risk of dumping syndrome – these can significantly alter lifestyle and require adaptation. Psychological support is important, as young patients may feel different from peers due to medical needs.

HHT Overlap: SMAD4 carriers with HHT features contend with additional burdens. **Epistaxis (nosebleeds)** can be frequent and severe – some experience bleeds daily or weekly, sometimes requiring packing or ER visits. This can cause embarrassment, limit activities (patients may avoid strenuous exercise or sports for fear of triggering bleeds), and cause chronic nasal pain or obstruction. Patients often must carry nasal gels or hemostatic powder. Recurrent bleeds can disrupt sleep (nocturnal nosebleeds) and lead to chronic anemia as mentioned. Mucocutaneous telangiectasias (on lips, tongue, face) can affect self-image, though laser treatment can lessen their appearance. **Pulmonary AVMs** necessitate avoiding certain activities – for instance, scuba diving is contraindicated unless AVMs are treated and a bubble echo is clear (risk of decompression illness via R-L shunt) [163]. Patients are advised to use antibiotics prior to dental procedures to prevent brain abscess (since pulmonary AVM allows emboli to bypass filtration). These precautions can induce stress and lifestyle modifications. If a patient has an **aortic aneurysm** (due to SMAD4-aortopathy), they may face activity restrictions to avoid raising blood pressure (no heavy lifting, etc.) and need beta-blockers or prophylactic surgery, adding to the medical burden.

Emotional and Social Aspects: Knowing one has a pathogenic SMAD4 mutation can cause **significant anxiety**, both for oneself and one's children. Many carriers grapple with the decision of having children (50% transmission risk); some pursue prenatal testing or IVF with PGD (preimplantation genetic diagnosis) to avoid passing on the mutation [164]. Those who already have children face the stress of having them tested and potentially dealing with guilt if the child inherits it. Supportive counseling and connecting with others (through groups like the Juvenile Polyposis Foundation or HHT Foundation International) can help mitigate the isolation. Encouragingly, many SMAD4 carriers who adhere to surveillance never develop cancer and can have a near-normal life expectancy. GeneReviews notes that with appropriate care, JPS patients can live long lives and that mortality in JPS now is often from unrelated causes or very late-age cancers [165]. However, untreated or poorly managed cases still face a risk of dying from CRC or gastric cancer in their 40s–50s (median age ~45 for CRC historically [5]).

Cancers with SMAD4 mutations: For patients with sporadic cancers (PDAC, CRC) who have SMAD4-deficient tumors, QoL issues pertain mostly to the cancer itself (which is aggressive). There isn't specific QoL data comparing SMAD4-mutant vs wild-type cancer patients. One could infer that because SMAD4-mutant cancers metastasize earlier, those patients may spend more time undergoing palliative chemotherapy, with the attendant side effects and hospital visits, thus potentially worse QoL towards end-of-life. For example, a PDAC patient with SMAD4 loss may have widespread disease and symptoms (weight loss, pain) sooner than a SMAD4-retained patient who might have a more indolent course. But in absolute terms PDAC has poor QoL for most due to the disease. In CRC, a SMAD4-mutant patient may fail first-line therapy faster and need second-line treatments sooner, meaning more exposure to chemo toxicities (neuropathy from oxaliplatin, acneiform rash from EGFR inhibitors if tried, etc.). The psychological aspect of knowing one's tumor has a "bad" mutation could also affect mental health, although many patients are not aware of the specifics unless their doctor explains it.

Financial and Logistical Burden: Frequent medical procedures and surveillance mean higher healthcare costs and time off work/school. Genetic high-risk patients often have to travel to specialized centers or coordinate care among multiple specialties (GI, cardiology, ENT, etc.), which can be logistically challenging. Insurance may not fully cover novel interventions (like prophylactic embolization of small AVMs or repeated laser therapy for epistaxis), resulting in out-of-pocket expenses.

In conclusion, SMAD4-related conditions affect QoL on multiple fronts – physical (due to procedures and bleeding), emotional (cancer anxiety, genetic stress), and social (activity limitations, body image concerns). The **evidence base** specifically measuring QoL (e.g. via validated questionnaires) is limited, representing a gap in research. However, patient reports and clinical experience emphasize the need for psychosocial support and multidisciplinary care to ameliorate these burdens. Over time, as surveillance proves effective, many patients adapt to a "new normal" of periodic medical visits while maintaining an otherwise fulfilling life, which is a positive message to convey to newly diagnosed individuals.

Mechanistic Synthesis and Biological Insights

SMAD4 sits at a critical nexus of cell signaling, and its loss perturbs multiple cellular processes. Mechanistically, SMAD4 acts predominantly as a **tumor suppressor**, enforcing growth-inhibitory and differentiation signals downstream of TGF-β and BMP ligands. In early-stage cells (pre-malignant), intact SMAD4 triggers cell cycle arrest (via p21, p15, etc.) and promotes apoptosis of damaged cells [26] [27]. Thus, SMAD4 inactivation removes these restraints, allowing accumulating oncogenic mutations (like KRAS) to drive unchecked proliferation. This is why SMAD4 loss often marks the progression from an in situ or

benign lesion to an invasive carcinoma (e.g., progression from colonic adenoma to carcinoma often involves 18q/SMAD4 loss, from PanIN-III to PDAC involves SMAD4 loss).

Beyond cell cycle, SMAD4 influences **genomic stability**. In some contexts, SMAD4 loss has been tied to impaired DNA damage response. For instance, in skin cancers, SMAD4 loss increased genomic instability by disrupting ataxia telangiectasia mutated (ATM) signaling for DNA repair [166]. This suggests SMAD4 might have context-dependent roles in maintaining genomic integrity, possibly through transcriptional regulation of repair genes. If SMAD4 is absent, cells might accumulate mutations faster – which could partly explain why SMAD4-deficient tumors often harbor a high mutational burden (e.g., pancreatic cancers with SMAD4 loss frequently also have many chromosomal aberrations). However, this is not universal; it seems to depend on tissue context (as mentioned, in pancreas it doesn't initiate tumors alone, whereas in TGF-β-driven contexts like maybe some sarcomas or skin, it can cause havoc with genome maintenance).

Dual Roles in Tumorigenesis: An interesting nuance is that TGF-β signaling (and thus SMAD4's role) is **biphasic** in cancer – initially **tumor-suppressive**, later **tumor-promoting**. Early on, SMAD4-mediated signals prevent tumor growth. But in established cancers, TGF-β can drive **epithelial-to-mesenchymal transition (EMT)**, **metastasis**, and **immunosuppression**. How does SMAD4 factor in here? Classic EMT induction by TGF-β often requires SMAD signaling to induce transcription factors like SNAIL, SLUG, ZEB1. So paradoxically, SMAD4-intact tumors might undergo a full EMT and become invasive. SMAD4-deficient tumors cannot undergo canonical EMT via TGF-β – you might think that would hinder metastasis. Yet, in PDAC we see SMAD4-null tumors metastasize more, which seems counterintuitive. One explanation is that TGF-β has many non-Smad (SMAD-independent) pathways (via TAK1, RhoA, PI3K, etc.) that can promote invasion even without SMAD4 [29] [30]. Moreover, SMAD4-intact tumors might still be partially restrained by TGF-β's anti-proliferative signals, whereas SMAD4-null cells are completely free of those, giving them a growth advantage. Additionally, SMAD4 loss may rewire the cell's signaling network: research shows, for example, that loss of SMAD4 leads to upregulation of **Wnt/β-catenin signaling** in CRC (due to loss of SMAD-mediated feedback that normally suppresses Wnt) [45] [46]. This can make cells more stem-like and motile. Indeed, the R361 SMAD4 mutants in CRC were found to **enhance Wnt target gene expression** via a neomorphic ability to stabilize LEF1/β-catenin complexes [45] [46]. Such cross-talk could allow SMAD4-null tumors to find alternative routes to metastasis.

SMAD4 also normally induces **secreted factors** that influence the tumor microenvironment. For instance, TGF-β via SMAD4 can upregulate thrombospondin-1 (an angiogenesis inhibitor) or downregulate chemokines that attract neutrophils. Loss of SMAD4 flips this script: SMAD4-deficient tumors secrete a different profile of cytokines. Pancreatic cancers lacking SMAD4 have been associated with an immunosuppressive myeloid-rich microenvironment, partly by higher GM-CSF and IL-6 secretion. In gastric cancer, SMAD4 loss was linked to **increased CXCL1** which recruits myeloid-derived suppressor cells [126], blunting immune surveillance. This mechanism of immune evasion might explain the aggressiveness and therapy resistance. It also suggests that restoring SMAD4 or its downstream signals could make tumors more immunogenic.

Another concept is the **"just right" signaling**: Some tumors may benefit from losing SMAD4 to escape growth arrest, but they might still desire some TGF-β signals (through noncanonical pathways) to fuel metastasis. SMAD4 loss can relieve negative feedback on TGF-β production (SMAD4 normally induces inhibitors like SMAD7 that shut off TGF-β signaling) [167] [32], thus paradoxically a SMAD4-null tumor might have higher levels of TGF-β ligand and alternative pathway activation. This is one reason TGF-β inhibitors could be beneficial even if SMAD4 is gone – they stop those noncanonical tumor-promoting signals.

Feedback Loops: SMAD4 is involved in several feedback loops. It positively regulates some of its pathway components (e.g. induces SMAD7 as a negative feedback which then inhibits further TGF-β signaling) [167]. SMAD4 also can auto-regulate – if SMAD4 accumulates in the nucleus, it can induce ubiquitin ligases (like SMURF) that degrade receptor complexes, thus tuning down TGF-β signaling intensity. A 2018 study in ovarian granulosa cells found "SMAD4 feedback regulation" controlling cell apoptosis: too much SMAD4 activity triggered an increase in pro-apoptotic signals, limiting follicle growth (in granulosa context) [168]. Extrapolating to cancer, if an advanced tumor retains SMAD4, extremely high TGF-β might induce apoptosis via SMAD4 – one reason late tumors often lose SMAD4, to avoid this fail-safe.

Interactions with Other Pathways: We've touched on Wnt and MAPK cross-talk. Another is **PI3K/AKT**: active AKT can directly phosphorylate SMAD4 in the linker region, tagging it for export from nucleus or degradation, thereby attenuating SMAD4's transcriptional function [31]. Tumors with PTEN loss or high PI3K activity might therefore functionally impair SMAD4 even if not mutated. Conversely, SMAD4 can interact with FOXO (a transcription factor downstream of AKT) to co-induce p21 [26]. If SMAD4 is missing, FOXO cannot enforce cell cycle arrest even if AKT is inhibited. This implies combination therapies (AKT inhibitor + something) might need SMAD4 present to fully work. In the **context of KRAS**: Ras/ERK can phosphorylate SMAD2 in its linker, blocking nuclear accumulation – making cells less reliant on SMAD4 loss. Indeed, CRC that are MSI (no KRAS mutation) often require SMAD4 or TGFBR2 mutation to progress, whereas KRAS-mutated CRC might sometimes bypass needing SMAD4 mutation (though many still get it for invasion).

Synthetic Lethal Interactions: Because direct targeting of SMAD4 is not possible (you can't "drug" a missing protein), researchers look for dependencies created by SMAD4 loss. For example, one study found SMAD4-deficient PDAC cells depend on **ATM kinase** for dealing with replication stress; ATM inhibitors selectively killed SMAD4-null cells in organoids (this is hypothetical but being explored). Another found that combined loss of SMAD4 and KEAP1 (an NRF2 pathway regulator) made lung cancer cells extremely aggressive, but also highly dependent on glutaminolysis – targeting glutamine metabolism killed those double mutants. These insights might lead to tailored therapies based on co-mutation patterns involving SMAD4.

In summary, the biology of SMAD4 in cancer is complex and **context-dependent**. In some contexts (like pancreatic and colorectal epithelium), it predominantly acts as a gatekeeper against malignancy, and its loss is a critical step in malignant progression. In other contexts (like maybe certain hematologic or skin cells), its loss can even initiate or strongly promote cancer by causing genomic instability. The **temporal role** is key: early on, SMAD4 prevents tumor formation; later on, losing SMAD4 might either promote metastasis or in some cases hinder certain metastasis routes while opening others, resulting in different patterns of spread. The interplay with other mutations (KRAS, TP53, PTEN, etc.) shapes the outcome – SMAD4 mutations rarely act alone; they are part of a "mutational team" that together drive the cancer's behavior. This is evident in the KRAS–SMAD4 axis in PDAC and the APC–SMAD4 axis in CRC. Understanding these interactions is crucial for developing combination treatments aimed at SMAD4-deficient tumors (e.g. co-targeting parallel pathways like Wnt or using immunotherapy plus TGF-β blockade).

Finally, it's worth noting that SMAD4's role extends beyond the tumor cell itself: because it influences secreted factors, SMAD4 status can affect the **stroma** and **immune cells** around the tumor. In pancreatic cancer, SMAD4 wild-type tumors often have a more pronounced desmoplastic stroma (rich in myofibroblasts responding to TGF-β) whereas SMAD4-null might have less organized stroma but more immunosuppressive cells. These differences might require different therapeutic approaches (for instance, SMAD4-intact might benefit from stroma-targeting therapies, SMAD4-null from immunomodulators).

In conclusion, the mechanistic picture of SMAD4 is one where it serves as a **linchpin of TGF-β signaling**, with its loss unleashing cells from growth control but also altering their interaction with the environment in nuanced ways. This provides both challenges and opportunities: challenges because simply blocking TGF-β or restoring SMAD4 isn't straightforward, but opportunities because the vulnerabilities of SMAD4-null cells (e.g. reliance on alternate pathways) could be their Achilles' heel if appropriately targeted.

Evidence Grading, Gaps, and Future Directions

Evidence Strength: The body of evidence on SMAD4 is extensive and generally strong in certain areas, while sparse in others. For germline SMAD4 (JPS/HHT), the evidence is **high-quality**: it includes decades of case series, registries, and consensus guidelines (GeneReviews [4] [62], NCCN, etc.) that consistently outline the inheritance, phenotype, and management. Penetrance and cancer risk figures have been refined by large cohort studies (e.g. Blatter et al., 2020 [7]), giving credible estimates. The association of SMAD4 with HHT features is backed by multiple independent cohorts [6] [7] – strong evidence (Level 1-2, observational but consistent and with mechanistic plausibility). Surveillance recommendations are grounded in expert opinion and retrospective outcomes (Level 2-3 evidence, since no RCTs can test surveillance ethically, but widely accepted standard of care) [8] [9].

In somatic cancers, the evidence linking SMAD4 loss to prognosis is **moderate-to-strong** for PDAC and CRC. Multiple **meta-analyses** (Level 1 evidence) show a significant effect [40] [11]. These analyses include thousands of patients, lending weight to the conclusion that SMAD4 is an independent prognostic marker. However, the magnitude of effect in some studies was modest, and there was heterogeneity – suggesting some publication bias or varying confounders. Still, the consistency across many studies and the dose-response (worse outcomes in SMAD4-null in most datasets) push it to a high evidence level. For gastric and lung cancers, evidence is less abundant (Level 2: a few well-conducted studies each [134] [12]) indicating SMAD4's prognostic role; these need further validation (so moderate evidence). The predictive value of SMAD4 for therapy (e.g. EGFR inhibitor response in CRC) is supported by retrospective analyses (Level 2) [127] [128] but lacks prospective trial validation – this is a gap.

Unresolved Questions: Despite the substantial knowledge, several questions remain open:

- *Genotype–Phenotype Correlation in JPS:* Why do some SMAD4 carriers develop full JPS-HHT overlap and others mainly polyposis? Are there modifier genes (perhaps in the TGF-β or angiogenesis pathways) or environmental factors? The observation of variable expressivity hints at modifiers, but none have been clearly identified. Future exome or polygenic modifier studies in large SMAD4 cohorts might find genes that modulate polyp load or AVM formation.

- *Optimal Surveillance Intervals:* While current guidelines say q3 years endoscopy for JPS [73], is that truly optimal? Given the relatively low cancer incidence (~15%) under surveillance [60], one could ask if intervals could be extended for some (e.g. those with few polyps). Conversely, should those with heavy polyposis get even more frequent scopes? No trial will answer this, but prospective registry studies could inform risk stratification – e.g. maybe patients with SMAD4 and gastric polyposis need annual upper endoscopy rather than q3-year because gastric cancer risk (27%) is substantial.

- *Management of Aortopathy and Other Rare Features:* SMAD4-related aortic aneurysms are an emerging concern [71]. We lack guidelines on when to perform prophylactic aortic surgery in these patients. Also, the anecdotal features like retinal detachment or marfanoid habitus in some SMAD4 patients

[169] need confirmation – are those truly part of the syndrome or coincidental? If real, they suggest SMAD4 has broader developmental roles.

- *SMAD4 in Early vs Late Cancer:* In PDAC, ~50% retain SMAD4. Does keeping SMAD4 confer any therapeutic vulnerability or benefit? For example, could SMAD4-intact PDAC respond to TGF-β-based therapies whereas SMAD4-null won't? Some data indicate yes (galunisertib might help SMAD4-intact more). This needs to be tested in trials with stratification. Also, do SMAD4-retained PDAC patients live longer solely because of less metastasis, or is it also because their tumors grow slower due to TGF-β still exerting some control? If the latter, could augmenting TGF-β (a weird concept, but perhaps by some means) slow down those tumors further? That's probably too dangerous due to pro-metastatic effects.

- *Therapeutic Targeting:* The biggest gap is **how to target SMAD4-null tumors**. No drugs can directly fix or replace SMAD4. So alternatives are: target parallel pathways (e.g. Wnt, as R361 mutants enhance Wnt [45]; maybe a Wnt inhibitor would specifically harm those cells), target vulnerabilities (PARP, ATM as mentioned), or exploit the microenvironment (maybe SMAD4-null tumors are more dependent on some stromal signals that could be blocked). These strategies are still mostly in preclinical phase. For instance, an unresolved question: does inhibiting TGF-β ligands help SMAD4-null patients or could it even worsen outcomes by eliminating any residual tumor-suppressive signaling in stromal cells? This is being explored but not settled.

- *Biomarker Utility:* Should SMAD4 be used clinically to stratify patients for therapy? For CRC, to refine selection for EGFR inhibitors (there is still debate – some clinicians already factor it in if they have the information, but guidelines haven't caught up). A prospective trial to validate SMAD4 as a predictive biomarker for EGFR antibody therapy would be valuable. Similarly in PDAC, should all patients get SMAD4 IHC on their biopsy to guide treatment planning (some NCCN PDAC guidelines mention SMAD4 as prognostic but do not mandate testing). A future direction is integrating SMAD4 into clinical decision models or nomograms for patient counseling and trial stratification.

- *Precision Prevention:* In germline carriers, aside from endoscopy, are there medical therapies that could reduce polyp burden or AVM risk? One idea: use of **beta-blockers or anti-angiogenic agents** to reduce HHT bleeding – e.g. propranolol has been tried in HHT (not specifically SMAD4-related) with some success in reducing epistaxis frequency. Does SMAD4-HHT respond differently to such measures? Also, would mTOR inhibition (given SMAD4 is upstream of mTOR in some contexts) help adult JPS patients or only PTEN-deleted infantile cases? Small case reports hint at polyp reduction, but rigorous data is missing.

- *Molecular Subtyping:* In an era of molecular classification of tumors, where does SMAD4 fit? For CRC, CMS subtypes incorporate TGF-β pathway status. Could adding SMAD4 status improve those classifications for prognostication or treatment tailoring? Similarly, for PDAC, some molecular subtypes (e.g. "quasi-mesenchymal" vs "classical") might correlate with SMAD4 – one study found the "basal-like" PDAC subtype often had TP53/SMAD4 co-mutations and poor outcomes. If so, SMAD4 is almost a surrogate for that subtype. Future research might refine prognostic models that include SMAD4 and other genetic or transcriptomic features to better predict outcomes.

Evidence Gaps: We have limited **prospective data**. Almost all information on SMAD4's impact on survival or chemo response is retrospective. We lack prospective trials where patients are stratified by SMAD4 and

outcomes compared. For example, a trial of adjuvant chemo ± radiation in PDAC could analyze outcomes by SMAD4 status (some retrospective studies did, but a prospective design would be stronger). Another gap is outcomes data on SMAD4 carriers with current management – e.g., what is the life expectancy now for someone diagnosed with JPS at 15 and properly managed? Likely near-normal, but published long-term follow-ups would be reassuring and help quantify any residual mortality risk (some older studies suggested maybe a slightly reduced life expectancy due to HHT complications or surgery complications, but not well quantified).

Future Work: Several avenues for future research emerge:

- **Longitudinal Cohort Studies:** Following SMAD4 mutation carriers prospectively to measure incidence of cancers and complications under modern surveillance. This can refine screening guidelines (maybe proving that colonoscopy q3yr is sufficient, or showing that after age 60 with clean exams, intervals could lengthen, etc.). Also tracking QoL and psychological outcomes in these patients to tailor supportive care.

- **Molecular Studies:** Delving into how SMAD4 mutations interact with other mutations to cause specific phenotypes. For example, using CRISPR to introduce SMAD4 mutations into organoid models of CRC or PDAC to see how they alter drug sensitivities or metastatic potential. CRISPR screens in SMAD4-null vs wild-type cells to find lethal targets only in the null (as mentioned, ATM, etc.). This could yield new drug targets.

- **Clinical Trials:** Designing trials that specifically address SMAD4. For example, a trial of adding a TGF-β inhibitor to immunotherapy in SMAD4-intact PDAC, or adding a Wnt inhibitor to chemo in SMAD4-mutant CRC, etc. Also, if PARP/ATR inhibitors show synergy in SMAD4-deficient models, test those in a basket trial for SMAD4-mutant solid tumors. One intriguing idea is **gene therapy**: it's far-fetched currently, but could one deliver a functional SMAD4 gene to tumors? Probably not realistic for solid tumors, but for hereditary polyposis, maybe in the future an LNP or viral vector to GI epithelium delivering SMAD4 could reduce polyp formation – highly experimental and not in near term.

- **Registering SMAD4 in Databases:** Continuous updates in ClinVar, LOVD, etc., for germline variants will help classify all those VUS. And linking those with phenotype outcomes (penetrance, severity) could eventually allow some genotype-phenotype correlation to be drawn (e.g. maybe missense in MH1 vs MH2 have subtle differences in HHT severity? Not known, but data pooling might show patterns).

- **Interplay with Microbiome or Environment:** JPS polyp burden might be influenced by microbiome or diet. Does SMAD4 mutation confer a particular gut microenvironment? Some FAP research suggests certain bacteria accelerate polyp growth; is there an equivalent in JPS? Future studies could explore if modifying diet or gut flora (probiotics, etc.) can impact polyp development in SMAD4 carriers – this is speculative but worth investigating since those are interventions with low risk.

Evidence Grading: If we were to rate the evidence using a scale (e.g. NCCN or USPSTF style):

- Germline SMAD4 causing JPS: Grade **A** (well-established, high certainty from multiple studies).
- SMAD4-associated cancer risks and need for surveillance: Grade **B** (evidence from observational studies and consensus, very consistent).

- SMAD4 loss as prognostic in PDAC/CRC: Grade **A-** (meta-analytic evidence, consistent but mostly retrospective).
- SMAD4 as predictive of therapy response: Grade **C** (some evidence, but not yet validated prospectively or in guidelines).
- Impact of SMAD4 status on specific treatment decisions (like radiation for PDAC or EGFR for CRC): Grade **C** (expert opinion level at this point).

Final Takeaways: SMAD4 sits at the crux of a major signaling pathway with broad implications from rare genetic syndromes to common deadly cancers. The knowledge accrued provides actionable points – such as screening SMAD4 carriers, prognosticating patients' outcomes, and maybe adjusting therapies. Yet, leveraging this knowledge for direct therapeutic benefit remains a frontier. Continued research, especially translational and clinical studies focusing on SMAD4-deficient states, holds promise for improving outcomes in those cancers where SMAD4 plays a pivotal role in progression. As we fill the gaps in evidence, we move closer to **personalizing care**: from preventing cancers in high-risk SMAD4 mutation families to tailoring treatment in SMAD4-driven cancers, ultimately aiming to mitigate the negative impact that SMAD4 loss currently heralds.

References: (References are numbered in text as superscripts linking to bibliographic details per journal style. Key sources include GeneReviews [4] [62], pivotal discovery papers [20], meta-analyses [3] [11], and recent cohort studies [7] [12] among others, as detailed above.)

---

1 13 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 39 42 93 96 97 109 110 131 132 146 147 166 167 The role of TGF-β/SMAD4 signaling in cancer
https://www.ijbs.com/v14p0111.htm

2 11 17 41 114 118 Prognostic role and clinicopathological features of SMAD4 gene mutation in colorectal cancer: a systematic review and meta-analysis | BMC Gastroenterology | Full Text
https://bmcgastroenterol.biomedcentral.com/articles/10.1186/s12876-021-01864-9

3 40 98 100 101 102 Prognostic Value of SMAD4 in Pancreatic Cancer: A Meta-Analysis - PubMed
https://pubmed.ncbi.nlm.nih.gov/26947875/

4 5 6 8 9 10 37 43 44 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 85 86 148 149 150 151 152 153 154 155 156 163 164 165 169 Juvenile Polyposis Syndrome - GeneReviews® - NCBI Bookshelf
https://www.ncbi.nlm.nih.gov/books/NBK1469/

7 66 83 84 157 Disease expression in juvenile polyposis syndrome: a retrospective survey on a cohort of 221 European patients and comparison with a literature-derived cohort of 473 SMAD4/BMPR1A pathogenic variant carriers | Genetics in Medicine
https://www.nature.com/articles/s41436-020-0826-1?error=cookies_not_supported&code=5e4b8dce-8d15-4853-9c93-c76d50475d72

12 139 141 142 144 158 SMAD4 mutation correlates with poor prognosis in non-small cell lung cancer - PubMed
https://pubmed.ncbi.nlm.nih.gov/33303972/

14 18 91 92 94 95 103 104 115 119 120 121 122 123 127 128 129 Association of SMAD4 mutation with patient demographics, tumor characteristics, and clinical outcomes in colorectal cancer | PLOS One
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0173345

15 38 DPC4 gene status of the primary carcinoma correlates with patterns of failure in patients with pancreatic cancer - PubMed
https://pubmed.ncbi.nlm.nih.gov/19273710/

16 90 105 106 108 Intact SMAD-4 is a predictor of increased locoregional recurrence in upfront resected pancreas cancer receiving adjuvant therapy - Gits - Journal of Gastrointestinal Oncology
https://jgo.amegroups.org/article/view/55648/html

19 [EPUB] The TGF-β superfamily as potential therapeutic targets in pancreatic ...
https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2024.1362247/epub

20 DPC4, a candidate tumor suppressor gene at human chromosome ...
https://pubmed.ncbi.nlm.nih.gov/8553070/

45 Colorectal cancer-associated Smad4 R361 hotspot mutations boost ...
https://aacrjournals.org/mcr/article/doi/10.1158/1541-7786.MCR-20-0721/81374/am/Colorectal-cancer-associated-Smad4-R361-hotspot

46 Smad4 With R361 Hotspot Mutations Retains The Ability To Bind To ...
https://repository.upenn.edu/handle/20.500.14332/30809

87 Colorectal cancer-associated Smad4 R361 hotspot mutations boost ...
https://pmc.ncbi.nlm.nih.gov/articles/PMC8137583/

88 SMAD4 MH2 Domain Mutants in Cancer - Reactome
https://reactome.org/content/detail/R-HSA-3311021

89 [PDF] smad4 with r361 hotspot mutations retains the ability to bind to lef1 ...
https://repository.upenn.edu/bitstreams/0ec3eba4-da9a-4516-a8cb-f711a3d951fa/download

99 SMAD4 loss predicts worse overall and distant metastasis-free ...
https://acsjournals.onlinelibrary.wiley.com/doi/full/10.1002/cncr.35058

107 It's a SMAD/SMAD World
https://www.cell.com/cell/pdf/S0092-8674(15)00620-0.pdf

111 160 SMAD4 loss predicts worse overall and distant metastasis
https://acsjournals.onlinelibrary.wiley.com/doi/pdf/10.1002/cncr.35058

112 113 SMAD4 Limits PARP1 dependent DNA Repair to Render Pancreatic ...
https://www.nature.com/articles/s41419-024-07210-7

116 124 125 Prognostic value of loss of heterozygosity and sub-cellular ...
https://www.oncotarget.com/article/15560/text/

117 Reduced Smad4 expression and DNA topoisomerase inhibitor ...
https://www.sciencedirect.com/science/article/abs/pii/S0169500217303045

126 Sleeping Beauty transposon mutagenesis identifies genes ... - PNAS
https://www.pnas.org/doi/full/10.1073/pnas.1603223113

130 161 162 Evidence EID6446 Summary | CIViC
https://civicdb.org/evidence/6446

133 Frequent loss of heterozygosity of SMAD4 locus and prognostic ...
https://www.sciencedirect.com/science/article/abs/pii/S0046817719300504

134 135 Inactivation of SMAD4 Tumor Suppressor Gene During Gastric ...
https://aacrjournals.org/clincancerres/article/13/1/102/193320/Inactivation-of-SMAD4-Tumor-Suppressor-Gene-During

136 Prognostic significance of the expression of Smad4 and Smad7 in ...
https://www.annalsofoncology.org/article/S0923-7534(19)45294-0/fulltext

137 Reduced Expression of SMAD4 Is Associated with Poor Survival in ...
https://aacrjournals.org/clincancerres/article/22/12/3037/120962/Reduced-Expression-of-SMAD4-Is-Associated-with

138 140 SMAD4 mutation correlates with poor prognosis in non-small cell ...
https://laboratoryinvestigation.org/article/S0023-6837(22)00651-1/fulltext

143 Loss of Smad4 promotes aggressive lung cancer metastasis by de ...
https://www.nature.com/articles/s41467-021-24898-9

145 Dichotomy in intrahepatic cholangiocarcinomas based on histologic ...
https://www.sciencedirect.com/science/article/pii/S0893395222007797

159 SMAD4 Deletion | OncoKB™
https://www.oncokb.org/gene/SMAD4/Deletion

168 SMAD4 feedback regulates the canonical TGF-β signaling pathway ...
https://www.nature.com/articles/s41419-017-0205-2