


titanic_ball_eta3e-5: exact gradient, LASSO anchor (first 600 iterations)



ALL RUNS STOPPED AT ITERATION 600 -- a truncation of the notebook's longer runs (identical W_0 and Gaussian stream), not a separate experiment. TITANIC -- ball $\|w\|^2 \leq 2$ (radius $\sqrt{2}$). $d = 10$ (intercept w_0 + nine features), no mini-batching: the EXACT gradient is used at every step. $\sigma = 10$, $\lambda = 7.12$, $\delta = 0.01082$, so a in $[0.50, 1]$. Block strengths $s = (5, 5, 5)$ on $w_1..w_9$ only. Step $\eta = 3e-05$. $R = 150$ coupled replicates (same W_0 , same Gaussian stream) on one fixed stratified 80/20 split ($n_{\text{train}} = 712$, $n_{\text{test}} = 179$). Paired difference AT ITERATION 600, non-reversible minus reversible: training $+0.0107$ ($t = +5.72$), test $+0.0128$ ($t = +4.82$). Bands: mean $\pm$ 1 SD ($\text{ddof} = 1$) across replicates -- repeat-run variability at a fixed split, not a confidence or credible interval. The ACCURACY AXIS IS $[0.4, 1]$, so the lower band edge runs off the bottom of the panel during the first few hundred iterations: about a quarter of replicates start below 0.4 and the lowest state visited is near 0.27 (see figures/axis_explainer.png).