



magic_lp: per-block strengths at $\eta = 1e-05$ (mix)



MAGIC -- smoothed ℓ_p $g(w) = \sum_i (w_i^2 + \epsilon^2)^{p/2} \leq \Lambda$, $p = 2.4$, $\epsilon = 0.2$, $\Lambda = 4$. $d = 10$ (intercept w_0 + 9 features). $U = \sum \text{softplus}(-y x \cdot w + w_0^2 / (2 \sigma^2)) + \lambda \sum_{j \geq 1} |w_j|$; U_0 replaces $|w_j|$ by $\sqrt{w_j^2 + \delta^2}$; $a = \exp(U - U_0)$. EXACT gradient. $\sigma = 10$, $\lambda_{\text{lasso}} = 152 = 0.01 n_{\text{train}}$, $\delta = 0.000506$ (a in $[0.50, 1]$). Step $\eta = 1e-05$. Per-block strengths $s = (s_1, s_2, s_3)$ on $l_1 = \text{PC1}.. \text{PC3}$, $l_2 = \text{PC4}.. \text{PC6}$, $l_3 = \text{PC7}.. \text{PC9}$. Isotropic vs gradient-balanced vs reversed-balanced vs hand-picked. $R = 100$ independent replicates on one fixed stratified 80/20 split ($n_{\text{train}} = 15216$, $n_{\text{test}} = 3804$). Band: mean $\pm$ 1 SD (ddof = 1) for the reversible arm -- repeat-run variability at a fixed split, not a confidence or credible interval. All triples are reported in results/blocks/blocks.csv; nothing here was selected.