

## P118. The length of the sport practice in swimming: A survival analysis approach

Pedro Sobreiro<sup>1,2</sup>, Alfredo Silva<sup>1,2</sup>, Abel Santos<sup>1,2</sup>, Hugo Louro<sup>1,2</sup>, Guedes de Carvalho<sup>3,2</sup>, Ana Conceição<sup>1,4</sup>

1. Sport Sciences School of Rio Maior, ESDRM-IPSANTARÉM, Rio Maior, Portugal; [sobreiro@esdrm.ipsantarem.pt](mailto:sobreiro@esdrm.ipsantarem.pt); 2. Life Quality Research Centre, CIEQV, Santarém, Portugal; 3. University Beira Interior, Faculty of Human and Social Sciences, Covilhã, Portugal; 4. Research Center in Sports Sciences, Health Sciences & Human Development, CIDESD, Vila Real, Portugal

### INTRODUCTION

The main concern of a sport organisation should be creating and retaining customers, where retention is fundamental (Tharrett & Bedford, 2012) and increasing the customer lifetime value is essential for the evaluation of the performance of the organisations (Gupta et al., 2006).

Retention in sports contributes to the increase of the athletes' health promotion (Siedentop, 1983; Seefeldt & Vogel, 1986), economic benefits (Shephard, 1986; Wang, Pratt, Macera, Zheng, & Heath, 2004), and bigger capacity for the athletes to be developed into international competitors (Green & Oakley, 2001). The duration in the sport practice can be targeted using the dropout, analysed using survival analysis, which is well-suited to study the timing of events in longitudinal data (Singer & Willett, 1993). The aim of this study is to predict the survival time in the swimming practice and the factors that contribute to a longer duration.

### METHODS

The duration is analysed using survival analysis, measuring the time that a swimmer practices in a sport facility. The dataset as  $N = 2653$  customers ( $n = 1536$  female  $19.62 \pm 15.79$  years) and  $n = 1117$  ( $20.03 \pm 16.5$  years) from a Portuguese sport facility, retrieved using the e@sport (Cedis, Portugal) software, corresponding to the period from 1 June 2014 until 31 October 2017. All data was pseudonymized ensuring that it is not attributed to an identified or identifiable natural person.

### RESULTS

The survival time of the swimmers during the first 12 months of practise in a sport facility, showed that there is a risk of dropout of 52% (Figure 1), with an estimated median survival of more 22 months.

The multivariate Cox's regression identified the variables "Days without frequency", "Total amount billed", "Number access", "Number of contract renewals" and "month" contributing to the increase of the time of survival.

### CONCLUSIONS

There is an increasing interest in the dropout problem in sports, especially in the rapidly growing children's and youth sports, in which attrition is extremely high (Salguero, Gonzalez-Boto, Tuero, & Márquez, 2003). The understanding of which factors contribute to the survival time allows to identify risk factors and when should measures to reduce the dropout be applied, giving these

information to the managers to define lines of actions to reduce the dropout. The "Days without frequency", "Total amount billed", "Number access", "Number of contract renewals" and "month" increase the survival time of the swimmers giving indicators to support the development of countermeasures to reduce dropout.

### References

- Green, M., & Oakley, B. (2001). Elite sport development systems and playing to win: uniformity and diversity in international approaches. *Leisure Studies*, 20(4), 247–267. doi:10.1080/02614360110103598
- Gupta, S., Hanssens, D., Hardie, B., Kahn, W., Kumar, V., Lin, N., ... Sriram, S. (2006). Modeling Customer Lifetime Value. *Journal of Service Research*, 9(2), 139–155. doi:10.1177/1094670506293810
- Salguero, A., Gonzalez-Boto, R., Tuero, C., & Márquez, S. (2003). Identification of dropout reasons in young competitive swimmers. *The Journal of Sports Medicine and Physical Fitness*, 43(4), 530–534.




Figure 1. Kaplan- Meier survivor curve of the swimmers, between June 2014 until October 2017.

- Seefeldt, V., & Vogel, P. (1986). *The Value of Physical Activity*. Reston, VA: American Alliance for Health, Physical Education. <https://eric.ed.gov/?id=ED289866>
- Shephard, R. J. (1986). *Economic benefits of enhanced fitness*. Champaign, Illinois: Human Kinetics. Obtido de <https://www.cabdirect.org/cabdirect/abstract/19861840075>
- Siedentop, D. (1983). *Developing teaching skills in physical education* (2nd ed). Palo Alto, Calif: Mayfield Pub. Co.
- Singer, J. D., & Willett, J. B. (1993). It's About Time: Using Discrete-Time Survival Analysis to Study Duration and the Timing of Events. *Journal of Educational Statistics*, 18(2), 155–195. doi:10.3102/10769986018002155
- Tharrett, S. J., & Bedford, P. (2012). *Why People Join, Leave, and Stay with Health /Fitness Clubs: The Ultimate Handbook of Member Retention*. Healthy Learning.
- Wang, G., Pratt, M., Macera, C. A., Zheng, Z.-J., & Heath, G. (2004). Physical activity, cardiovascular disease, and medical expenditures in U.S. adults. *Annals of Behavioral Medicine*, 28(2), 88–94. doi:10.1207/s15324796abm2802\_3