

produce both a higher $n_0(\text{IP})$ and a larger acetyl proton chemical shift. Least-squares analysis of the correlation between $n_0(\text{IP})$ and chemical shift yields the equation

$$\text{IP}(n_0) = -2.45 + 4.71\delta$$

with a correlation coefficient of 0.97 for seven *para*-substituted acetophenones.

Experimental

Photoelectron spectra were determined on a Perkin-Elmer PS-18 photoelectron spectrometer with heated probe. Xenon and argon were used as internal calibrants and the resolution as measured on the 15.76 eV peak of Ar was 25–30 meV. Peak positions (band maxima) listed in Table 1 represent the average of at least five determinations. The acetophenones were commercial samples and were used without further purification.

Acknowledgments

Financial support by the Natural Sciences and Engineering Research Council of Canada and St. Francis Xavier University Council for Research is acknowledged. The cooperation and assistance of Professor K. N. Houk of Louisiana State University in whose laboratory the photoelectron spectra were determined is also gratefully acknowledged.

1. D. W. TURNER, C. BAKER, A. D. BAKER, and C. R. BRUNDLE. *Molecular photoelectron spectroscopy*. Wiley Interscience, New York. 1970. Chaps. 10, 11.
2. A. D. BAKER, D. P. MAY, and D. W. TURNER. *J. Chem. Soc. B*, 22 (1968).
3. T. KOBAYASHI and S. NAGAKURA. *Bull. Chem. Soc. Jpn.* 47, 2563 (1974).

4. R. EGDELL, J. C. GREEN, and C. N. R. RAO. *Chem. Phys. Lett.* 33, 600 (1975).
5. C. N. R. RAO. *Tetrahedron*, 32, 1561 (1976).
6. E. NAGY-FELSOBUKI and J. B. PEEL. *J. Electron Spectrosc.* 16, 397 (1979).
7. E. J. MCALDUFF, B. M. LYNCH, and K. N. HOUK. *Can. J. Chem.* 56, 495 (1978).
8. E. J. MCALDUFF and T. CHAN. *Can. J. Chem.* 56, 2714 (1978).
9. A. PULLMAN and B. PULLMAN. *Quantum biochemistry*. Wiley, London. 1963.
10. R. J. HODGES, D. E. WEBSTER, and P. B. WELLS. *J. Chem. Soc. Dalton*, 2571 (1972); 2577 (1972).
11. T. KOENIG, R. WIELOSEK, L. L. MILLER, and Y.-H. SO. *J. Am. Chem. Soc.* 99, 7061 (1977).
12. M. THOMPSON, P. A. HEWITT, and D. S. WOOLSCROFT. *Handbook of X-ray and ultraviolet photoelectron spectroscopy*. Heyden Press, London. 1977. Chapt. 10, p. 349.
13. C. D. JOHNSON. *The Hammett equation*. Cambridge University Press, London. 1973.
14. R. DITCHFIELD, J. E. DEL BENE, and J. A. POPLE. *J. Am. Chem. Soc.* 94, 703 (1972).
15. R. O. LOUTFY and R. O. LOUTFY. *Tetrahedron*, 29, 2251 (1973).
16. R. O. LOUTFY and R. O. LOUTFY. *J. Phys. Chem.* 76, 1650 (1972).
17. R. O. LOUTFY and R. O. LOUTFY. *J. Phys. Chem.* 77, 336 (1973).
18. R. O. LOUTFY and R. O. LOUTFY. *Can. J. Chem.* 50, 4050 (1972).
19. P. R. YOUNG and W. P. JENCKS. *J. Am. Chem. Soc.* 101, 3288 (1979).
20. Y. YUKAWA and Y. TSUNO. *Bull. Chem. Soc. Jpn.* 32, 965 (1959); 32, 971 (1959).
21. D. W. EARLS, J. R. JONES, and T. G. RUMNEY. *J. Chem. Soc. Perkin II*, 878 (1975).
22. G. A. CAPLIN. *Org. Magn. Reson.* 6, 99 (1974).

Erratum: A reformulation of the theory of unimolecular reactions

ANDREW W. YAU AND HUW O. PRITCHARD

Centre for Research in Experimental Space Science, York University, Downsview, Ont., Canada M3J 1P3

Received January 10, 1980

(Ref: *Can. J. Chem.* 56, 1389 (1978))

Five small but substantive errors in this paper have come to our attention:

- (i) In [14], γ_i should read λ_i .
- (ii) In eq. [26], κ_{rr} and κ_{ss} should be interchanged in the numerator.
- (iii) In eq. [42], $\tilde{n}_j$ should read $\tilde{n}_i$.
- (iv) In Table 2, the value of B_r for $\text{C}_2\text{H}_5\text{NC}$ should be 33.26 cm^{-1} .
- (v) Dr. J. M. Simmie (Galway) has found that the deposited algorithm yields slightly different

answers on different computing machines: the trouble is caused by a ubiquitous mixed-mode FORTRAN operation in the 59th instruction of the program which should read $\text{NMAX(I)} = 0.5\text{E}0 * (1.\text{E}0/\text{XE(I)} - 1.\text{E}0) + 1.\text{E}-4$; adding 1×10^{-4} at this point causes all machines to give the correct answers, or subtracting 1×10^{-4} causes all machines to give the answers deposited as a test case with the algorithm.