

CSA and BJS versus TWFE (aligned samples)

20 studies · Filled: paired $|t| > 1.96$ · Hollow: otherwise · Cross: fragile inference


CSA vs TWFE

263 retained / 353 comparisons; 31 capped (diamonds)



BJS vs TWFE

288 retained / 357 comparisons; 41 capped (diamonds)



Study 1 (2017)

Study 7 (2019)

Study 2 (2017)

Study 8 (2020)

Study 4 (2018)

Study 9 (2020)

Study 5 (2018)

Study 10 (2021)

Study 6 (2018)

Study 11 (2021)

Study 12 (2022)

Study 17 (2024)

Study 13 (2023)

Study 18 (2024)

Study 14 (2023)

Study 20 (2024)

Study 15 (2023)

Study 21 (2025)

Study 16 (2024)

Study 22 (2025)

Circles: static specifications. Squares: event-study averages, with pre and post entering separately. Pre averages are diagnostics, not treatment effects. Shares use retained comparisons, including flagged points. Significant + Reversal: opposite nonzero signs and a filled marker. Diamonds: coordinates capped at ± 5 ; displayed gaps may differ from actual gaps. Excluded if either estimate has effective clusters < 5 or maximum cluster leverage $> 50\%$. Crosses: effective clusters < 10 or maximum leverage $> 25\%$ for either estimate or their difference, or prior inference warnings; significance fill suppressed. Leverage uses squared outcome-linear weights by the existing bootstrap clusters; it is a design diagnostic, not a variance guarantee. Exact-union sample alignment does not equate estimator weights or estimands. Both axes divide by this pair's saved TWFE SE; the plotted gap is not the paired-difference t statistic. Common horizons are unchanged. Fill uses paired-bootstrap normal tests, without multiplicity adjustment. Dashed line: equality. Study 8 uses unrestricted TWFE; its restricted published fit is retained separately. Study 4 clusters 143 builders despite only three treatment-rollout regions.