


CSA and BJS versus TWFE

20 studies · Filled: paired $|t| > 1.96$ · Hollow: otherwise · Cross: fragile inference


CSA vs TWFE

263 retained / 353 comparisons; 32 capped (diamonds)



BJS vs TWFE

288 retained / 357 comparisons; 43 capped (diamonds)



- | | | | |
|------------------|-------------------|-------------------|-------------------|
| ● Study 1 (2017) | ● Study 7 (2019) | ● Study 12 (2022) | ● Study 17 (2024) |
| ● Study 2 (2017) | ● Study 8 (2020) | ● Study 13 (2023) | ● Study 18 (2024) |
| ● Study 4 (2018) | ● Study 9 (2020) | ● Study 14 (2023) | ● Study 20 (2024) |
| ● Study 5 (2018) | ● Study 10 (2021) | ● Study 15 (2023) | ● Study 21 (2025) |
| ● Study 6 (2018) | ● Study 11 (2021) | ● Study 16 (2024) | ● Study 22 (2025) |

Circles: static specifications. Squares: event-study averages, with pre and post entering separately. Pre averages are diagnostics, not treatment effects. Shares use retained comparisons, including flagged points. Significant + Reversal: opposite nonzero signs and a filled marker. Diamonds: coordinates capped at ± 5 ; displayed gaps may differ from actual gaps. Excluded if either estimate has effective clusters < 5 or maximum cluster leverage $> 50\%$. Crosses: effective clusters < 10 or maximum leverage $> 25\%$ for either estimate or their difference, or prior inference warnings; significance fill suppressed. Leverage uses squared outcome-linear weights by the existing bootstrap clusters; it is a design diagnostic, not a variance guarantee. TWFE retains its eligible sample; modern estimators use their own supported targets. Both axes divide by this pair's saved TWFE SE; the plotted gap is not the paired-difference t statistic. Common horizons are unchanged. Fill uses paired-bootstrap normal tests, without multiplicity adjustment. Dashed line: equality. Study 8 uses unrestricted TWFE; its restricted published fit is retained separately. Study 4 clusters 143 builders despite only three treatment-rollout regions.