

Characteristic	Overall N = 221 ¹	Intervention N = 108 ¹	Placebo N = 113 ¹
Age ²	42 (20 - 68)	41 (20 - 66)	43 (21 - 68)
Years of education ³	13 (4 - 21)	12 (7 - 21)	13 (4 - 20)
Highest education ³			
No formal education	17 (7.7%)	10 (9.3%)	7 (6.2%)
Lower secondary school	88 (40%)	42 (39%)	46 (41%)
Intermediate secondary school	70 (32%)	36 (33%)	34 (30%)
Upper secondary school	30 (14%)	13 (12%)	17 (15%)
University degree	16 (7.2%)	7 (6.5%)	9 (8.0%)
Employment status ³			
Unemployed	61 (28%)	29 (27%)	32 (28%)
In education/training	11 (5.0%)	6 (5.6%)	5 (4.4%)
Unskilled worker	8 (3.6%)	5 (4.6%)	3 (2.7%)
Worker	61 (28%)	32 (30%)	29 (26%)
Employee	51 (23%)	25 (23%)	26 (23%)
Self-employed	8 (3.6%)	4 (3.7%)	4 (3.5%)
Retired	21 (9.5%)	7 (6.5%)	14 (12%)
Marital status ³			
Single	144 (65%)	79 (73%)	65 (58%)
Married	44 (20%)	15 (14%)	29 (26%)
Divorced	29 (13%)	12 (11%)	17 (15%)
Widowed	4 (1.8%)	2 (1.9%)	2 (1.8%)
Gender ⁴			
Female	2 (0.9%)	2 (1.9%)	0 (0%)
Male	216 (98%)	105 (97%)	111 (99%)
Trans/Transsexual	1 (0.5%)	0 (0%)	1 (0.9%)
Other	1 (0.5%)	1 (0.9%)	0 (0%)
Missing	1	0	1
Sexual orientation ⁴			
Exclusively heterosexual	145 (66%)	72 (67%)	73 (65%)
Predominantly heterosexual	24 (11%)	10 (9.3%)	14 (13%)

¹ Mean (Min - Max); n (%)

² data extracted from case files by trained staff (SSQ-CF).

³ information provided by the participant (SSQ-P).

⁴ assessed with the ICD-11 screener (LIT), a self-report measure which items follow the ICD-11 diagnostic criteria for paraphilic disorders.

⁵ information provided by supervision officer (SSQ-SO); in the German legal system, probation service refers to supervision and support during a suspended sentence or post-release support after prison, while post-release supervision (Führungsaufsicht) is a court-ordered measure of control and assistance after imprisonment in a high security hospital.

⁶ data extracted from case files by trained staff; since the Static-99 is only defined for hands-on offenders, we used the reduced Static-99 as mentioned by [LIT]; all staff member have been trained for the Static-99.

⁷ information provided by supervision officer (SSQ-SO).

⁸ assessed with the PHQ-D (LIT).

Characteristic	Overall N = 221 ¹	Intervention N = 108 ¹	Placebo N = 113 ¹
Bisexual	26 (12%)	12 (11%)	14 (13%)
Predominantly homosexual	3 (1·4%)	2 (1·9%)	1 (0·9%)
Exclusively homosexual	11 (5·0%)	5 (4·6%)	6 (5·4%)
Asexual	3 (1·4%)	3 (2·8%)	0 (0%)
Other	8 (3·6%)	4 (3·7%)	4 (3·6%)
Missing	1	0	1
Offense type²			
Hands-on (at least one conviction §176ff StGB)	89 (40%)	44 (41%)	45 (40%)
Hands-off (only §184b StGB)	132 (60%)	64 (59%)	68 (60%)
Type of supervision⁵			
post-release supervision	45 (20%)	20 (19%)	25 (22%)
community supervision	176 (80%)	88 (81%)	88 (78%)
Static recidivism risk⁶			
Low-medium	180 (81%)	87 (81%)	93 (82%)
Medium-high risk	41 (19%)	21 (19%)	20 (18%)
Term of imprisonment (month)⁷			
Missing	3	1	2
Treatment ordered by court⁷			
Yes	124 (56%)	62 (57%)	62 (55%)
No	97 (44%)	46 (43%)	51 (45%)
Additional treatment²			
Yes	70 (32%)	33 (31%)	37 (33%)
No	151 (68%)	75 (69%)	76 (67%)
Major depressive syndrome⁸			
Yes	12 (5·4%)	6 (5·6%)	6 (5·3%)
No	209 (95%)	102 (94%)	107 (95%)
Other depressive syndromes⁸			
Yes	20 (9·0%)	12 (11%)	8 (7·1%)
No	201 (91%)	96 (89%)	105 (93%)
Panic syndrome⁸			

¹ Mean (Min - Max); n (%)

² data extracted from case files by trained staff (SSQ-CF).

³ information provided by the participant (SSQ-P).

⁴ assessed with the ICD-11 screener (LIT), a self-report measure which items follow the ICD-11 diagnostic criteria for paraphilic disorders.

⁵ information provided by supervision officer (SSQ-SO); in the German legal system, probation service refers to supervision and support during a suspended sentence or post-release support after prison, while post-release supervision (Führungsaufsicht) is a court-ordered measure of control and assistance after imprisonment in a high security hospital.

⁶ data extracted from case files by trained staff; since the Static-99 is only defined for hands-on offenders, we used the reduced Static-99 as mentioned by [LIT]; all staff member have been trained for the Static-99.

⁷ information provided by supervision officer (SSQ-SO).

⁸ assessed with the PHQ-D (LIT).

Characteristic	Overall N = 221 ¹	Intervention N = 108 ¹	Placebo N = 113 ¹
Yes	5 (2.3%)	2 (1.9%)	3 (2.7%)
No	216 (98%)	106 (98%)	110 (97%)
Other anxiety syndromes⁸			
Yes	3 (1.4%)	3 (2.8%)	0 (0%)
No	218 (99%)	105 (97%)	113 (100%)
Bulimia nervosa⁸			
Yes	0 (0%)	0 (0%)	0 (0%)
No	221 (100%)	108 (100%)	113 (100%)
Binge eating⁸			
Yes	8 (3.6%)	3 (2.8%)	5 (4.4%)
No	213 (96%)	105 (97%)	108 (96%)
Alcohol syndrome⁸			
Yes	11 (5.0%)	6 (5.6%)	5 (4.4%)
No	210 (95%)	102 (94%)	108 (96%)
Somatoform syndrome⁸			
Yes	17 (7.7%)	8 (7.4%)	9 (8.0%)
No	204 (92%)	100 (93%)	104 (92%)
Psychosocial functioning			
Not difficult at all	130 (59%)	60 (56%)	70 (62%)
Somewhat difficult	76 (34%)	37 (34%)	39 (35%)
Very difficult	13 (5.9%)	10 (9.3%)	3 (2.7%)
Extremely difficult	2 (0.9%)	1 (0.9%)	1 (0.9%)
6D30 Exhibitionistic disorder⁴			
Yes	1 (0.5%)	1 (0.9%)	0 (0%)
No	219 (100%)	107 (99%)	112 (100%)
Missing	1	0	1
6D31 Voyeuristic disorder⁴			
Yes	5 (2.3%)	4 (3.7%)	1 (0.9%)
No	215 (98%)	104 (96%)	111 (99%)
Missing	1	0	1

¹ Mean (Min - Max); n (%)

² data extracted from case files by trained staff (SSQ-CF).

³ information provided by the participant (SSQ-P).

⁴ assessed with the ICD-11 screener (LIT), a self-report measure which items follow the ICD-11 diagnostic criteria for paraphilic disorders.

⁵ information provided by supervision officer (SSQ-SO); in the German legal system, probation service refers to supervision and support during a suspended sentence or post-release support after prison, while post-release supervision (Führungsaufsicht) is a court-ordered measure of control and assistance after imprisonment in a high security hospital.

⁶ data extracted from case files by trained staff; since the Static-99 is only defined for hands-on offenders, we used the reduced Static-99 as mentioned by [LIT]; all staff member have been trained for the Static-99.

⁷ information provided by supervision officer (SSQ-SO).

⁸ assessed with the PHQ-D (LIT).

Characteristic	Overall N = 221 ¹	Intervention N = 108 ¹	Placebo N = 113 ¹
6D32 Pedophilic disorder⁴			
Yes	7 (3.2%)	4 (3.7%)	3 (2.7%)
No	213 (97%)	104 (96%)	109 (97%)
Missing	1	0	1
6D33 Coercive sexual sadism disorder⁴			
Yes	2 (0.9%)	2 (1.9%)	0 (0%)
No	218 (99%)	106 (98%)	112 (100%)
Missing	1	0	1
6D34 Frotteuristic disorder⁴			
Yes	1 (0.5%)	1 (0.9%)	0 (0%)
No	219 (100%)	107 (99%)	112 (100%)
Missing	1	0	1
6D35 Other paraphilic disorder (non-consenting)⁴			
Yes	1 (0.5%)	1 (0.9%)	0 (0%)
No	219 (100%)	107 (99%)	112 (100%)
Missing	1	0	1

¹ Mean (Min - Max); n (%)

² data extracted from case files by trained staff (SSQ-CF).

³ information provided by the participant (SSQ-P).

⁴ assessed with the ICD-11 screener (LIT), a self-report measure which items follow the ICD-11 diagnostic criteria for paraphilic disorders.

⁵ information provided by supervision officer (SSQ-SO); in the German legal system, probation service refers to supervision and support during a suspended sentence or post-release support after prison, while post-release supervision (Führungsaufsicht) is a court-ordered measure of control and assistance after imprisonment in a high security hospital.

⁶ data extracted from case files by trained staff; since the Static-99 is only defined for hands-on offenders, we used the reduced Static-99 as mentioned by [LIT]; all staff member have been trained for the Static-99.

⁷ information provided by supervision officer (SSQ-SO).

⁸ assessed with the PHQ-D (LIT).