

Characteristic	Placebo N = 113	Intervention N = 108	p-hat	95% CI	t	p-value	q-value
Module 1			0.540	0.464; 0.615	1.03	0.31	>0.99
Mean (SD)	1 (18)	4 (20)					
Median (Q1, Q3)	0 (-8, 12)	4 (-4, 12)					
Min, Max	-68, 64	-60, 76					
Module 2			0.492	0.41; 0.575	-0.178	0.86	>0.99
Mean (SD)	1 (12)	0 (16)					
Median (Q1, Q3)	0 (-4, 8)	0 (-8, 8)					
Min, Max	-40, 40	-48, 36					
Missing	19	17					
Module 3			0.447	0.358; 0.54	-1.12	0.26	>0.99
Mean (SD)	1 (19)	-2 (19)					
Median (Q1, Q3)	0 (-8, 12)	0 (-12, 6)					
Min, Max	-48, 72	-100, 48					
Missing	35	36					
Module 4			0.426	0.331; 0.527	-1.45	0.15	0.89
Mean (SD)	4 (18)	-1 (17)					
Median (Q1, Q3)	0 (-2, 8)	0 (-12, 8)					
Min, Max	-44, 64	-36, 48					
Missing	41	46					
Module 5			0.496	0.391; 0.603	-0.064	0.95	>0.99
Mean (SD)	-1 (18)	-2 (15)					
Median (Q1, Q3)	0 (-4, 4)	0 (-8, 8)					
Min, Max	-100, 40	-60, 28					
Missing	54	55					
Module 6			0.528	0.414; 0.639	0.479	0.63	>0.99
Mean (SD)	2 (12)	2 (13)					
Median (Q1, Q3)	0 (-4, 8)	0 (0, 8)					
Min, Max	-24, 32	-40, 28					
Missing	60	66					

Abbreviations: , 95% CI = Confidence interval of the estimated relative effect, Q1 = 25th percentile, Q3 = 75th percentile, *p-hat* = estimated relative effect, *q-value* = Holm-Bonferroni adjusted p-value, *t* = Brunner–Munzel test statistic for unpaired samples