

Module	pre N = 106	post N = 106	p	q-value
Module 1 (post)			<0·0001	<0·0001
Mean (SD)			12 (7)	11 (8)
Median (Q1, Q3)			11 (7, 16)	9 (4, 16)
Min, Max			0, 31	0, 35
Module 2 (post)			0·050	0·21
Mean (SD)			10 (8)	11 (8)
Median (Q1, Q3)			8 (4, 16)	10 (5, 14)
Min, Max			0, 35	0, 36
Missing			15	15
Module 3 (post)			0·041	0·21
Mean (SD)			12 (9)	10 (8)
Median (Q1, Q3)			10 (5, 18)	8 (5, 14)
Min, Max			0, 36	0, 32
Missing			31	31
Module 4 (post)			0·98	>0·99
Mean (SD)			10 (7)	10 (7)
Median (Q1, Q3)			8 (5, 14)	8 (5, 14)
Min, Max			0, 29	0, 37
Missing			38	38
Module 5 (post)			0·78	>0·99
Mean (SD)			9 (8)	10 (8)
Median (Q1, Q3)			7 (4, 13)	7 (4, 14)
Min, Max			0, 37	1, 32
Missing			51	51
Module 6 (post)			0·62	>0·99
Mean (SD)			10 (8)	10 (8)
Median (Q1, Q3)			7 (4, 14)	7 (4, 13)
Min, Max			1, 32	1, 35
Missing			53	53

Abbreviation: Q1 = 25th percentile, Q3 = 75th percentile, p_{adj} = Holm-Bonferroni adjusted p.