

Characteristic	Intervention N = 108	Placebo N = 113	p-hat	95% CI	t	p-value	q-value
<b>Module 1 (advised: 28 days)</b>			0.423	0.348, 0.501	-1.93	0.053	0.27
Mean (SD)	114 (110)	87 (76)					
Median (Q1, Q3)	69 (43, 151)	58 (36, 112)					
Min, Max	24, 691	23, 506					
Missing	2	7					
<b>Module 2 (advised: 28 days)</b>			0.486	0.401, 0.572	-0.315	0.75	>0.99
Mean (SD)	71 (61)	66 (56)					
Median (Q1, Q3)	45 (33, 78)	44 (33, 74)					
Min, Max	25, 312	22, 390					
Missing	24	22					
<b>Module 3 (advised: 28 days)</b>			0.407	0.319, 0.502	-1.92	0.055	0.27
Mean (SD)	73 (52)	60 (44)					
Median (Q1, Q3)	57 (36, 90)	42 (32, 74)					
Min, Max	27, 308	24, 246					
Missing	39	38					
<b>Module 4 (advised: 28 days)</b>			0.380	0.289, 0.481	-2.31	0.021	0.13
Mean (SD)	73 (60)	51 (35)					
Median (Q1, Q3)	46 (33, 91)	37 (30, 50)					
Min, Max	25, 325	24, 173					
Missing	47	45					
<b>Module 5 (advised: 28 days)</b>			0.494	0.384, 0.605	-0.101	0.92	>0.99
Mean (SD)	62 (46)	56 (34)					
Median (Q1, Q3)	44 (33, 84)	44 (33, 73)					
Min, Max	23, 223	28, 186					
Missing	58	58					
<b>Module 6 (advised: 28 days)</b>			0.387	0.28, 0.506	-1.86	0.063	0.27
Mean (SD)	43 (41)	38 (43)					
Median (Q1, Q3)	33 (23, 46)	27 (22, 35)					
Min, Max	15, 279	15, 317					
Missing	66	60					

Abbreviations: 95% *CI* = Confidence interval of the estimated relative effect, *Q1* = 25th percentile, *Q3* = 75th percentile, *p-hat* = estimated relative effect, *q-value* = Holm-Bonferroni adjusted p-value, *t* = Brunner–Munzel test statistic for unpaired samples