| Module | pre N = 98 | post N = 98 | p-hat | 95% CI | t | p-value | q-value |
|---|---|---|---|---|---|---|---|
| **Module 1 (post)** | | | 0·512 | 0.467, 0.557 | 0·535 | 0·59 | >0·99 |
| Mean (SD) | 42 (11) | 42 (11) | | | | | |
| Median (Q1, Q3) | 43 (34, 51) | 44 (34, 50) | | | | | |
| Min, Max | 12, 60 | 13, 60 | | | | | |
| **Module 2 (post)** | | | 0·488 | 0.45, 0.526 | -0·641 | 0·52 | >0·99 |
| Mean (SD) | 42 (12) | 41 (13) | | | | | |
| Median (Q1, Q3) | 44 (34, 50) | 42 (30, 53) | | | | | |
| Min, Max | 13, 60 | 12, 60 | | | | | |
| Missing | 16 | 16 | | | | | |
| **Module 3 (post)** | | | 0·520 | 0.483, 0.557 | 1·09 | 0·28 | >0·99 |
| Mean (SD) | 42 (14) | 43 (13) | | | | | |
| Median (Q1, Q3) | 44 (30, 53) | 46 (35, 53) | | | | | |
| Min, Max | 12, 60 | 12, 60 | | | | | |
| Missing | 24 | 24 | | | | | |
| **Module 4 (post)** | | | 0·511 | 0.475, 0.546 | 0·588 | 0·56 | >0·99 |
| Mean (SD) | 43 (13) | 43 (13) | | | | | |
| Median (Q1, Q3) | 46 (35, 53) | 47 (34, 54) | | | | | |
| Min, Max | 12, 60 | 12, 60 | | | | | |
| Missing | 32 | 32 | | | | | |
| **Module 5 (post)** | | | 0·516 | 0.478, 0.554 | 0·841 | 0·40 | >0·99 |
| Mean (SD) | 44 (13) | 44 (13) | | | | | |
| Median (Q1, Q3) | 48 (34, 54) | 48 (34, 55) | | | | | |
| Min, Max | 12, 60 | 14, 60 | | | | | |
| Missing | 43 | 43 | | | | | |

Abbreviations: , *95% CI* = Confidence interval of the estimated relative effect, *Q1* = 25th percentile, *Q3* = 75th percentile, *p-hat* = estimated relative effect, *q-value* = Holm-Bonferroni adjusted p-value, *t* = Brunner–Munzel test statistic for paired samples