

Characteristic	Placebo N = 98	Intervention N = 97	p-hat	95% CI	t	p-value	q-value
<b>Module 1 (post)</b>			0·594	0.512; 0.671	2·23	0·025	0·13
Median (Q1 – Q3)	1 (-3 – 5)	3 (-2 – 8)					
<b>Module 2 (post)</b>			0·586	0.497; 0.671	1·89	0·059	0·18
Median (Q1 – Q3)	0 (-4 – 3)	1 (-2 – 4)					
Missing	16	15					
<b>Module 3 (post)</b>			0·458	0.365; 0.554	-0·859	0·39	0·39
Median (Q1 – Q3)	0·0 (-2·0 – 4·0)	0·0 (-3·0 – 2·0)					
Missing	24	29					
<b>Module 4 (post)</b>			0·432	0.334; 0.535	-1·30	0·19	0·39
Median (Q1 – Q3)	0·0 (-2·0 – 4·0)	-1·0 (-5·0 – 2·0)					
Missing	32	39					
<b>Module 5 (post)</b>			0·621	0.508; 0.723	2·09	0·036	0·14
Median (Q1 – Q3)	0·0 (-2·0 – 4·0)	2·0 (0·0 – 6·0)					
Missing	43	50					

Abbreviations: , 95% CI = Confidence interval of the estimated relative effect, Q1 = 25th percentile, Q3 = 75th percentile, p-hat = estimated relative effect, q-value = Holm-Bonferroni adjusted p-value, t = Brunner–Munzel test statistic for unpaired samples