

Characteristic	Placebo N = 98	Intervention N = 97	p-hat	95% CI	t	p-value	q-value
<b>Module 1 (post)</b>			0.610	0.528; 0.685	2.62	0.0090	0.045
Median (Q1 – Q3)	1 (-3 – 4)	4 (-1 – 6)					
<b>Module 2 (post)</b>			0.559	0.47; 0.645	1.30	0.19	0.40
Median (Q1 – Q3)	0.0 (-3.0 – 2.0)	1.0 (-1.0 – 4.0)					
Missing	16	15					
<b>Module 3 (post)</b>			0.468	0.373; 0.564	-0.659	0.51	0.51
Median (Q1 – Q3)	1.0 (-2.0 – 3.0)	0.0 (-2.0 – 3.0)					
Missing	24	29					
<b>Module 4 (post)</b>			0.420	0.323; 0.525	-1.50	0.13	0.40
Median (Q1 – Q3)	0.0 (-2.0 – 3.0)	-0.5 (-4.0 – 2.0)					
Missing	32	39					
<b>Module 5 (post)</b>			0.608	0.495; 0.71	1.87	0.061	0.24
Median (Q1 – Q3)	0.0 (-2.0 – 2.0)	1.0 (0.0 – 4.0)					
Missing	43	50					

Abbreviations: , 95% CI = Confidence interval of the estimated relative effect, Q1 = 25th percentile, Q3 = 75th percentile, *p-hat* = estimated relative effect, *q-value* = Holm-Bonferroni adjusted p-value, *t* = Brunner–Munzel test statistic for unpaired samples