

Module	pre N = 97	post N = 97	p-hat	95% CI	t	p-value	q-value
Module 1 (post)			0·582	0.531, 0.633	3·18	0·0020	0·010
Mean (SD)	42 (11)	44 (9)					
Median (Q1, Q3)	43 (34, 49)	47 (39, 51)					
Min, Max	14, 60	22, 60					
Module 2 (post)			0·527	0.485, 0.57	1·28	0·21	0·62
Mean (SD)	45 (9)	46 (10)					
Median (Q1, Q3)	47 (41, 52)	48 (39, 54)					
Min, Max	23, 60	12, 60					
Missing	15	15					
Module 3 (post)			0·513	0.463, 0.563	0·519	0·61	0·62
Mean (SD)	47 (9)	48 (9)					
Median (Q1, Q3)	48 (40, 55)	48 (42, 55)					
Min, Max	27, 60	24, 60					
Missing	29	29					
Module 4 (post)			0·468	0.413, 0.522	-1·19	0·24	0·62
Mean (SD)	47 (9)	46 (10)					
Median (Q1, Q3)	48 (42, 55)	47 (39, 56)					
Min, Max	24, 60	23, 60					
Missing	39	39					
Module 5 (post)			0·572	0.524, 0.619	3·02	0·0040	0·016
Mean (SD)	46 (11)	49 (9)					
Median (Q1, Q3)	47 (37, 57)	49 (41, 57)					
Min, Max	23, 60	23, 60					
Missing	50	50					

Abbreviation: 95% CI = Confidence interval of the estimated relative effect, Q1 = 25th percentile, Q3 = 75th percentile, p-hat = estimated relative effect, q-value = Holm-Bonferroni adjusted p-value, t = Brunner–Munzel test statistic for paired samples.