


# Counting structures on the $n \times k$ grid graph

Peter Kagey


University of Southern California

Graduate Student Combinatorics Conference  
April 2019


# Commodore 64 (1/3)




# Commodore 64 (2/3)




# Commodore 64 (3/3)



# Javascript



# Javascript



## Counting grids




Figure 1: An illustration of the bijection between tiles with diagonal markings and tiles with quarter circles in opposite corners.

A295229: Number of tilings of the  $n \times n$  grid, using diagonal lines to connect the grid points.




Figure 2: An example of the  $a(2) = 6$  different ways to fill the  $2 \times 2$  grid with diagonal tiles up to dihedral action of the square.

$$a(n) = \begin{cases} \frac{1}{8}(2^{n^2} + 2 \cdot 2^{n(n+1)/2} + 3 \cdot 2^{n^2/2} + 2 \cdot 2^{n^2/4}) & n \text{ even} \\ \frac{1}{8}(2^{n^2} + 2 \cdot 2^{n(n+1)/2} + 2^{(n^2+1)/2}) & n \text{ odd} \end{cases}$$



## Other tiles




Figure 3: Eleven box-drawing characters placed on an  $15 \times 8$  grid

# Baby's first corollary

## Corollary (of Burnside's Lemma)


Let

- ▶  $t$  be the number of tiles,
- ▶  $q$  be the number of tiles symmetric under a  $90^\circ$  rotation,
- ▶  $h$  be the number of tiles symmetric under a  $180^\circ$  rotation,
- ▶  $d$  be the number of tiles symmetric under a diagonal reflection, and
- ▶  $v$  be the number of tiles symmetric under a vertical reflection.

Then the number of tilings up to symmetries of the square is

$$a(n) = \begin{cases} \frac{1}{8} \left( t^{n^2} + 2qt^{\frac{n^2-1}{4}} + ht^{\frac{n^2-1}{2}} + (v^n + d^n)t^{\frac{n^2-n}{2}} \right) & n \text{ odd} \\ \frac{1}{8} \left( t^{n^2} + 3t^{\frac{n^2}{2}} + 2t^{\frac{n^2}{4}} + 2d^n t^{\frac{n^2-n}{2}} \right) & n \text{ even} \end{cases}$$

# Pipe Mania



## Leaf-free subgraphs of the grid graph




Figure 4: One of the  $a_4(12) = 42650154782713601$  (42 quadrillion) subgraphs on the  $12 \times 4$  grid graph  $G_{12,4} = P_{12} \square P_4$ .

The number of leaf-free subgraphs of  $G_{n,2}$  grid, obeys the recurrence

$$a_2(1) = 1, \quad a_2(2) = 2$$

$$a_2(n) = 5a(n-1) - 5a(n-2).$$

## Leaf-free subgraphs: intermediate states




Figure 5: An example of a leaf-free subgraph with its states labeled




Figure 6: Two examples of transitions from states to their children

## Example: A System of Recurrences

The  $1 \times 2$  grid has initial conditions

$$a_{00}(1) = a_{11}(1) = 1$$

$$a_{10}(1) = a_{01}(1) = a_{12}(1) = a_{21}(1) = a_{22}(1) = 0,$$

and satisfies the system of first order homogeneous difference relations

$$a_{00}(n+1) = a_{00}(n) + a_{22}(n)$$

$$a_{01}(n+1) = a_{01}(n) + a_{21}(n) + a_{22}(n)$$

$$a_{10}(n+1) = a_{10}(n) + a_{12}(n) + a_{22}(n)$$

$$a_{11}(n+1) = a_{00}(n) + a_{11}(n) + a_{12}(n) + a_{21}(n) + 2a_{22}(n)$$

$$a_{12}(n+1) = a_{01}(n) + a_{21}(n) + a_{22}(n)$$

$$a_{21}(n+1) = a_{10}(n) + a_{12}(n) + a_{22}(n)$$

$$a_{22}(n+1) = a_{11}(n) + a_{12}(n) + a_{21}(n) + a_{22}(n).$$

# A single recurrence from a system of recurrences

## Theorem (Corollary of Cayley–Hamilton theorem)

*In a system of first order homogeneous linear difference equations,*

$$\begin{aligned}a^{(1)}(n+1) &= \alpha_{11}a^{(1)}(n) + \dots + \alpha_{1k}a^{(k)}(n) \\ \vdots &= \vdots \\ a^{(k)}(n+1) &= \alpha_{k1}a^{(1)}(n) + \dots + \alpha_{kk}a^{(k)}(n)\end{aligned}$$

*each equation satisfies the recurrence*

$$a^{(i)}(n) = -\beta_{k-1}a^{(i)}(n-1) - \dots - \beta_1a^{(i)}(n-k-1) - \beta_0a^{(i)}(n-k)$$

*for  $n > k$  where  $A = \{\alpha_{ij}\}_{i,j=1}^k$  is the coefficient matrix and*

$$m_A(x) = x^k + \beta_{k-1}x^{k-1} + \dots + \beta_1x + \beta_0$$

*is the minimal polynomial of  $A$ .*

## A single recurrence from a system of recurrences

$$\underbrace{\begin{bmatrix} a_{00}(n) \\ a_{01}(n) \\ a_{10}(n) \\ a_{11}(n) \\ a_{12}(n) \\ a_{21}(n) \\ a_{22}(n) \end{bmatrix}}_{\vec{a}(n)} = \underbrace{\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{bmatrix}}_{A^{n-1}}^{n-1} \underbrace{\begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}}_{\vec{a}(1)}$$

Let  $x^k + \beta_{k-1}x^{k-1} + \dots + \beta_1x + \beta_0$  be the minimal polynomial of  $A$ . Then

$$\begin{aligned} A^k &= -\beta_{k-1}A^{k-1} - \dots - \beta_1A - \beta_0 \\ A^{n-1}\vec{a}(1) &= -\beta_{k-1}A^{n-2}\vec{a}(1) - \dots - \beta_1A^{n-k}\vec{a}(1) - \beta_0A^{n-k-1}\vec{a}(1) \\ \vec{a}(n) &= -\beta_{k-1}\vec{a}(n-1) - \dots - \beta_1\vec{a}(n-k+1) - \beta_0\vec{a}(n-k) \end{aligned}$$



## Some conjectural recurrences

For  $k = 3, 4, 5$ ,  $a_k(n)$  the number of leaf-free subgraphs of the  $n \times k$  grid graph is conjectured to satisfy


$$\begin{aligned} a_3(n) = & 12a_3(n-1) - 6a_3(n-2) - 20a_3(n-3) \\ & - 5a_3(n-4) \end{aligned}$$

$$\begin{aligned} a_4(n) = & 36a_4(n-1) - 7a_4(n-2) - 201a_4(n-3) \\ & + 49a_4(n-4) + 20a_4(n-5) - 5a_4(n-6) \end{aligned}$$

$$\begin{aligned} a_5(n) = & 103a_5(n-1) + 1063a_5(n-2) - 1873a_5(n-3) \\ & - 20274a_5(n-4) + 44071a_5(n-5) - 10365a_5(n-6) \\ & - 20208a_5(n-7) + 5959a_5(n-8) + 2300a_5(n-9) \\ & - 500a_5(n-10) \end{aligned}$$

For  $k = 6$ , this is conjectured to be an 18-order recurrence.

# Subgraphs which satisfy linear recurrences



## Theorem (Faase, 1994)

Let  $G$  be an arbitrary finite graph and let  $H_n$  denote either the path graph  $P_n$  or the cycle graph  $C_n$  on  $n$  vertices, and let  $s(n)$  count the number of subgraphs  $S$  of the Cartesian product  $G \square H_n$  subject to any combination of the following properties:

1. Restrictions on degree
2. Connectivity
3. Acyclicity

Then  $s(n)$  satisfies a linear recurrence.

# Examples: subgraphs which satisfy linear recurrences

Let  $G_{n,k}$  be a grid graph, then the following classes of subgraphs satisfy linear recurrences:

- ▶ Leaf-free subgraphs
  - ▶ Degree set  $D = \{0, 2, 3, 4\}$
- ▶ Spanning tree (mazes)
  - ▶ Degree set  $D = \{1, 2, 3, 4\}$
  - ▶ Connected
  - ▶ Acyclic
- ▶ Hamiltonian paths
  - ▶ Degree set  $D = \{1, 2\}$
  - ▶ Connected
  - ▶ Acyclic
- ▶ Perfect matchings (domino tilings)
  - ▶ Degree set  $D = \{1\}$




Figure 7: A correspondence between mazes and spanning trees

## Generalizing domino tilings

# More subgraphs which satisfy linear recurrences

## Theorem

Let  $H$  be a group and  $X$  be an arbitrary finite graph upon which  $H$  acts. Then let  $s(n)$  count the number of subsets of  $X \square P_n$  subject to any combination of the following properties:

1. Exactly/fewer than/more than  $m$  vertices of degree  $d$
2. Exactly/fewer than/more than  $c$  connected components on exactly/fewer than/more than  $d$  vertex.
3. An odd/even number of vertices of degree  $d$
4. Exactly  $\ell$  connected components
5. Fixed by the induced action of  $h \in H$

Then  $s(n)$  satisfies a linear recurrence.

## Corollary

The number of subgraphs of  $X \square P_n$  (with appropriate vertex degree, connectivity, or acyclicity restrictions) counted up to the group action of  $H$  satisfies a linear recurrence.


## Up to horizontal/vertical reflection

The number of no-leaf subgraphs of the  $2 \times n$  grid satisfies the two term recurrence

$$a_2(n) = 5a_2(n-1) - 5a_2(n-2).$$

The number of no-leaf subgraphs of the  $2 \times n$  grid up to horizontal/vertical reflection is conjectured to satisfy the eight term recurrence

$$\begin{aligned} s(n) = & 8s(n-1) - 16s(n-2) - 20s(n-3) + 95s(n-4) \\ & - 60s(n-5) - 80s(n-6) + 100s(n-7) - 25s(n-8) \end{aligned}$$



## Example: Möbius ladder (Guy, 1967)




Figure 8: The Möbius ladder  $M_{16}$  on 16 vertices.

### Fact

*The number of leaf-free subgraphs on the Möbius ladder on  $2n$  vertices is equal to the number of leaf-free subgraphs on the  $(n+2) \times 2$  grid graph.*

# Cycling around (a generalization of the Möbius ladder)

## Theorem

Let  $G$  be an arbitrary finite graph with vertex set  $V = \{v_1, v_2, \dots, v_m\}$ , and let  $E \subseteq V \times V$ . Then let  $H_n$  be the graph Cartesian product  $G \square P_n$  together with the edges  $\{((1, v_i), (n, v_j)) : (v_i, v_j) \in E\}$

Next let  $s(n)$  count the number of subgraphs of  $H_n$  subject to any combination of the following properties:

1. Restrictions on degree
2. Connectivity
3. Acyclicity

Then  $s(n)$  satisfies a linear recurrence.

Note that  $E = \{(v_1, v_1), (v_2, v_2), \dots, (v_m, v_m)\}$  recovers  $G \square C_n$  and  $E = \emptyset$  recovers  $G \square P_n$ .



## More exotic connections

### Theorem

*Let  $G$  be an arbitrary finite graph with vertex set  $V = \{v_1, v_2, \dots, v_m\}$ , and let  $E \subseteq V \times V$ . Then let  $H_n$  be the disjoint union product  $\underbrace{G \sqcup G \sqcup \dots \sqcup G}_{n \text{ times}}$  together with the edges*


$$\{((k, v_i), (k+1, v_j)) : (v_i, v_j) \in E, k \in [n-1]\}.$$

*Next let  $s(n)$  count the number of subgraphs of  $H_n$  subject to any combination of the properties mentioned in the previous theorems. Then  $s(n)$  satisfies a linear recurrence.*

Note that  $E = \{(v_1, v_1), (v_2, v_2), \dots, (v_m, v_m)\}$  recovers  $G \square P_n$ .

## Example: king graph

Let  $P_5$  be the path labeled  $v_1, v_2, \dots, v_5$  in the obvious way, and let  $E = \{(v_i, v_{i+1}) : i < 5\} \cup \{(v_i, v_i) : i \in [5]\} \cup \{(v_i, v_{i-1}) : i > 1\}$ .



**Figure 9:** The number of ways a king can tour every square of the  $5 \times 15$  chessboard is given by the number of Hamiltonian subgraphs in the  $5 \times 15$  king graph

The number of kings tours on a  $k \times n$  chessboard is a linear recurrence in  $n$ .

# Generalizations

- ▶ Is there always a nice algorithm for counting subgraphs of the  $n \times n$  grid graph satisfying particular properties?
  - ▶ There are nice ways of counting domino tilings and spanning trees.
- ▶ Given some graph and some properties, what is the order of recurrence that counts the number of subgraphs satisfying those properties?