

# Closed Geodesics: Topological Existence Theorems

Peter Kagey

April 25, 2019

A note to begin: I wrote this paper with a particular audience in mind, namely me in February of this year. I tried to capture a few things: (1) intuition for the topic (2) rigorous definitions, (3) early theorems and consequences, (4) modern theorems, and (5) conjectures and open problems.

## 1 Preliminaries

The intuition in this paper is to think of a creature who lives in a manifold walking in a “straight” line in some sense at a constant (nonzero) speed. In particular, the examples I try to keep in mind is the torus  $\mathbb{R}^2/\mathbb{Z}^2$ , the round sphere  $S^2 \subset \mathbb{R}^3$ , a very “wobbly” sphere, and (to avoid thinking about only so-called “geodesically complete” 2-manifolds)  $\mathbb{R}^3 \setminus \{0\}$ .

In the case where  $\mathbb{R}^n$ , the notion of acceleration along a path at some point in time is easy to naively define, since the notion of adding velocity vectors at two different points along the curve can be done naively, and the second derivative works in the expected way

$$\gamma''(t) = \lim_{h \rightarrow 0} \frac{\gamma'(t+h) - \gamma'(t)}{h}.$$

However, one has to be much more careful when talking about acceleration in a general manifold. In particular,  $T_{\gamma(t)}$  and  $T_{\gamma(t+h)}$  are generally different tangent spaces, and the notion of adding their vectors is decidedly more subtle. In particular, if a manifold is isometrically embedded into Euclidean space via  $f: M \rightarrow \mathbb{R}^n$ , the velocity along the image of a curve is preserved, but the acceleration is not quite the same—instead, the acceleration in  $M$  corresponds to the acceleration in  $f(M)$  of the projection of the acceleration to the tangent plane. However, capturing this notion intrinsically is more subtle.

**Definition 1.1** (Connection). Let  $\pi: E \rightarrow M$  be a smooth vector bundle over  $M$ . Then a connection in  $E$  is any map that sends a tangent vector field and a section of  $E$  to another section of  $E$

$$\nabla: \mathfrak{X}(M) \times \Gamma(E) \rightarrow \Gamma(E)$$

denoted  $\nabla_X Y := \nabla(X, Y)$  and satisfying the following three properties:

- (i)  $\nabla_{f_1 X_1 + f_2 X_2} Y = f_1 \nabla_{X_1} Y + f_2 \nabla_{X_2} Y$ ,
- (ii)  $\nabla_X (a_1 Y_1 + a_2 Y_2) = a_1 \nabla_X Y_1 + a_2 \nabla_X Y_2$ , and
- (iii)  $\nabla_X (fY) = f \nabla_X Y + \underbrace{(Xf)}_{\in C^\infty(M)} Y$ .

Connections are generalizations of directional derivatives with respect to  $X$ , a tangent vector field.

*Note 1.1.* In the case where  $\Gamma(E) = C^\infty(M)$ ,  $\nabla(X, f) := Xf$  satisfies the above properties. In particular,  $\nabla$  can be thought of as a prescription for extending  $X: C^\infty(M) \rightarrow C^\infty(M)$  to act on other vector bundles.

**Example 1.1.** Consider the case of  $M = \mathbb{R}^3$  with the section of the tangent bundle given by  $(x, 1, e^y) = X \in \mathfrak{X}(\mathbb{R}^3)$  and the rank 2 vector bundle  $(x^2, y + z) = Y \in \Gamma(E)$ . Then taking the derivative of  $Y$  along  $X$  yields

$$\nabla_X Y = \underbrace{\begin{bmatrix} \frac{\partial x^2}{\partial x} & \frac{\partial x^2}{\partial y} & \frac{\partial x^2}{\partial z} \\ \frac{\partial(y+z)}{\partial x} & \frac{\partial(y+z)}{\partial y} & \frac{\partial(y+z)}{\partial z} \end{bmatrix}}_{dY(x,y,z)} \underbrace{\begin{bmatrix} x \\ 1 \\ e^y \end{bmatrix}}_X = \begin{bmatrix} 2x^2 \\ 1 + e^y \end{bmatrix}.$$

The matrix multiplication perspective makes it clearer to see that in this case (i)  $\nabla$  is  $C^\infty(\mathbb{R}^3)$ -linear in the first term, because the column vector is  $C^\infty(\mathbb{R}^3)$ -linear, (ii)  $\nabla$  is  $\mathbb{R}$ -linear in the second term because both partial derivatives and matrices are  $\mathbb{R}$ -linear, and (iii) holds by doing the product rule entrywise in the Jacobian matrix  $d(fY)(x, y, z)$ .

These conditions capture, in some sense, the “essential qualities” of taking the directional derivative of a vector bundle over a given section of the tangent bundle.

**Definition 1.2** (Covariant derivative along a curve). Let  $\nabla$  be a connection in tangent bundle  $TM$ , and let  $\gamma$  be a curve in  $M$ . Then the **covariant derivative along  $\gamma$**  is the (unique) map

$$D_\gamma: \mathfrak{X}(\gamma) \rightarrow \mathfrak{X}(\gamma)$$

satisfying the following three properties

- (i)  $D_\gamma(aV + bW) = aD_\gamma V + bD_\gamma W$
- (ii)  $D_\gamma(fV) = f'V + fD_\gamma V$
- (iii)  $D_\gamma V(t) = \nabla_{\gamma'(t)} \tilde{V}$  for every extension  $\tilde{V}$  of  $V$ .

*Note 1.2.* This definition is sensible, because (i) and (ii) correspond to (ii) and (iii) in the definition of a connection, and the last condition is roughly equivalent to (i) in the definition of a connection: that is, the covariant derivative is  $C^\infty(M)$ -linear in the first component, so it only depends on the vector field pointwise.

When on a Riemannian manifold, it is common to use a particular connection called the Levi-Civita connection, which is compatible with the metric in the expected way, and which has some other nice properties.

**Definition 1.3** (Levi-Civita connection). The **Levi-Civita connection** is a connection  $\nabla$  such that

- (a) the metric is preserved under  $\nabla$ , that is,  $\nabla_{(-)} g \equiv 0$ , and
- (b) for any vector fields  $X, Y \in (M)$ ,  $\nabla_X Y - \nabla_Y X = [X, Y]$ .

*Note 1.3.* When the Levi-Civita connection exists, it is unique.

*Note 1.4.* A connection that satisfies the second condition in the definition of the Levi-Civita connection is called “torsion-free”, because the torsion is defined as

$$T(X, Y) := \nabla_X Y - \nabla_Y X - [X, Y].$$

**Definition 1.4** (Acceleration of a curve). For every smooth map from an interval to the manifold,  $\gamma: I \rightarrow M$ , define the **acceleration of  $\gamma$**  as the vector field  $D_\gamma \gamma'$  along  $\gamma$ .

**Definition 1.5** (Geodesic). A smooth curve  $\gamma$  is called a **geodesic** (with respect to  $\nabla$ ) if it’s acceleration is zero:  $D_\gamma \gamma' \equiv 0$ .

*Theorem 1.1* (Existence and uniqueness for geodesics). Let  $M$  be a smooth manifold and  $\nabla$  a connection. Then for every  $p \in M$ ,  $v \in T_p M$  and  $t_0 \in \mathbb{R}$ , there exists an open interval  $I \subset \mathbb{R}$  containing  $t_0$  and a geodesic  $\gamma: I \rightarrow M$  satisfying  $\gamma(t_0) = p$  and  $\gamma'(t_0) = v$ .

Furthermore, any two geodesics with the same initial conditions agree on their common domain.

*Proof idea.* This result follows from moving to local coordinates in a way that respects the connection, and then making an appeal to uniqueness and existence for (second order) ordinary differential equations.  $\square$

**Definition 1.6** (Geodesic extension). Given a geodesic  $\gamma: I \rightarrow M$ , an **extension of  $\gamma$**  is a map  $\tilde{\gamma}: \tilde{I} \rightarrow M$  such that

- (i)  $I$  is a subset of  $\tilde{I}$ ,
- (ii) the map  $\tilde{\gamma}$  is itself a geodesic, that is  $D_{\tilde{\gamma}}\tilde{\gamma}' \equiv 0$ , and
- (iii) the map agrees on the original interval, that is  $\tilde{\gamma}|_I = \gamma$ .

*Note 1.5.* A *proper* extension is an extension in which condition (i) is strengthened such that  $I$  is a *proper* subset of  $\tilde{I}$ .

*Note 1.6.* A geodesic with no proper extension is called a **maximal** geodesic.

**Definition 1.7** (Geodesically complete manifold). A manifold is called geodesically complete (with respect to  $\nabla$ ) if every geodesic curve  $\gamma$  has an “all-time” extension to  $\tilde{\gamma}: \mathbb{R} \rightarrow M$ .

**Definition 1.8** (Closed geodesic). A geodesic  $\gamma: I \rightarrow M$  is called **closed** if there exist two points  $t_1 \neq t_2 \in I$  such that  $\gamma(t_1) = \gamma(t_2)$  and  $\gamma'(t_1) = \gamma'(t_2)$ .

*Note 1.7.* Every closed geodesic  $\gamma$  has an all-time extension  $\tilde{\gamma}: \mathbb{R} \rightarrow M$ .

## 2 Properties of geodesics

An arbitrary manifold is not endowed with any “measuring tape” and creatures living on the manifold don’t have any notion of their speed when they travel. Of course, it’s natural to believe that having one gives the other—that is if our creature has a speedometer, it also has an odometer, and vice versa.

A Riemannian manifold comes equipped with an inner product on the tangent space at each point, and an inner product is equivalent to assigning a magnitude to each vector (by the polarization identity). That is, a Riemannian manifold is special because each tangent vector in  $TM$  has a notion of its magnitude or speed. This section will develop the natural way of turning the speed at each point to the length of a curve, and a natural way to define the distance between two points derived from the length of the *curves* between two points.

**Definition 2.1.** The **length of a vector**  $v \in TM$  in a Riemannian manifold with inner product  $g$  is

$$\|v\|_g = \sqrt{g(v, v)} = \sqrt{\langle v, v \rangle}.$$

**Definition 2.2.** The **length of a** (piecewise differentiable) **curve**  $\gamma: I \rightarrow M$  in a Riemannian manifold is

$$L(\gamma) = \int_I \|\gamma'(t)\|_g dt$$

*Theorem 2.1.* Let  $(M, g)$  be a Riemannian manifold. Then  $g$  induces a metric  $d: M \times M \rightarrow \mathbb{R}$  on  $M$ , namely

$$d(a, b) = \inf \{L(\gamma) : \gamma \text{ is a path from } a \text{ to } b\}$$

*Proof.* In order to show that  $d$  is a metric, it is necessary and sufficient to show that

1.  $d(a, b) = 0$  if and only if  $a = b$ .

( $\implies$ ) (A sketch, as this proof is a bit involved and technical.)

Choose some chart  $(U, \varphi)$  around  $p$  but not  $q$ , then in any compact set  $K \subset U$  around  $p$  there exists some  $C \in (0, \infty)$  such that for all  $x \in K$  and  $v \in T_x K$ , all vectors  $\|\varphi_* v\| \leq C\|v\|_g$ , which means that all curves in  $K$  have length greater than  $1/C$  times the length of their image under  $\varphi$ .

Then let  $R = \varphi^{-1}(B_\varepsilon(\varphi(p)))$  be the preimage of a small ball around  $\varphi(p)$ . By construction,  $p$  is inside this region,  $q$  is outside the region, and the distance to the boundary of the region is at least  $\varepsilon$ . Thus  $d(p, q) \geq \varepsilon$  by the Jordan Curve Theorem.

( $\Leftarrow$ ) If  $a = b$ , the constant path  $\gamma(t) = a = b$  has length 0, and the length of a path is strictly non-negative because the integrand is nonnegative:  $\|v\|_g \geq 0$ .

2.  $d(x, y) = d(y, x)$

If  $\gamma: (t_0, t_1) \rightarrow M$  is a path from  $a$  to  $b$ , then  $\bar{\gamma}: (t_0, t_1) \rightarrow M$  defined by  $\bar{\gamma}(t) = \gamma(t_1 + t_0 - t)$  is a path from  $b$  to  $a$  which has the same image and same length.

3.  $d(a, b) \leq d(a, c) + d(c, b)$

The set of pairs of paths from  $a$  to  $c$  and  $c$  to  $b$  is in length preserving bijection (via concatenation) with the set of paths from  $a$  to  $b$  through  $c$ , which is a subset of the set of paths from  $a$  to  $b$ .

□

**Definition 2.3.** A connected Riemannian manifold  $(M, g)$  is called **geodesically complete** if every geodesic curve  $\gamma: I \rightarrow M$  there is an extension of  $\gamma$  with  $\mathbb{R}$  as the domain,  $\tilde{\gamma}: \mathbb{R} \rightarrow M$ .

While the distance  $d(a, b)$  function on  $M$  is defined as the infimum under all paths from  $a$  to  $b$ , in the case of a geodesically complete (connected) manifold, it is equivalent to define distance as the *minimum* over all paths instead.

*Theorem 2.2.* In a geodesically complete Riemannian manifold, there exists a geodesic (with respect to the Levi-Civita connection)  $\gamma$  from  $(a, b)$  such that  $L(\gamma) = d(a, b)$ .

*Proof idea.* This follows from the Hopf-Rinow Theorem (below) along with some technical applications using the exponential map. □

Now is perhaps an appropriate time to introduce the exponential map. The idea is that the exponential map takes initial conditions of a geodesic (position and velocity), and (when possible) returns the position of a particle flowing along such a geodesic after unit time. The exponential map takes a point in the tangent bundle  $(p, v) \in TM$  and returns the position of a geodesic starting at  $p$  and flowing in the direction of  $v$  after unit time.

**Definition 2.4** (Exponential Map). Let

$$\mathcal{D} = \{(p, v) \in TM : \exists \text{ a geodesic curve } \gamma: [0, 1] \rightarrow M \text{ with } \gamma(0) = p \text{ and } \gamma'(0) = v\}$$

Then the **exponential map** is a map

$$\exp: \mathcal{D} \rightarrow M$$

satisfying  $\exp((p, v)) = \gamma(1)$  where  $\gamma: [0, 1] \rightarrow M$  is any geodesic satisfying  $\gamma(0) = p$  and  $\gamma'(0) = v$ .

*Note 2.1.* A geodesically complete (connected) manifold is equivalently defined as one in which  $\mathcal{D} = TM$ .

### 3 Manifolds as metric spaces

It is interesting to ask about whether or not a manifold is geodesically complete, that is whether or not the creature in the manifold can travel forever in a “straight line” starting at any point and heading in any direction.

Clearly the creature does not run into any trouble flying through space ( $\mathbb{R}^3$ ) or walking on the surface of a sphere ( $S^2$ ), but the creature clearly *does* run into trouble if it is on the surface of a disc ( $D^2 \subset \mathbb{R}^2$ ) or if it is flying toward the origin, but that point has been removed ( $\mathbb{R}^3 \setminus \{0\}$ ).

The Hopf-Rinow Theorem says that these two examples capture the extent of what can go wrong during the creature’s journey. In particular, it says that that properties of geodesics and the induced metric agree with each other and with the topology.

**Definition 3.1** (Complete metric space). A complete metric space is a metric space in which every Cauchy sequence is convergent.

*Theorem 3.1* ([4]). Let  $(M, g)$  be a Riemannian manifold, and let  $(M, d)$  be the induced metric space. Then the following are equivalent:

- (i)  $(M, d)$  is metrically complete.
- (ii)  $(M, g)$  is geodesically complete.
- (iii)  $(M, g)$  is geodesically complete at some point.
- (iv) Every closed, bounded set in  $M$  is compact.

And furthermore, between any two points  $p, q \in M$ , there exists a length-minimizing geodesic connecting these two points.

*Proof sketch.*

(i)  $\implies$  (ii) The idea is to assume that  $M$  is not complete and arrive at a contradiction. In particular, this means there is some geodesic curve  $\gamma: [a, b) \rightarrow M$  which can not be extended to  $\tilde{\gamma}: [a, b] \rightarrow M$ —however this can't happen because we can use the limit of a Cauchy sequence to give a value for  $\tilde{\gamma}(b)$  such that  $\tilde{\gamma}$  is a geodesic.

(ii)  $\implies$  (iii) If  $(M, g)$  is geodesically complete at every point, then in particular it is geodesically complete at some point.

(iii)  $\implies$  (iv) Suppose  $(M, g)$  is geodesically complete at  $p$ . For any closed and bounded  $A \subset M$ , we can always find a ball  $B_R(p)$  around  $A$ . Since the exponential map is continuous, we can see that a closed subset of a compact set is itself compact.

(iv)  $\implies$  (i) Let  $(x_n)$  be a Cauchy sequence in  $M$ . Since  $\{x_n\}$  is bounded, its closure is compact and so  $(x_n)$  has a convergent subsequence, and because it is Cauchy, it converges to this subsequence.  $\square$

## 4 Closed geodesics

*Theorem 4.1.* Let  $\pi_1(M)$  be the fundamental group of  $M$ . Then each equivalence class of loops  $[\alpha] \in \pi_1$  has a (closed) geodesic representative, in particular it contains a curve of minimal length which is a geodesic.

*Proof idea.* [8] summaries the proof as

One way to see why this is true is via a Morse-theory approach, with a full proof being given in Klingenberg - Lectures on Closed Geodesics. The gist of it is to pick a loop close to the infimum on the class and flow it under (minus) the gradient flow of the energy functional in the free loop space (this has an obvious analogy with the finite-dimensional case when we flow via gradient and converge to a critical point).

$\square$

*Note 4.1.* In a survey paper, [9] writes

one of the first successes of the calculus of variations was to establish rigorously that such a minimizing procedure is effective and produces a closed geodesic. The situation is subtler if the manifold is simply connected, and the question was answered in the affirmative by Lyusternik and Fet in their celebrated 1951 paper

*Theorem 4.2* ([6]). Every compact Riemannian manifold  $(M, g)$  has at least one closed geodesic.

*Proof.* In a report from the International Workshop on Geodesics in August 2010, [3] writes

Lyusternik and Fet considered the energy functional on the loop space  $\Lambda M$  and showed that the topology of  $\Lambda M$  is complicated enough so that the energy functional must have critical points with nonzero energy (which are non trivial closed geodesics).

□

**Conjecture 4.1** ([3]). Every compact Riemannian manifold of dimension greater than 1 contains infinitely many geometrically distinct non-constant closed geodesics.

*Note 4.2.* This particular conjecture in the above report by [3]:

How many closed geodesics must exist on a closed manifold? For surfaces, the answer is known: there are always infinitely many geometrically different closed geodesics. This is easily proved using Birkhoffs first argument when the fundamental group is infinite. The remaining cases of the sphere and the projective plane were settled by Bangert (1993) and Franks (1992). In higher dimensions Rademacher has shown that a closed manifold with a generic Riemannian metric admits infinitely many geometrically different closed geodesics Rademacher (1989).

The report gives a heuristic for why this is hard to prove:

It is important to distinguish geometrically different geodesics from repetitions of the same geodesic. This distinction is difficult to make.

*Note 4.3.* In 1999 [1] writes

The question of which Riemannian manifolds admit simple closed geodesics is still a mystery. It is not known whether all closed Riemannian manifolds contain simple closed geodesics.

*Theorem 4.3* ([7]; [2]). Every Riemannian manifold with the topology of a sphere has at least three simple closed geodesics.

*Proof.* The history of the proof from the [5] paper:

The existence of three simple periodic geodesics on every Riemannian 2-sphere was first proven by L.A. Lusternik and L.G. Shnirelman. They considered the space  $\Pi M$  of non-parametrized curves on a 2-dimensional Riemannian manifold  $M$  diffeomorphic to  $S^2$ , as well as its subset  $\Pi_0 M$  that consists of all constant curves (and, thus, can be identified with  $M$ ). [One can then] consider the three relative homology classes of the pair  $(\Pi S^2, \Pi_0 S^2)$  with coefficients in  $\mathbb{Z}_2$ , where we regard  $S^2$  as the unit round sphere in  $\mathbb{R}^3$

- 1) The 1-dimensional class represented by the relative 1-cycle  $z_1$  formed by all circles on  $S^2$  in planes parallel to the  $XZ$ -plane in the ambient  $R^3$ ;
- 2) The 2-dimensional class represented by the relative cycle  $z_2$  formed by all circles in planes parallel to the  $Z$ -axis; and
- 3) The 3-dimensional class represented by the relative 3-cycle  $z_3$  formed by all round circles on the sphere (including points regarded as degenerate circles).

Lusternik and Shnirelman described a curve shortening flow in  $\Pi M$ , [and] they observed that this flow gets stuck on critical points representing simple periodic geodesics, when applied to  $z_1$ ,  $z_2$ ,  $z_3$ , and proved that if two of these three geodesics coincide, then there is a whole critical level with a 1-parametric set of distinct simple periodic geodesics. Some errors in the construction of their curve shortening flow had been later corrected by W. Ballman [and others]. Alternatively, one can prove the existence of three simple periodic geodesics using either the curvature flow and Graysons theorem or an especially simple curve shortening flow constructed by J. Hass and P. Scott.

□

In particular, a recent theorem shows an even stronger result: the lengths of all three geodesics on a sphere are bounded by the “size” of the manifold.

**Definition 4.1.** The **diameter** of a Riemannian manifold  $(M, g)$  is defined by

$$\text{diam}(M) := \sup_{p, q \in M} d(p, q) \in \mathbb{R}_{\geq 0} \cup \{+\infty\}$$

**Example 4.1.** Consider the unit circle  $S^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$  as a sub-manifold of  $\mathbb{R}^2$  with the induced inner product. Then  $\text{diam}(S^1) = \pi$ , with the distance between two antipodal points, say  $p = (1, 0)$  and  $q = (-1, 0)$ , being half of the circumference of the circle (in the ordinary sense).

**Example 4.2.** Consider the torus  $T^2 = \mathbb{R}^2/\mathbb{Z}^2$  with the inner product inherited from  $\mathbb{R}^2$ . Then the diameter of the torus  $\text{diam}(T^2) = \sqrt{2}/2$ , with this distance achieved by  $p = (0, 0)$  and  $q = (\frac{1}{2}, \frac{1}{2})$ .




Figure 1: Illustrations of diameters on  $S^1$  and  $T^2$ , and geodesics that achieve them.

*Theorem 4.4* ([5]). Let  $M$  be a Riemannian 2-sphere. Then there exist three distinct simple geodesics with lengths that do not exceed  $20d$  where  $d$  is the diameter of  $M$ . Furthermore, if no simple closed geodesics of length  $\leq 2d$ , then there are three distinct simple periodic geodesics on  $M$  with lengths  $\leq 5d$ ,  $10d$ , and  $20d$  respectively.

*Proof idea.* The above authors give their proof idea as follows:

The main idea of the proof of [the above theorem] is to express three homology classes of the space of non-parametrized curves that are used in classical proofs of the Lyusternik-Schnirelmann theorem by cycles that consist of simple closed curves “mainly made” of curves in a meridian-like family that connects two fixed points of  $M$ . [...] We attempt to construct such a family where the lengths of all meridians are bounded by  $\text{const } d$  for an appropriate  $\text{const}$ . Our repeated attempts can be blocked only by appearance of different short simple periodic geodesics of index 0. So, we either get three short simple periodic geodesics of index 0, or our third attempt to construct a meridional slicing succeeds. Once one of our attempts succeeds, and we get a slicing of  $M$  into short meridians, the original proof of the Lyusternik-Schnirelmann theorem yields the desired upper bounds.

□

*Note 4.4.* From [5]:

[There is a] classical result of M. Morse, who proved that the fourth periodic [not necessarily simple] geodesic becomes uncontrollably large for ellipsoids with distinct but very close semi-axes.

## References

- [1] Colin Adams, Joel Hass, and Peter Scott. “Simple Closed Geodesics in Hyperbolic 3-Manifolds”. In: *Bulletin of the London Mathematical Society* 31.1 (1999), pp. 81–86. DOI: 10.1112/S0024609398004883. eprint: <https://londmathsoc.onlinelibrary.wiley.com/doi/pdf/10.1112/S0024609398004883>.
- [2] Werner Ballmann. “Der Satz von Lusternik und Schnirelmann”. In: *Math. Schriften* 102 (1978), pp. 1–25.
- [3] Keith Burns and Vladimir S. Matveev. *Open problems and questions about geodesics*. 2013. eprint: arXiv:1308.5417.

- [4] Heinz Hopf and Willi Rinow. *About the concept of the complete differential geometric surface*. July July 1931. DOI: <https://doi.org/10.1007/BF01601813>.
- [5] Yevgeny Liokumovich, Alexander Nabutovsky, and Regina Rotman. *Lengths of three simple periodic geodesics on a Riemannian 2-sphere*. 2014. eprint: [arXiv:1410.8456](https://arxiv.org/abs/1410.8456).
- [6] L.A. Lyusternik and A.I. Fet. “Variational problems on closed manifolds”. In: *Dokl. Akad. Nauk SSSR (N.S.)* 81 (Jan. 1951), pp. 17–18.
- [7] Lazar Lyusternik and Lev Schnirelmann. “Sur le problème de trois géodésiques fermées sur les surfaces de genre 0”. In: *Comptes Rendus de l’Académie des Sciences de Paris* 189 (1929), pp. 269–271.
- [8] Aloizio Macedo. *Closed geodesic on a non-simply connected Riemannian manifold*. URL: <https://math.stackexchange.com/a/2844405/121988>. (accessed: 04.25.2019).
- [9] Alexandru Oancea. *Morse theory, closed geodesics, and the homology of free loop spaces*. 2014. eprint: [arXiv:1406.3107](https://arxiv.org/abs/1406.3107).