## Supporting Information for "Ship-based lidar evaluation of Southern Ocean low clouds in the storm-resolving general circulation model ICON and the ERA5 and MERRA-2 reanalyses"

Peter Kuma<sup>1,2,3,4</sup>, Frida A.-M. Bender<sup>1,2</sup>, Adrian J. McDonald<sup>3</sup>, Simon P.

Alexander<sup>5,6</sup>, Greg M. McFarquhar<sup>7,8</sup>, John J. Cassano<sup>9,10,11</sup>, Graeme E.

Plank<sup>3</sup>, Sean Hartery<sup>3</sup>, Simon Parsons<sup>3</sup>, Sally Garrett<sup>12</sup>, Alex J.

Schuddeboom<sup>3</sup>, and Anna Possner<sup>4</sup>

<sup>1</sup>Department of Meteorology (MISU), Stockholm University, Stockholm, Sweden

<sup>2</sup>Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

<sup>3</sup>School of Physical and Chemical Sciences, University of Canterbury, Christchurch, Aotearoa/New Zealand

<sup>4</sup>Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Frankfurt am Main, Hesse, Germany

<sup>5</sup>Australian Antarctic Division, Kingston, Tasmania, Australia

<sup>6</sup> Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania,

## Australia

<sup>7</sup>Cooperative Institute of Severe and High Impact Weather Research and Operations, University of Oklahoma, Norman, OK, USA

<sup>8</sup>School of Meteorology, University of Oklahoma, Norman, OK, USA

<sup>9</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

<sup>10</sup>National Snow and Ice Data Center, University of Colorado, Boulder, CO, USA

<sup>11</sup>Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, CO, USA

<sup>12</sup>New Zealand Defence Force, Wellington, Aotearoa/New Zealand

## Contents of this file

- 1. Figure S1
- 2. Figure S2
- 3. Figure S3

Corresponding author: P. Kuma, Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Altenhöferallee 1, Frankfurt am Main, 60438, Hesse, Germany. (peter@peterkuma.net)

\*Current affiliation: Department of

Physics & Atmospheric Science, Dalhousie

University, Halifax, Canada

<sup>†</sup>Current affiliation: New South Wales

Department of Planning and Environment,

Sydney, New South Wales, Australia




Figure S1. Cloud occurrence by height for the 31 voyages and one sub-Antarctic station (MICRE) in observations (O) and simulated by the ALCF from the ICON model (I), MERRA-2 (M), and ERA5 reanalysis data (E). The numbers in the legend indicate the total cloud fraction and the number of days of data. Multiple lines of ICON profiles are for each of the four years of model data available.




Figure S2. The same as Fig. 9, but only for profiles that contain (a) fog, (b) cloud at 500 m, and (c) cloud at 1.5 km.




Figure S3. The same as Fig. 11, but for virtual potential temperature.