

Peter Harding

November 21, 2023

Abstract

Deakin Uni Physics for the Life Sciences Notes

1 Constants

$$c = 3.00 \times 10^8 \text{ m/sec}$$

$$e = 1.60 \times 10^{-19} \text{ C}$$

$$g = 9.8 \text{ m/sec}^2$$

$$1 \text{ atm} = 1.01 \times 10^5 \text{ Pa} = 760 \text{ mmHg}$$

$$\text{Coulomb's } K = 9 \times 10^9 \text{ Nm}^2\text{C}^{-2}$$

$$\text{Speed of Sound} = 343 \text{ m/sec}$$

$$1 \text{ Cal} = 4.186 \text{ J}$$

$$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$$

$$\text{Electron Mass} = 9.11 \times 10^{-31} \text{ Kg}$$

$$\text{Proton Mass} = 1.67 \times 10^{-27} \text{ Kg}$$

$$\text{Atomic Mass Unit} = 1.67 \times 10^{-27} \text{ Kg}$$

$$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2$$

$$R = 8.31 \text{ J/mol} - \text{K}$$

$$\text{Threshold of hearing} = I_0 = 1.0 \times 10^{-12} \text{ W/m}^2$$

$$1 \text{ curie} = 3.7 \times 10^{10} \text{ Bq}$$

$$k_B = 1.38 \times 10^{-23} \text{ J/K}$$

$$R = 8.31 \text{ J/mol} - \text{K}$$

$$\text{Speed of Light} = 3.0 \times 10^8 \text{ m/sec}$$

$$\hbar = 1.05 \times 10^{-34} \text{ J} \cdot \text{s} = 6.58 \times 10^{-16} \text{ eV} \cdot \text{s}$$

$$\text{Density of Water} = 1000 \text{ kg/m}^3$$

2 Conversion Formulae

$$T = T_c + 273$$

$$T(^{\circ}C) = \frac{5}{9}[T(^{\circ}F) - 32^{\circ}]$$

$$n = \frac{M \text{ (in grams)}}{M_{mol}} = \frac{N}{N_A}$$

2.1 Trigonometry

$$\textit{cosine} = \frac{\textit{adjacent}}{\textit{hypotenuse}}$$

$$\textit{sine} = \frac{\textit{opposite}}{\textit{hypotenuse}}$$

$$\textit{tangent} = \frac{\textit{opposite}}{\textit{adjacent}}$$

2.2 Pythagorean Theorem

$$a^2 = b^2 + c^2$$

3 Formulae

3.1 Area and Volume Formulae

Circle	$A = 2\pi r$	$V = \pi r^2$
Cylinder	$A = 2\pi r h$	$V = \pi r^2 h$
Sphere	$A = 4\pi r^2$	$V = \frac{4}{3}\pi r^3$
Cube	$A = 6h^2$	$V = h^3$

3.2 Length of a vector

Length of a Vector	$ V = \sqrt{V_x^2 + V_y^2}$
--------------------	------------------------------

4 Kinematics

Equations of Motion	$\Delta x = x_j - x_i$ $v = \frac{\Delta x}{\Delta t}$ $a = \frac{\Delta v}{\Delta t}$ $v_f^2 = v_i^2 + 2a\Delta x$ $v_f = v_i + a\Delta t$ $\Delta x = v_i\Delta t + \frac{1}{2}a\Delta t^2$
---------------------	--

Net Force	$F_{net} = F_1 + F_2 + \dots + F_N$
-----------	-------------------------------------

Newton's Second Law	$F = ma$
---------------------	----------

Drag Force	$D = \frac{1}{2}C_d\rho v^2 A$
------------	--------------------------------

Friction Force	$f_{s,max} = \mu_s n$	$f_k = \mu_k n$
----------------	-----------------------	-----------------

Spring Force (Hook's Law)	$F = -kx$
---------------------------	-----------

Conservation of Energy (without transfer)	$\Delta K + \Delta U + \Delta E_{th} = 0$
---	---

Conservation of Energy (with transfer)	$\Delta K + \Delta U + \Delta E_{th} = w + q$
--	---

Kinetic Energy:	$K = \frac{1}{2}mv^2$
-----------------	-----------------------

Gravitational Potential Energy:	$U_g = mgy$
---------------------------------	-------------

Spring Potential Energy:	$U_x = \frac{1}{2}kx^2$
--------------------------	-------------------------

Work:	$W = Fd(\cos\theta)$
-------	----------------------

Power:	$P = \frac{\Delta E}{\Delta t}$
--------	---------------------------------

Mechanical Power:	$P = \frac{W}{\Delta t} = Fv$
-------------------	-------------------------------

Energy Efficiency:	$e = \frac{E_{out}}{E_{in}}$
--------------------	------------------------------

4.1 Thermal Properties

	$T = \frac{2}{3} \frac{K_{avg}}{k_b}$	
Thermal expansion (volume):	$\Delta V = \beta V, \Delta T$	
Thermal expansion (linear):	$\Delta L = \alpha L, \Delta T$	
Gas Pressure	$p = \frac{2}{3} \frac{N}{V} K_{avg}$	
Ideal Gas Law (multiple forms)	$pV = NK_bT$	$= nN_A k_b T = nRT$
Mass of a Substance	$m = \rho V$	
Work Done by a Gas	$W_{gas} = p\Delta V$	
Energy Conservation for Interacting Systems	$Q_{net} = Q_1 + Q_2 + \dots = 0$	
Heat Equations for Solids and Liquids	$Q = mC\Delta T$	
Heat Equations for Gasses		
Constant Volume	$Q = nC_v\Delta T$	
Constant Pressure	$Q = nC_p\Delta T$	
Heat Required for a Phase Change		
Fusion (melting/freezing)	$Q = \pm ML_f$	
Vapourization (boiling/condensing)	$Q = \pm ML_v$	
Rate of Conduction Across a Temperature Gradient	$Q = \frac{kA}{L}\Delta T$	
Rate of Radiative Heat Transfer	$\frac{Q}{\Delta T} = e \sigma AT^4$	

5 Fluids

Fluid Pressure	$p = \frac{F}{A}$
Hydrostatic Pressure	$p = p_0 + \rho g d$
Gauge Pressure	$p_g = p - p_{atm}$
Pressure Gradient in a Viscous Fluid	$\Delta p = 8\pi\nu \frac{LV_{avg}}{A}$
Buoyancy Force	$F_b = \rho_f V_f g$

6 Oscillations

Frequency-period relationship $f = \frac{1}{T}$

Frequency of mass-spring oscillator $f = \frac{1}{2\pi} \sqrt{\frac{k}{m}}$

Frequency of pendulum oscillator (small angle of displacement) $f = \frac{1}{2\pi} \sqrt{\frac{g}{L}}$

7 Waves

$$\text{Wave speed of a stretched string} \quad v_{string} = \sqrt{\frac{T_s}{\mu}}$$

$$\text{Wave speed in a gas} \quad v_{sound} = \sqrt{\frac{\gamma RT}{M}}$$

$$\text{Wave speed} \quad v = f\lambda = \frac{\lambda}{T}$$

$$\text{Sound intensity} \quad \beta = (10dB)\log_{10}\left(\frac{I}{I_0}\right)I = \frac{P_{source}}{4\pi r^2}$$

8 Doppler Effect

$$\text{Observed frequency (source approaching at } v_s) \quad f_+ = \frac{f_0}{1 - v_a/v}$$

$$\text{Observed frequency (source receding at } v_s) \quad f_- = \frac{f_0}{1 + v_a/v}$$

$$\text{Observed frequency (observer approaching source at } v_o) \quad f_+ = \frac{1 + v_o}{v} f_0$$

$$\text{Observed frequency (observer receding from source at } v_o) \quad f_- = \frac{1 - v_o}{v} f_0$$

9 Optics

$$\text{Speed of light in a transparent medium} \quad v = \frac{c}{n}$$

$$\text{Snall's Law} \quad n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$\text{Critical angle (total internal reflection)} \quad \theta_c = \sin^{-1} \left(\frac{n_2}{n_1} \right)$$

$$\text{Optical magnification} \quad m = \frac{s'}{s} = \frac{h'}{h}$$

$$\text{Len power} \quad P = \frac{1}{f}$$

$$\text{Thin lens equation} \quad \frac{1}{s} + \frac{1}{s'} = \frac{1}{f}$$

$$\text{Light gathering ability} \quad f - \text{number} = \frac{f}{d}$$

$$\text{Simple magnifier} \quad M = \frac{25 \text{ cm}}{f}$$

10 electric Fields and Forces

Coulomb's law	$F_{1on2} = F_{2on1} = \frac{K q_1 q_2 }{r^2}$
Electric field at point defined by charge q	$E(x,y,z) = \frac{F_{on\ q}(x,y,z)}{q}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$
xxx	$p = \frac{F}{A}$

11 Quantum Numbers

Bohr energy of an hydrogen atom	$E_n = \frac{13.6 \text{ eV}}{n^2}$ where $n = 1, 2, 3, 4, \dots$
Angular momentum of an electron's orbit	$L = \sqrt{\ell(\ell+1)}\hbar$ where $\ell = 0, 1, 2, 3, \dots, n-1$
Magnetic quantum number	$m = -\ell, -\ell+1, \dots, 0, \ell-1, \ell$
Spin quantum number	$m_s = -\frac{1}{2}$ or $+\frac{1}{2}$

12 Nuclear Physics

Half life	$N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{t_{1/2}}}$
Exponential decay	$N = N_0 e^{-\frac{t}{\tau}}$
Activity	$R = \frac{N}{\tau} = \frac{0.693}{t_{1/2}} N$
Binding energy	$B = (Zm_H + Nm_n - m_{atom}) \times (931.49 \text{ MeV/u})$

13 Periodic Tables