

1 **Susceptibility of European freshwater fish to climate change: species profiling based on
2 life-history and environmental characteristics**

3

4 Running head: European freshwater fish and climate change

5

6 Ivan Jarić^{1,2,3*}, Robert J. Lennox⁴, Gregor Kalinkat², Gorčin Cvijanović³ and Johannes
7 Radinger^{2,5}

8

9 ¹ Biology Centre of the Czech Academy of Sciences, Institute of Hydrobiology, Ceske
10 Budejovice, Czech Republic

11 ² Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany

12 ³ Institute for Multidisciplinary Research, University of Belgrade, Serbia

13 ⁴ Fish Ecology and Conservation Physiology Laboratory, Department of Biology, Carleton
14 University, Ottawa, Ontario, Canada

15 ⁵ GRECO, Institute of Aquatic Ecology, University of Girona, Spain

16

17 * Corresponding author: Ivan Jarić, Biology Centre of the Czech Academy of Sciences,
18 Institute of Hydrobiology, Na Sádkách 702/7, 370 05 České Budějovice, Czech Republic, E-
19 mail: ivan.jaric@hbu.cas.cz, phone: +420 38 777 5855, fax: +420 385 310 24

20

21 Keywords: IUCN; Red List; extinction threat; global warming; climate change.

22

23

24

25 **Abstract**

26 Climate change is expected to strongly affect freshwater fish communities. Combined with
27 other anthropogenic impacts, the impacts will alter species distributions and contribute to
28 population declines and local extinctions. To provide timely management and conservation of
29 fishes, it is relevant to identify species that will be most impacted by climate change and those
30 that will be resilient. Species traits are considered a promising source of information on
31 characteristics that influence resilience to various environmental conditions and impacts. We
32 collated life history traits and climatic niches of 443 European freshwater fish species and
33 compared those identified as susceptible to climate change to those that are considered to be
34 resilient. Significant differences were observed between the two groups in their distribution,
35 life-history and climatic niches, with climate-change susceptible species being distributed
36 more southwardly within Europe, and being characterized by higher threat levels, lower
37 commercial relevance, lower vulnerability to fishing, smaller body size and warmer thermal
38 envelopes. We establish a list of species revealed to be of highest priority for further research
39 and monitoring regarding climate change susceptibility within Europe. The presented
40 approach represents a promising tool, to quickly assess large groups of species regarding their
41 susceptibility to climate change and other threats, and to identify research and management
42 priorities.

43

44 **Introduction**

45 As ectothermic organisms, fishes are intimately linked to local climatic conditions
46 through physiological mechanisms that delimit tolerance or resilience (Comte & Olden,
47 2017a). Zoogeography of fishes is therefore greatly influenced by the average and spread of

48 temperatures experienced in a given watershed (Pörtner & Farrell, 2008; Isaak & Rieman,
49 relative to seas and oceans, freshwater habitats are more drastically impacted by
50 changes in climate, especially due to changes in temperature and flow, and climate change is
51 projected to strongly affect freshwater fish communities (O'Reilly et al., 2003; Buisson et al.,
52 2008; Graham & Harrod, 2009; Harrod, 2016; Radinger et al., 2017). Combined with other
53 anthropogenic impacts (e.g. land use change and thermal pollution; Radinger et al., 2016;
54 Raptis et al., 2017), climate change will restrict or redraw thermal envelopes, contribute to
55 population declines and local extinctions, and overall shifts in the distribution of species.
56 Riverine fish species losses due to climate change and reduced water discharge are predicted
57 to reach 75% in some river basins (Xenopoulos et al., 2005). Phenological changes in fish
58 behaviour (Otero et al., 2014; Dempson et al., 2017; Hovel et al., 2017) have been also
59 detected and emphasize the powerful changes imposed by a changing climate. In Europe,
60 there is a broad range of climatic conditions experienced across the landscape and a diverse
61 ichthyofauna distributed throughout the lakes and rivers (Ficke et al., 2007). Within the IUCN
62 (International Union for Conservation of Nature and Natural Resources) Red List, as much as
63 33% of European freshwater fish species are recognized as threatened by climate change
64 (IUCN, 2017).

65 Efforts to preserve ecosystem integrity must focus on maintaining species richness
66 and diversity to ensure that the services provided by freshwater ecosystems are maintained.
67 Conservation is often limited by funding and therefore must undergo triage to identify
68 priorities and allocate resources efficiently (McDonald-Madden et al., 2011). To provide
69 timely management and conservation and allocate resources efficiently, it is important to
70 identify those species that will be most impacted by climate change and those that might be
71 rather resilient. Species traits are considered as a promising source of information on

72 characteristics that influence resilience to various environmental conditions and impacts
73 (Jiguet et al., 2007; Comte & Olden, 2017b). Species traits represent any morphological,
74 physiological or phenological feature that is measurable at the individual level of a species
75 (Floeter et al., 2018). Trait-based evaluation has been demonstrated to be linked to the risk
76 status of species and can be used to investigate mechanisms that contribute to imperilment,
77 make predictions about unassessed species, or rank and prioritize species based on their
78 relative risk (Olden et al., 2007; Bland & Böhm, 2016; Comte & Olden, 2018).

79 Here we assess various ecological and life-history characteristics of European
80 freshwater fish species to identify traits that are characteristic for those that are susceptible to
81 the effects of climate change. Automated scraping of an online trait database and calculation
82 of climate envelopes using IUCN range maps overlaid on climate maps allowed us to collate
83 species-specific data on life history, distribution, climatic niches, as well as data on threat and
84 economic status. This allowed us to compare species identified as susceptible to climate
85 change with those that are considered to be resilient. Results of the study will contribute to a
86 better understanding of the expected climate change effects on European freshwater fish
87 fauna. We also establish a list of European species of highest priority for further research and
88 monitoring regarding climate change susceptibility. The method allows to extrapolate results
89 and characterize rare and less studied species, with scarce autecological information.

90

91 **Materials and methods**

92 *Dataset*

93 Our analysis comprised comparisons of in total 443 European freshwater fishes
94 between those that were identified as threatened by climate change (n=148) within the IUCN
95 Red List Database (IUCN, 2017) and those without climate change listed as a threat (n=295).

96 A list of native European freshwater fish species belonging to 25 families, mainly to
97 Cyprinidae (45%) and Salmonidae (20%), was obtained from the IUCN Red List database
98 (IUCN, 2017). It included both exclusively freshwater species, as well as those that partly
99 enter brackish and saltwater. Obtained data also comprised IUCN Red List classification and
100 maps of their distributional range within Europe. In addition, we obtained information
101 whether climate change was indicated as one of the threats for each species, which is based on
102 threat analyses and expert judgement by IUCN species experts. Overall, the dataset comprised
103 33% species which were categorized as susceptible to climate change.

104 In addition, for each species we collated trait information related to their life
105 history, ecology, fishery and threat status, and spatial and bioclimatic data variables (Table 1).
106 Life-history data were obtained from the FishBase database (Froese & Pauly, 2017) by using
107 the rfishbase R package (Boettiger et al., 2012, 2017). Traits with low data coverage (i.e.
108 those that were available for less than 1% of all species) were excluded from the analysis.
109 Bioclimatic spatial data were obtained from the MERRAclim database (Vega et al., 2017) and
110 included 19 variables related to temperature and humidity (Table 1). Global Human Footprint
111 map (map of anthropogenic impacts on the environment) was obtained from WCS and
112 CIESIN (2005) and the spatial elevation data were obtained from USGS (2010). Based on the
113 distributional range maps of each species (IUCN, 2017), mean values within each species'
114 range were estimated for each of the spatial variables using the intersect tool in ArcGIS
115 (version 10.5) and the *extract* function in the R (version 3.4.3; R Core Team, 2017) package
116 raster (version 2.6-7; Hijmans, 2017). Range maps were also used to estimate the number of
117 watersheds covered by each species based on WRI (2006), as well as the area and coordinates
118 of the range centroid for each species. General descriptive statistics and information on data
119 sources of all variables used in the analysis is presented in Table 1 and Supplementary

120 material S1. The complete dataset is presented in Supplementary material S2.

121

122 *Statistical Analysis*

123 We calculated boosted regression trees (BRT) to evaluate the relationship between the
124 membership of a species to the group of susceptible vs. non-susceptible species and the 45
125 explanatory variables. BRT are a statistical learning method that combines and averages
126 (boosting) many simple single regression trees to form collective models of improved
127 predictive performance (Elith et al., 2008). BRT can accommodate continuous and categorical
128 variables, are not affected by missing values or transformation or outliers and are considered
129 to effectively select relevant variables, identify variable interactions and avoid overfitting
130 (Elith et al., 2008, Radinger et al., 2015).

131 Specifically, we first fitted an initial global BRT model (R package *dismo*, *gbm.step*,
132 version 1.1-4; Hijmans et al., 2017) using the complete set of explanatory variables. An
133 automatized stepwise backward selection of explanatory variables (*gbm.simplify*) was applied
134 to eliminate non-informative variables based on model-internal cross-validation of changes in
135 a models' predictive deviance (Hijmans et al., 2017). Thereafter, we calculated a final BRT
136 model (*gbm.step*) based on the selected set of explanatory variables. For all BRT modeling
137 steps, tree complexity and learning rate was set to 3 and 0.001, respectively, to achieve the
138 recommended number of more than 1000 regression trees (Elith et al., 2008). All other model
139 settings were set to default or were automatically adjusted by the boosting algorithm. We
140 calculated a 10-fold cross validation of the BRT model as already implemented in the
141 algorithm. In addition, we extracted the mean AUC (area under the receiver operating
142 characteristic curve) as a measure of the model's predictive quality. The AUC is a threshold-
143 independent rank-correlation coefficient with high values typically indicating a strong

144 agreement between the model prediction and the membership of species to the susceptible vs.
145 non-susceptible group (Hijmans & Elith, 2017).

146 The relative importance (%) of each explanatory variable in the final BRT model was
147 quantified based on the number of times each variable was used for splitting, weighted by the
148 squared improvement at each split and averaged over all trees (Elith et al., 2008). For BRT
149 models with Gaussian distribution the relative variable importance equals the reduction of
150 squared error attributable to a given variable. Differences between groups were also assessed
151 by bootstrapping, by sampling each group independently and estimating the difference based
152 on confidence intervals (functions *two.boot* and *boot.ci*, R package *simpleboot*, version 1.1-3;
153 Peng 2008). Differences were considered to be significant if 95% confidence intervals did not
154 overlap with zero.

155 Subsequently, species were ranked based on the subset of variables selected by the
156 BRT analysis (i.e., those with >1% variable importance score), and weighed by the
157 importance of each variable, in order to estimate their position along the climate change
158 susceptibility continuum. For each species, the value of each variable was standardized based
159 on its position between the minimum (t_{\min}) and maximum values (t_{\max}) observed in the
160 dataset, with 0-1 possible range, and multiplied by the importance score (I_x) of the given
161 variable:

$$162 \quad R_{tx} = \frac{t_x - t_{\min}}{t_{\max} - t_{\min}} \times I_x \quad (1)$$

163 where R_{tx} represents the rank value of variable t in species x , and t_x is the value of variable t
164 for species x . For variables where the lower endpoint (t_{\min}) is associated with the climate
165 change susceptibility, equation should be adjusted as follows:

$$166 \quad R_{tx} = \frac{t_{\max} - t_x}{t_{\max} - t_{\min}} \times I_x \quad (2)$$

167 Summing of all ranking scores across all variables yielded the total species ranking score,
168 which can range from 0 to 100, with higher values indicating stronger climate change
169 susceptibility.

170

171 **Results**

172 Our analyses indicated substantial differences between the two groups, climate
173 change susceptible versus non-susceptible species. The BRT analysis selected 35 most
174 relevant variables, which were subsequently assessed for their relative importance to
175 discriminate between the two groups (Figure 1 and Supplementary material S3). The BRT
176 model with the selected set of explanatory variables was successfully modeled
177 (Supplementary material S3) with a good cross-validated AUC of 0.87 (standard error =
178 0.014). Out of all explanatory variables, latitude of the species range centroid was selected as
179 by far the most relevant variable (31% variable importance), followed by the IUCN Red List
180 classification (8%), commercial relevance (6%) and vulnerability to fishing (6%). Climate
181 susceptible species were characterized by more southwardly positioned distribution range
182 centroids (41.6° vs. 47.8° N as a mean value in susceptible and non-susceptible species,
183 respectively), smaller range sizes ($175 \times 10^3 \text{ km}^2$ vs. $1686 \times 10^3 \text{ km}^2$), and lower elevations
184 within their ranges (717.7 m vs. 892.2 m a.s.l.), with a higher proportion of exclusively
185 freshwater species (93% vs. 66%; Figure 2). Susceptible species were also characterized by a
186 smaller maximum body length (23.4 cm vs. 41.0 cm), higher proportion of threatened species
187 (63% vs. 31%), lower proportion of commercially relevant species (25% vs. 74% of highly
188 commercial, commercial and minor commercial species), and lower vulnerability to
189 overfishing (32.6 vs. 38.5 vulnerability index; Figure 3), as well as by higher temperature-
190 related values (Supplementary material S4). Bootstrapping indicated significant differences

191 between the groups in each of the variables. Species that are susceptible to climate change are
192 mainly distributed within the Mediterranean region, while the non-susceptible species
193 distribution mainly covers central and northern European regions, as well as the Carpathian
194 region (Figure 4).

195 Species ranking based on the association of their traits with climate change
196 susceptibility characteristics is presented in Table 2 and Supplementary material S5. The five
197 top-ranked climate susceptible species were Acheron spring goby (*Knipowitschia milleri*),
198 Corfu toothcarp (*Valencia letourneuxi*), *Iberochondrostoma almacai*, Evia barbel (*Barbus*
199 *euboicus*) and Malaga chub (*Squalius malacitanus*). Most of the species with the high climate-
200 susceptibility ranks are also classified as highly threatened according to the IUCN
201 classification (Table 2). Interestingly, the highest ranked species, *K. milleri*, was not classified
202 within the IUCN Red List as threatened by climate change. Other high-ranking species that
203 were not recognized as threatened by climate change were Malaga chub (*Squalius*
204 *malacitanus*), Almiri toothcarp (*Aphanius almiriensis*), and Trichonis dwarf goby
205 (*Economidichthys trichonis*). Species with the lowest ranking scores, i.e. with low climate
206 change susceptibility, were humpback whitefish (*Coregonus pidschian*), Arctic flounder
207 (*Liopsetta glacialis*), northern pike (*Esox lucius*), burbot (*Lota lota*), and European perch
208 (*Perca fluviatilis*). A complete list of all species' rankings is presented in Supplementary
209 material S5.

210

211 **Discussion**

212 In the present study, significant differences in life-history and climatic niches were
213 observed between the European freshwater species susceptible to climate change and those
214 that are not, such as species body size, range size, distribution and thermal envelopes. The

215 latitude of the species range centroid was by far the most influential trait. Overall, southern
216 regions with the warmer, Mediterranean climate comprised a higher proportion of species
217 susceptible to climate change (Figure 4). These results support recent findings that the species
218 from lower latitudes and tropical, warm-water habitats, are at greater risk from climate change
219 and warming (Payne & Smith, 2016; Payne et al., 2016; Comte & Olden, 2017b). In such
220 species, evolved towards higher upper thermal tolerances, specialization to thermal extremes
221 is accompanied by a reduced physiological flexibility and adaptation capacity to respond to
222 changing environmental conditions (Payne & Smith, 2016; Payne et al., 2016; Comte &
223 Olden, 2017b). Such heat-tolerant species are also adapted to temperatures close to their
224 physiological thermal limits, with a narrow safety margin for further temperature increases
225 (Sinclair et al., 2016; Comte & Olden, 2017b). Freshwater basins in Southern Europe are also
226 of particular conservation concern due to an elevated pressure by a range of anthropogenic
227 impacts that further exacerbate effects of climate change, such as pollution, water resource
228 development and consumption, and biological invasions (Xenopoulos et al., 2005; Clavero &
229 García-Berthou, 2005; Walther et al., 2009; Vörösmarty et al., 2010; Comte & Olden, 2017a).

230 Climate-susceptible species were also characterized by a smaller body and range
231 size (Figures 2, 3). These traits, which are also related to a lower dispersal ability (Radinger &
232 Wolter, 2014), are well recognized as predictors of climate change susceptibility in fish (e.g.
233 Ficke et al., 2007; Isaak & Rieman, 2013; Chessman, 2013; Pearson et al., 2014; Radinger et
234 al., 2017). Smaller-bodied fish are facing elevated overall extinction risk in freshwater habitat
235 due to multiple threats, such as habitat loss and fragmentation (Olden et al., 2007; Kalinkat et
236 al., 2017; Kopf et al., 2017), which explains higher threat level observed in climate-
237 susceptible species in the present study. Observed lower commercial relevance and lower
238 vulnerability to fishing of climate-susceptible species both stem from a lower body size and

239 related faster life history of such species.

240 It is important to acknowledge certain limitations of the data sources used in this
241 study, such as species and trait coverage, reliability of methods applied for threat and
242 extinction risk classification, and potential assessors' biases (Clavero & García-Berthou, 2005;
243 Keith et al., 2014; Trull et al., 2018). Furthermore, species that are not classified within IUCN
244 Red List as threatened by climate change can comprise also those that are not yet assessed for
245 their major threats. Nevertheless, the focus of our study on a well-studied group such as
246 European species ensured that the basic life history data and IUCN Red List assessments were
247 mostly available (Kopf et al., 2017). IUCN Red List is sometimes considered to underestimate or
248 improperly account for climate change as a threat, mostly due to ambiguous definitions and
249 criteria (Trull et al., 2018). However, recent studies have indicated that the IUCN
250 classification is more efficient in detecting species vulnerable to climate change than
251 anticipated (Keith et al., 2014; Pearson et al., 2014). Notwithstanding all the caveats, the
252 databases used in the present study still represent the most comprehensive sources of data and
253 the best available knowledge (Olden et al., 2007; Vega et al., 2017).

254 Trait-based risk assessments are increasingly used for species profiling (Pacifici et
255 al., 2015; Liu et al., 2017; MacLean & Beissinger, 2017). The approach presented in this study
256 might be considered a valid and promising approach to be used as a screening tool, i.e. to
257 quickly assess large groups of species regarding their susceptibility to climate change and
258 other threats based on species traits, and to identify research and management priorities. Our
259 results indicate that the European environmental policy related to climate change mitigation
260 and adaptation (EEA, 2012, 2017) should be mainly focused on Mediterranean region. This is
261 especially important since this region is also predicted to have the highest frequency of
262 droughts and extreme high temperatures, strongest reduction in precipitation and river

263 discharges, the highest aggregate climate change impact and the lowest adaptation capacity
264 (Milly et al., 2005; Dankers & Feyen, 2008; Fischer & Schär, 2010; ESPON Climate, 2011;
265 Stagge et al., 2011; Rojas et al., 2012; Jacob et al., 2014; Russo et al., 2014). Moreover,
266 Mediterranean region was also identified as a European priority area regarding climate change
267 impacts for other species groups. A similar distributional pattern of species susceptible to
268 climate change was previously also reported for aquatic insects such as Plecoptera,
269 Ephemeroptera and Trichoptera (Hering et al., 2009; Conti et al., 2014), mammals (Levinsky
270 et al., 2007), as well as for terrestrial species in general (Pacifici et al., 2015).

271 Species ranking conducted here indicated priority species for further research and
272 monitoring regarding climate change (e.g. *V. letourneuxi*, *I. almacai* and *B. euboicus*; Table 2).
273 Moreover, it also identified species whose IUCN Red List status potentially needs to be
274 reconsidered or updated, such as highly ranked but apparently non-susceptible species (e.g. *K.
275 milleri*), or highly ranked species without a proper threat category (e.g. *K. goernerii*, classified
276 as Data Deficient species). As such, it has a potential to be used as a “Robin Hood Approach”
277 (Punt et al., 2011), where assessments based on information-rich species are used to evaluate
278 and categorize those that are information-poor. There is a need for climate change
279 vulnerability assessments that would be based on quantitative approaches and consistent set of
280 criteria, such as trait-based approaches advocated by IUCN (Foden et al., 2013; Trull et al.,
281 2018). The approach presented here could be a good step in that direction.

282

283 **Acknowledgements**

284 IJ acknowledges the sponsorship provided by the The J. E. Purkyně Fellowship of the
285 Academy of Sciences of the Czech Republic and funding from the European Union’s Horizon
286 2020 research and innovation programme through the project ClimeFish (grant No. 677039),

287 as well as funding provided by the Alexander von Humboldt Foundation and the German
288 Federal Ministry of Education and Research (BMBF). GK acknowledges funding from the
289 BMBF through the project “GLANCE” (Global Change Effects in River Ecosystems; 01
290 LN1320A). JR acknowledges funding through the 2015-2016 BiodivERsA COFUND call for
291 research proposals and the Spanish Ministry of Economy, Industry and Competitiveness
292 (project ODYSSEUS, BiodivERsA3-2015-26, PCIN-2016-168). GC acknowledges funding
293 from the Ministry of Education, Science and Technological Development, Republic of Serbia
294 (grant number 173045). Thanks are extended to Ryder Burt for assistance with geographic
295 information processing.

296

297 **References**

- 298 Bland, L. M., & Böhm, M. (2016). Overcoming data deficiency in reptiles. *Biological
299 Conservation*, 204, 16-22.
- 300
- 301 Boettiger, C., Lang, D. T., & Wainwright, P. (2012). rfishbase: exploring, manipulating and
302 visualizing FishBase data from R. *Journal of Fish Biology*, 81, 2030–2039.
- 303
- 304 Boettiger, C., Chamberlain, S., Lang, D. T., & Wainwright, P. (2017). Package ‘rfishbase’. R
305 Interface to 'FishBase'. Version 2.1.2. <https://github.com/ropensci/rfishbase>. Accessed 15
306 January 2018
- 307
- 308 Buisson, L., Thuiller, W., Lek, S., Lim, P., & Grenouillet, G. (2008). Climate change hastens
309 the turnover of stream fish assemblages. *Global Change Biology*, 14, 2232–2248.
- 310

311 Chessman, B. C. (2013). Identifying species at risk from climate change: traits predict the
312 drought vulnerability of freshwater fishes. *Biological Conservation*, 160, 40-49.

313

314 Clavero, M., & García-Berthou, E. (2005). Invasive species are a leading cause of animal
315 extinctions. *Trends in Ecology & Evolution*, 20, 110.

316

317 Comte, L., & Olden, J. D. (2017a). Climatic vulnerability of the world's freshwater and
318 marine fishes. *Nature Climate Change*, 7, 718.

319

320 Comte, L., & Olden, J. D. (2017b). Evolutionary and environmental determinants of
321 freshwater fish thermal tolerance and plasticity. *Global Change Biology*, 23, 728-736.

322

323 Comte, L., & Olden, J. D. (2018). Evidence for dispersal syndromes in freshwater fishes.
324 *Proceedings of the Royal Society B*, 285, 20172214.

325

326 Conti, L., Schmidt-Kloiber, A., Grenouillet, G., & Graf, W. (2014). A trait-based approach to
327 assess the vulnerability of European aquatic insects to climate change. *Hydrobiologia*, 721,
328 297-315.

329

330 Dankers, R., & Feyen, L. (2008). Climate change impact on flood hazard in Europe: An
331 assessment based on high resolution climate simulations. *Journal of Geophysical Research*,
332 113, D19105.

333

334 Dempson, B., Schwarz, C. J., Bradbury, I. R., Robertson, M. J., Veinott, G., Poole, R., &

- 335 Colbourne, E. (2017). Influence of climate and abundance on migration timing of adult
336 Atlantic salmon (*Salmo salar*) among rivers in Newfoundland and Labrador. *Ecology of*
337 *Freshwater Fish*, 26, 247-259.
- 338
- 339 EEA (2012). *Climate Change, Impacts and Vulnerability in Europe 2012: An Indicator-Based*
340 *Report*. EEA Report No 12/2012. Copenhagen: European Environment Agency.
- 341
- 342 EEA (2017). *Climate Change, Impacts and Vulnerability in Europe 2016: An Indicator-Based*
343 *Report*. EEA Report No 1/2017. Copenhagen: European Environment Agency.
- 344
- 345 Elith, J., Leathwick, J. R., & Hastie, T. (2008). A working guide to boosted regression trees.
346 *Journal of Animal Ecology*, 77, 802–813.
- 347
- 348 ESPON Climate (2011). Climate Change and Territorial Effects on Regions and Local
349 Economies (Scientific Report). Institute of Spatial Planning (IRPUD), TU Dortmund
350 University. http://www.espon.eu/main/Menu_Projects/Menu_AppliedResearch/climate.html.
351 Accessed 20 June 2018
- 352
- 353 Ficke, A. D., Myrick, C. A., & Hansen, L. J. (2007). Potential impacts of global climate
354 change on freshwater fisheries. *Reviews in Fish Biology and Fisheries*, 17, 581-613.
- 355
- 356 Fischer, E. M., & Schär, C. (2010). Consistent geographical patterns of changes in high-
357 impact European heatwaves. *Nature Geoscience*, 3, 398-403.
- 358

- 359 Floeter, S. R., Bender, M. G., Siqueira, A. C., & Cowman, P. F. (2018). Phylogenetic
360 perspectives on reef fish functional traits. *Biological Reviews*, 93, 131-151.
361
362 Foden, W. B., Butchart, S. H., Stuart, S. N., Vié, J. C., Akçakaya, H. R., Angulo, A., ... &
363 Donner, S. D. (2013). Identifying the world's most climate change vulnerable species: a
364 systematic trait-based assessment of all birds, amphibians and corals. *PloS One*, 8, e65427.
365
366 Froese, R., & Pauly, D. (Eds.) (2017). FishBase. World Wide Web electronic publication.
367 version 10/2017. <https://www.fishbase.org>. Accessed 15 January 2018
368
369 Graham, C. T., & Harrod, C. (2009). Implications of climate change for the fishes of the
370 British Isles. *Journal of Fish Biology*, 74, 1143-1205.
371
372 Harrod, C. (2016). Climate change and freshwater fisheries. In J. F. Craig (Ed.), *Freshwater
373 Fisheries Ecology* (pp. 641-694). John Wiley & Sons.
374
375 Hering, D., Schmidt-Kloiber, A., Murphy, J., Lücke, S., Zamora-Munoz, C., López-
376 Rodríguez, M. J., ... & Graf, W. (2009). Potential impact of climate change on aquatic insects:
377 a sensitivity analysis for European caddisflies (Trichoptera) based on distribution patterns and
378 ecological preferences. *Aquatic Sciences*, 71, 3-14.
379
380 Hijmans, R. J. (2017). Raster: Geographic Data Analysis and Modeling. R package version
381 2.6-7. <https://CRAN.R-project.org/package=raster>. Accessed 20 June 2018
382

- 383 Hijmans, R. J., & Elith, J. (2017). Species distribution modeling with R. R vignette.
- 384 <https://cran.r-project.org/web/packages/dismo/vignettes/sdm.pdf>. Accessed 20 June 2018
- 385
- 386 Hijmans, R. J., Phillips, S., Leathwick, J., & Elith, J. (2017). Dismo: Species distribution
- 387 modeling. <http://cran.r-project.org/package=dismo>. Accessed 20 June 2018
- 388
- 389 Hovel, R. A., Carlson, S. M., & Quinn, T. P. (2017). Climate change alters the reproductive
- 390 phenology and investment of a lacustrine fish, the three-spine stickleback. *Global Change*
- 391 *Biology*, 23, 2308-2320.
- 392
- 393 Isaak, D. J., & Rieman, B. E. (2013). Stream isotherm shifts from climate change and
- 394 implications for distributions of ectothermic organisms. *Global Change Biology*, 19, 742-751.
- 395
- 396 IUCN (2017). The IUCN Red List of Threatened Species. Version 2017-1.
- 397 <https://www.iucnredlist.org>. Accessed 15 January 2018
- 398
- 399 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., ... &
- 400 Georgopoulou, E. (2014). EURO-CORDEX: new high-resolution climate change projections
- 401 for European impact research. *Regional Environmental Change*, 14, 563-578.
- 402
- 403 Jiguet, F., Gadot, A. S., Julliard, R., Newson, S. E., & Couvet, D. (2007). Climate envelope,
- 404 life history traits and the resilience of birds facing global change. *Global Change Biology*, 13,
- 405 1672-1684.
- 406

- 407 Kalinkat, G., Jähnig, S. C., & Jeschke, J. M. (2017). Exceptional body size–extinction risk
408 relations shed new light on the freshwater biodiversity crisis. *Proceedings of the National
409 Academy of Sciences*, 201717087.
- 410
- 411 Keith, D. A., Mahony, M., Hines, H., Elith, J., Regan, T. J., Baumgartner, J. B., ... & Penman,
412 T. (2014). Detecting extinction risk from climate change by IUCN Red List criteria.
413 *Conservation Biology*, 28, 810-819.
- 414
- 415 Kopf, R. K., Shaw, C., & Humphries, P. (2017). Trait based prediction of extinction risk of
416 small bodied freshwater fishes. *Conservation Biology*, 31, 581-591.
- 417
- 418 Levinsky, I., Skov, F., Svenning, J. C., & Rahbek, C. (2007). Potential impacts of climate
419 change on the distributions and diversity patterns of European mammals. *Biodiversity and
420 Conservation*, 16, 3803-3816.
- 421
- 422 Liu, C., Comte, L., & Olden, J. D. (2017). Heads you win, tails you lose: Life history traits
423 predict invasion and extinction risk of the world's freshwater fishes. *Aquatic Conservation:
424 Marine and Freshwater Ecosystems*, 27, 773-779.
- 425
- 426 MacLean, S. A., & Beissinger, S. R. (2017). Species' traits as predictors of range shifts under
427 contemporary climate change: a review and meta analysis. *Global Change Biology*, 23,
428 4094-4105.
- 429
- 430 McDonald-Madden, E., Chadès, I., McCarthy, M. A., Linkie, M., & Possingham, H. P. (2011).

431 Allocating conservation resources between areas where persistence of a species is uncertain.

432 *Ecological Applications*, 21, 844-858.

433

434 Milly, P. C., Dunne, K. A., & Vecchia, A. V. (2005). Global pattern of trends in streamflow

435 and water availability in a changing climate. *Nature*, 438, 347-350.

436

437 Olden, J. D., Hogan, Z. S., & Zandeen, M. J. V. (2007). Small fish, big fish, red fish, blue fish:

438 size-biased extinction risk of the world's freshwater and marine fishes. *Global Ecology and*

439 *Biogeography*, 16, 694-701.

440

441 O'Reilly, C. M., Alin, S. R., Plisnier, P. D., Cohen, A. S., & McKee, B. A. (2003). Climate

442 change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature*, 424,

443 766-768.

444

445 Otero, J., L'Abée Lund, J. H., Castro-Santos, T., Leonardsson, K., Storvik, G. O., Jonsson,

446 B., ... & Dionne, M. (2014). Basin scale phenology and effects of climate variability on

447 global timing of initial seaward migration of Atlantic salmon (*Salmo salar*). *Global Change*

448 *Biology*, 20, 61-75.

449

450 Pacifici, M., Foden, W. B., Visconti, P., Watson, J. E., Butchart, S. H., Kovacs, K. M., ... &

451 Corlett, R. T. (2015). Assessing species vulnerability to climate change. *Nature Climate*

452 *Change*, 5, 215-225.

453

454 Payne, N. L., & Smith, J. A. (2017). An alternative explanation for global trends in thermal

- 455 tolerance. *Ecology Letters*, 20, 70-77.
- 456
- 457 Payne, N. L., Smith, J. A., Meulen, D. E., Taylor, M. D., Watanabe, Y. Y., Takahashi, A., ... &
- 458 Suthers, I. M. (2016). Temperature dependence of fish performance in the wild: links with
- 459 species biogeography and physiological thermal tolerance. *Functional Ecology*, 30, 903-912.
- 460
- 461 Pearson, R. G., Stanton, J. C., Shoemaker, K. T., Aiello-Lammens, M. E., Ersts, P. J., Horning,
- 462 N., ... & Akçakaya, H. R. (2014). Life history and spatial traits predict extinction risk due to
- 463 climate change. *Nature Climate Change*, 4, 217-221.
- 464
- 465 Peng, R. D. (2008). Simple bootstrap routines. <https://cran.r-project.org/package=simpleboot>.
- 466 Accessed 20 June 2018
- 467
- 468 Pörtner, H. O., & Farrell, A. P. (2008). Physiology and climate change. *Science*, 322, 690-692.
- 469
- 470 Punt, A. E., Smith, D. C., & Smith, A. D. (2011). Among-stock comparisons for improving
- 471 stock assessments of data-poor stocks: the “Robin Hood” approach. *ICES Journal of Marine*
- 472 *Science*, 68, 972-981.
- 473
- 474 R Development Core Team (2017). R: A language and environment for statistical computing.
- 475 R Foundation for Statistical Computing, Vienna, Austria.
- 476
- 477 Radinger, J., & Wolter, C. (2014). Patterns and predictors of fish dispersal in rivers. *Fish and*
- 478 *Fisheries*, 15, 456-473.

479

480 Radinger, J., Wolter, C., & Kail, J. (2015). Spatial Scaling of Environmental Variables
481 Improves Species-Habitat Models of Fishes in a Small, Sand-Bed Lowland River. *PLoS One*,
482 10, e0142813.

483

484 Radinger, J., Hölker, F., Horký, P., Slavík, O., Dendoncker, N., & Wolter, C. (2016).
485 Synergistic and antagonistic interactions of future land use and climate change on river fish
486 assemblages. *Global Change Biology*, 22, 1505-1522.

487

488 Radinger, J., Essl, F., Hölker, F., Horký, P., Slavík, O., & Wolter, C. (2017). The future
489 distribution of river fish: the complex interplay of climate and land use changes, species
490 dispersal and movement barriers. *Global Change Biology*, 23, 4970-4986.

491

492 Raptis, C. E., Boucher, J. M., & Pfister, S. (2017). Assessing the environmental impacts of
493 freshwater thermal pollution from global power generation in LCA. *Science of the Total
494 Environment*, 580, 1014-1026.

495

496 Rojas, R., Feyen, L., Bianchi, A., & Dosio, A. (2012). Assessment of future flood hazard in
497 Europe using a large ensemble of bias corrected regional climate simulations. *Journal of
498 Geophysical Research*, 117, D17109.

499

500 Russo, S., Dosio, A., Graversen, R. G., Sillmann, J., Carrao, H., Dunbar, M. B., ... & Vogt, J.
501 V. (2014). Magnitude of extreme heat waves in present climate and their projection in a
502 warming world. *Journal of Geophysical Research: Atmospheres*, 119, 12500-12512.

- 503
- 504 Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., ... &
- 505 Huey, R. B. (2016). Can we predict ectotherm responses to climate change using thermal
- 506 performance curves and body temperatures?. *Ecology Letters*, 19, 1372-1385.
- 507
- 508 Stagge, J. H., Rizzi, J., Tallaksen, L. M., & Stahl, K. (2015). Future meteorological drought:
- 509 projections of regional climate models for Europe. In *EGU General Assembly Conference*
- 510 *Abstracts* (Vol. 17). Technical Report No. 25, DROUGHT-R&SPI, European Commission.
- 511
- 512 Trull, N., Böhm, M., & Carr, J. (2018). Patterns and biases of climate change threats in the
- 513 IUCN Red List. *Conservation Biology*, 32, 135-147.
- 514
- 515 USGS (2010). Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010). U.S.
- 516 Geological Survey (USGS). https://topotools.cr.usgs.gov/gmted_viewer. Accessed 15 January
- 517 2018
- 518
- 519 Vega, G. C., Perttierra, L. R., & Olalla-Tárraga, M. Á. (2017). MERRAclim, a high-resolution
- 520 global dataset of remotely sensed bioclimatic variables for ecological modelling. *Scientific*
- 521 *Data*, 4, 170078.
- 522
- 523 Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., ...
- 524 & Davies, P. M. (2010). Global threats to human water security and river biodiversity. *Nature*,
- 525 467, 555-561.
- 526

- 527 Walther, G. R., Roques, A., Hulme, P. E., Sykes, M. T., Pyšek, P., Kühn, I., ... & Czucz, B.
- 528 (2009). Alien species in a warmer world: risks and opportunities. *Trends in Ecology and*
529 *Evolution*, 24, 686-693.
- 530
- 531 WCS, & CIESIN (2005). Last of the Wild Project, Version 2, 2005 (LWP-2): Global Human
- 532 Footprint Dataset (IGHP). Wildlife Conservation Society (WCS) and and Center for
- 533 International Earth Science Information Network (CIESIN), Columbia University. Palisades,
- 534 NY: NASA Socioeconomic Data and Applications Center (SEDAC).
- 535 <http://dx.doi.org/10.7927/H4GF0RFQ>. Accessed 15 January 2018
- 536
- 537 WRI (2006). World Resources Institute (WRI) Major Watersheds of the World Delineation.
- 538 FAO GeoNetwork, FAO - Aquaculture Management and Conservation Service (FIMA).
- 539 <http://www.fao.org/geonetwork/srv/en/metadata.show?id=30914>. Accessed 15 January 2018
- 540
- 541 Xenopoulos, M. A., Lodge, D. M., Alcamo, J., Märker, M., Schulze, K., & Van Vuuren, D. P.
- 542 (2005). Scenarios of freshwater fish extinctions from climate change and water withdrawal.
- 543 *Global Change Biology*, 11, 1557-1564.
- 544
- 545
- 546
- 547
- 548
- 549
- 550

Table 1. Variables used in the analysis, with their data sources, general descriptive statistics and coverage (proportion of species with available data). See Supplementary material I for more information.

Variable name	Data source	Median; mean ± SD (range); proportions (%) for categorical data	Coverage (%)
1 Game fish	FishBase	Yes: 14, No: 86	100
2 IUCN Red List status	FishBase/IUCN	EX: 3, CR: 12, EN: 11, VU: 16, NT: 4, LC: 48, DD: 6	100
3 Climate zone (Köppen climate classification)	FishBase	Subtropical: 26, temperate 73, polar: 1	100
4 Preferred habitat	FishBase	Pelagic: 3, pelagic-neritic: 3, benthopelagic: 64, demersal: 30	100
5 Minimum value of the water depth range (m)	FishBase	2; 11.2 ± 17.9 (0 - 100)	18
6 Maximum value of the water depth range (m)	FishBase	80; 106.7 ± 138.8 (1 - 700)	14
7 Freshwater preference	FishBase	Exclusively freshwater: 75, enters saltwater: 25	100
8 Maximum recorded body length (cm)	FishBase	21.7; 35.2 ± 60.5 (2.2 - 800)	98
9 Lateral body shape type	FishBase	Eell-like: 1, elongated: 38, fusiform/normal: 59, short and/or deep: 1	77
10 Aspect ratio of the caudal fin	FishBase	1.6; 1.6 ± 0.6 (0.4 - 3.4)	34
11 Trophic level	FishBase	3.3; 3.4 ± 0.6 (2.1 - 4.5)	13
12 Batch spawner	FishBase	Yes: 12, no: 88	55
13 Reproductive guild (first classification)	FishBase	Bearers: 1, guarders: 18, nonguarders: 81	53
14 Reproductive guild (second classification)	FishBase	Brood hiders: 10, clutch tenders: 5, external brooders: 10, internal live bearers: 2, nesters: 18, open water/substratum egg scatterers: 57	28
15 Maximum recorded longevity (years)	FishBase	9; 13.8 ± 17.4 (1 - 118)	30
16 Commercial importance	FishBase	Of no interest: 41, subsistence fisheries: 4, minor commercial: 15, commercial: 35, highly commercial: 5	40
17 Average global landings/production	FishBase	828.4; 10461.4 ± 23020.9 (0 - 104902.8)	12
18 Resilience to fishing pressure	FishBase	1: 3, 2: 12, 3: 58, 4: 27	100
19 Vulnerability to fishing	FishBase	32.9; 36.5 ± 16.6 (10 - 88.7)	100
20 Temperature tolerance (max - min reported T, °C)	FishBase	15; 14.4 ± 5.2 (1 - 32)	21
21 Number of inhabited freshwater basins	WRI (2006)/IUCN(range)	2; 8.5 ± 14.6 (0 - 82)	96
22 Global Human Footprint	WCS & CIESIN (2005)/IUCN(range)	6; 7.8 ± 7.8 (0 - 46)	96
23 Longitude of the centroid of species range	IUCN(range)	18.6; 17.2 ± 17.4 (-81.1 - 117.2)	96

24	Latitude of the centroid of species range	IUCN(range)	44.6; 45.8 ± 6.3 (34.4 - 70.1)	96
25	Range size (km ²)	IUCN(range)	32499; 1180481 ± 3799809 (13 - 35987250)	96
26	Mean elevation within the species range (m)	USGS (2010)/IUCN(range)	590.4; 833.8 ± 652.5 (3.9 - 2373.6)	96
27	Annual mean temperature (°C)	MERRAclim/IUCN(range)	13.5; 13.4 ± 4.1 (-3.7 - 22.7)	96
28	Mean diurnal temperature range (°C)	MERRAclim/IUCN(range)	19.9; 19.5 ± 2.9 (7.2 - 27.0)	96
29	Temperature isothermality (°C)	MERRAclim/IUCN(range)	44.9; 45.3 ± 3.9 (34.5 - 55.0)	96
30	Temperature seasonality (st. dev. x 100) (°C)	MERRAclim/IUCN(range)	780.6; 800.0 ± 180.6 (312.6 - 1515.3)	96
31	Maximum temperature of the warmest month (°C)	MERRAclim/IUCN(range)	35.7; 35.0 ± 4.6 (20.3 - 43.2)	96
32	Minimum temperature of the coldest month (°C)	MERRAclim/IUCN(range)	-8.6; -8.5 ± 7.6 (-36.4 - 8.8)	96
33	Temperature annual range (°C)	MERRAclim/IUCN(range)	43.3; 43.5 ± 7.5 (17.1 - 65.9)	96
34	Mean temperature of the most humid quarter (°C)	MERRAclim/IUCN(range)	22.9; 22.7 ± 3.5 (10.8 - 30.5)	96
35	Mean temperature of the least humid quarter (°C)	MERRAclim/IUCN(range)	4.2; 3.9 ± 5.8 (-19.8 - 16.5)	96
36	Mean temperature of the warmest quarter (°C)	MERRAclim/IUCN(range)	23.2; 23.1 ± 3.7 (11.0 - 31.0)	96
37	Mean temperature of the coldest quarter (°C)	MERRAclim/IUCN(range)	3.9; 3.6 ± 5.7 (-19.9 - 15.6)	96
38	Annual mean specific humidity (g of water / kg of air)	MERRAclim/IUCN(range)	7.2; 7.2 ± 0.9 (3.5 - 10.1)	96
39	Specific humidity of the most humid month (g of water / kg of air)	MERRAclim/IUCN(range)	11.2; 11.1 ± 1.0 (6.7 - 14.3)	96
40	Specific humidity of the least humid month (g water / kg air)	MERRAclim/IUCN(range)	4.0; 4.0 ± 1.1 (0.8 - 7.1)	96
41	Specific humidity seasonality (g water / kg air)	MERRAclim/IUCN(range)	255.4; 249.7 ± 52.3 (123.0 - 395.5)	96
42	Specific humidity of the most humid quarter (g water / kg air)	MERRAclim/IUCN(range)	10.4; 10.4 ± 1.0 (6.2 - 13.6)	96
43	Specific humidity of the least humid quarter (g water / kg air)	MERRAclim/IUCN(range)	4.5; 4.4 ± 1.1 (1.0 - 7.5)	96
44	Specific humidity of the warmest quarter (g water / kg air)	MERRAclim/IUCN(range)	10.3; 10.3 ± 1.0 (6.2 - 13.3)	96
45	Specific humidity of the coldest quarter (g water / kg air)	MERRAclim/IUCN(range)	4.5; 4.4 ± 1.2 (1.0 - 7.6)	96

Table 2. European freshwater fish species with the highest ranking scores, estimated based on the association of their traits with the climate change susceptibility characteristics, as indicated by the BRT model. Complete ranking list of all species is presented in Supplementary material 5.

Rank	Species	Climate change susceptibility	IUCN Red List category	Ranking score
1	<i>Knipowitschia milleri</i>	non-susceptible	CR	82.5
2	<i>Valencia letourneuxi</i>	susceptible	CR	82.1
3	<i>Iberochondrostoma almacai</i>	susceptible	CR	81.6
4	<i>Barbus euboicus</i>	susceptible	CR	81.4
5	<i>Squalius malacitanus</i>	non-susceptible	EN	80.9
6	<i>Aphanius baeticus</i>	susceptible	EN	80.7
7	<i>Knipowitschia goernerri</i>	susceptible	DD	80.5
8	<i>Squalius keadicus</i>	susceptible	EN	80.5
9	<i>Pelasgus laconicus</i>	susceptible	CR	80.4
10	<i>Tropidophoxinellus spartiaticus</i>	susceptible	VU	80.3
11	<i>Iberocypris palaciosi</i>	susceptible	CR	80.1
12	<i>Pelasgus epiroticus</i>	susceptible	CR	80.0
13	<i>Aphanius almiriensis</i>	non-susceptible	CR	79.8
14	<i>Anaecypris hispanica</i>	susceptible	EN	79.5
15	<i>Salaria economidisi</i>	susceptible	CR	79.5
16	<i>Squalius torgalensis</i>	susceptible	EN	79.4
17	<i>Cobitis trichonica</i>	susceptible	EN	78.9
18	<i>Valencia hispanica</i>	susceptible	CR	78.6
19	<i>Economidichthys trichonis</i>	non-susceptible	EN	78.6
20	<i>Knipowitschia thessala</i>	susceptible	EN	78.5

1 **Figure captions:**

2

3 **Figure 1.** Variables selected by the boosted regression tree (BRT) model as the most relevant
4 descriptors of climate change susceptibility in European freshwater fish species; 20 most
5 relevant variables are presented, which together account for 90% of the total relative variable
6 influence.

7

8 **Figure 2.** Violin-boxplots and barplots of the most relevant spatial variables in European
9 freshwater fish species indicated as either susceptible ($n = 148$) or non-susceptible ($n = 295$)
10 to climate change. Habitat preference: blue - exclusively freshwater species, green - species
11 that also enter saltwater.


12


13 **Figure 3.** Violin-boxplots and barplots of the most relevant life history traits and variables
14 related to threat and commercial status in European freshwater fish species indicated as either
15 susceptible ($n = 148$) or non-susceptible ($n = 295$) to climate change.







16


17 **Figure 4.** Richness of freshwater fish species across Europe indicated as either susceptible
18 (middle panel) or non-susceptible (left panel) to climate change, and the relative share of
19 susceptible species in the local total species richness (right panel).

20











Number of non-susceptible species

≤5	25	50	100	≥150
----	----	----	-----	------



Number of susceptible species

≤2	5	10	20	≥30
----	---	----	----	-----



Share (%) of susceptible species

0	25	50	75	100
---	----	----	----	-----