

1 Phylogenetically-informed crayfish conservation in the face of climate 2 change

3

4 Short title: Climate change threatens crayfish evolutionary history

5

6 Key words: crayfish, phylogenetic diversity, evolutionary history, conservation, prioritisation, EDGE
7 species, climate change.

8

9 Sebastian Pipins ^{a, b, c, d, e, f*}, Monika Böhm ^{g, h, i}, Lucie Bland ^{j, k}, Keith A. Crandall ^{k, l, m, n}, Md Anwar
10 Hossain ^o, James Rosindell ^{a, f}, Rikki Gumbs ^{b, f, g}

11

12 ^a Department of Life Sciences, Imperial College London, Silwood Park Campus, Berkshire, UK

13 ^b EDGE of Existence Programme, Zoological Society of London, London, UK

14 ^c Royal Botanic Gardens, Kew, UK.

15 ^d On the Edge, London, UK.

16 ^e Science and Solutions for a Changing Planet DTP, Grantham Institute, Imperial College London,
17 London, UK

18 ^f IUCN SSC Phylogenetic Diversity Task Force, London, UK

19 ^g Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY, UK

20 ^h Global Center for Species Survival, Indianapolis Zoo, 1200 West Washington Street, Indianapolis IN
21 46222, USA

22 ⁱIUCN SSC Freshwater Conservation Committee, IUCN Species Survival Commission, Rue Mauverney
23 28, 1196 Gland, Switzerland

24 ^jCentre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University,
25 Burwood, Victoria, Australia

26 ^kComputational Biology Institute, George Washington University, Washington DC, 20052, USA

27 ^lDepartment of Invertebrate Zoology, Smithsonian Institution, Washington, DC, 20560, USA

28 ^mBiodiverse Anthropocenes, University of Oulu, Oulu, FI-00014, Finland

29 ⁿIUCN SSC Freshwater Crustacean Specialist Group, Gland, Switzerland

30 ^oSchool of BioSciences, The University of Melbourne, Parkville, Vic. 3010, Australia

31

32

33 * Corresponding author.

34

35 E-mail address: sebastian.pipins19@imperial.ac.uk (S. Pipins).

36

37 **Abstract**

38

39 Crayfish are an ancient clade of freshwater decapods that play vital and diverse ecological roles in the
40 freshwater systems that they inhabit. One third of assessed crayfish species are threatened with extinction
41 and 87% are highly sensitive to climate change. However, the extent to which the evolutionary history of
42 crayfish is threatened, especially by climate change, remains unclear. To address this, we produced a
43 phylogenetically-informed species prioritisation for the conservation of crayfish and explored the
44 consequences of projected climate change scenarios on crayfish phylogenetic diversity. We used the
45 EDGE2 metric to highlight 70 priority species for conservation. Their combined threatened evolutionary
46 history accounts for 40.7% of the expected phylogenetic diversity loss for all 673 crayfish. Of these
47 priority species, 45.7% are vulnerable to climate change, with the majority (32 species) located in
48 Australia. Under a high warming scenario, individual species that are more evolutionarily distinct are
49 projected to be disproportionately affected. The extinction of climate change vulnerable species could
50 cause the loss of up to 26.3% more phylogenetic diversity than the random expectation, equivalent to 409
51 million years of evolutionary history. Our results highlight the need to consider climate change effects on
52 phylogenetic diversity to fully understand its impact on biodiversity. If actions were taken to avoid a
53 worst-case warming scenario in favour of an intermediate scenario, we could safeguard 251 million years
54 of unique crayfish evolutionary history.

55

56

57 Introduction

58 Despite covering less than 1% of the Earth's surface, freshwater ecosystems contain approximately one
 59 tenth of all recorded animal species (Balian et al., 2007). Freshwater biodiversity is vital for human health
 60 and well-being, but these ecosystems are in a state of crisis (Albert et al., 2020) due to a myriad of
 61 anthropogenic stressors (Dudgeon et al., 2006; Vörösmarty et al., 2010). Superimposed on these is the
 62 growing threat of climate change (Reid et al., 2019), which could threaten half of all freshwater fish
 63 globally (Darwall & Freyhof, 2016). Monitored populations of freshwater vertebrate species have
 64 declined on average by 85% since 1970 (WWF, 2024), a decline notably higher than their terrestrial or
 65 marine counterparts.

66

67 Crayfish comprise more than 690 species worldwide (Crandall & De Grave, 2017; DecaNet, 2024), split
 68 across three northern hemisphere families (Astacidae, Cambaridae, Cambaroididae) and one southern
 69 hemisphere family (Parastacidae). They play several important roles in freshwater communities (Reynolds
 70 et al., 2013), including as keystone species (Füreder et al., 2010), ecosystem engineers (Harvey et al., 2011),
 71 indicators of water quality (Pârvulescu et al., 2011) and some, like the noble crayfish (*Astacus astacus*), serve
 72 as umbrella species for the conservation of others (Edsman et al., 2010; Reynolds et al., 2013). Crayfish
 73 have been the subject of IUCN Red List assessments (Richman et al., 2015) and, together with other
 74 freshwater decapods, fish, and odonates, they are part of a global effort to assess the status of the world's
 75 freshwater fauna (Sayer, 2024). Around one-third of assessed crayfish are threatened with extinction, with
 76 European crayfish experiencing the greatest number of different threats and Australian species being
 77 most at risk from climate change (Richman et al., 2015).

78

79 To conserve crayfish biodiversity into the future, certain species and regions will have to be prioritised.
 80 One way of directing conservation efforts is to consider phylogenetic diversity (PD; Faith 1992), which is
 81 calculated as the sum of all branch lengths connecting a set of species on a phylogenetic tree (Faith,
 82 1992). The conservation of species without consideration of their evolutionary heritage can lead to

substantial losses from the tree of life (Gumbs et al., 2023a). Therefore, prioritising PD not only helps to capture variety among living organisms (Faith, 2008), but also deals with the issues of limited resources by highlighting priority species or areas for conservation that conserve evolutionary history (Crandall, 1998; Whiting et al., 2000). Recent work has also empirically demonstrated that maximising PD captures a greater proportion of diverse features that are beneficial to humanity (such as food and medicine) compared to the random selection of taxa (Molina-Venegas et al., 2021).

The PD conservation framework prioritises species that both represent a disproportionately large amount of evolutionary history and are threatened with extinction (Isaac et al., 2007; Mooers et al., 2008). One way of doing so is through the EDGE metric, which ranks species based on a combination of their irreplaceability (ED: Evolutionary Distinctiveness) and their vulnerability (GE: Global Endangerment) (Isaac et al., 2007). The metric has been applied to several taxonomic groups spanning vertebrates (Gumbs et al., 2018; Isaac et al., 2012; Isaac et al., 2007; Jetz et al., 2014; Stein et al., 2018), invertebrates (Curnick et al., 2015), and plants (Forest et al., 2018). More recently, the method for prioritising species has been updated (Gumbs et al., 2023b); the ‘EDGE2’ metric is now better able to incorporate uncertainty in phylogenetic and extinction risk data and partially weights a species’ value by the extinction risk of related species (Faith, 2008; Steel et al., 2007). The EDGE2 metric has since been applied to all jawed vertebrates (Gumbs et al., 2024), but so far no invertebrate group has been assessed despite the metric’s greater capacity to deal with data-poor clades.

Recent research has led to a number of exciting developments for crayfish, including descriptions of new species, taxonomic revisions (Crandall & De Grave, 2017), the production of a comprehensive molecular phylogeny (Stern et al., 2017), and a trait-based climate change vulnerability analysis (Hossain et al., 2018). Whilst we know that climate change is predicted to confer significant changes in the PD of some clades such as eucalypts (Gonzalez-Orozco et al., 2016), we do not know how it will affect the evolutionary history of most other groups, such as crayfish. In this study, we therefore determined the extent of crayfish PD threatened by climate change and the PD that could be saved by avoiding a high warming

110 scenario. We also produced a priority EDGE list for the conservation of crayfish. Our findings highlight
111 the most irreplaceable and vulnerable crayfish globally, with the potential to guide future conservation
112 action in the face of climate change.

113

Materials and Methods

EDGE2

EDGE2 scores were calculated by assigning each species a probability of extinction (a GE2 score) from a distribution of possible values corresponding to its IUCN Red List category (Gumbs et al., 2023b). The median values of each distribution are as follows: Least Concern = 0.060625, Near Threatened = 0.12125, Vulnerable = 0.2425, Endangered = 0.485, Critically Endangered = 0.97. The extinction risk probabilities for Data Deficient or unassessed species are drawn from a pool of all possible values, with the median value being roughly equivalent to the Vulnerable Red List category. We believe this is an appropriate representation given the link between data-poor species and elevated extinction risks (Bland et al., 2015; Borgelt et al., 2022; Howard & Bickford, 2014).

The branch lengths of a phylogenetic tree are then multiplied by the product of the probabilities of extinction for all descendant species to estimate the expected PD loss for each branch (Gumbs et al., 2023b). Summing the branch lengths that connect a species from the tip to the root of the phylogeny gives its EDGE2 score, which is a measure of the amount of threatened evolutionary history a species is responsible for. ED2 scores are calculated in the same way but preclude the focal species' probability of extinction from the calculation. Summing all branches of an extinction risk-weighted tree gives the total expected PD loss for a given clade. The EDGE2 calculation accounts for phylogenetic complementarity, meaning that the extinction probabilities of related species will influence each other's scores (Steel et al., 2007). EDGE2 scores therefore give greater precedence to species on long, isolated branches of a phylogenetic tree, or to threatened and unique species whose closest relatives are also threatened.

To calculate EDGE2, we first revised the most comprehensive crayfish phylogeny to date (Stern et al., 2017) by inserting known missing species following the methods described in Gumbs, et al. (2023b). We identified missing species based on the crayfish revision of Crandall & De Grave (2017). There were 144 differences between the tip labels in Stern et al.'s phylogeny and the species listed in Crandall & De Grave, largely a result of recent taxonomic revisions to certain groups and new species descriptions. After matching the taxonomies and removing species not listed in Crandall & De Grave, the Stern et al. phylogeny was left with 435 species tips. We imputed missing species into the phylogenetic tree using the function *congeneric.impute* in the *pez* package in R (version 3.6.1) (Pearse et al., 2015), which inserts missing species to a random site within the crown group of the lowest taxonomic rank available (genus, family etc.). At least 23 new species of crayfish have been described since Crandall & De Grave (2017), including 17 species in the family Cambaridae (Bloom et al., 2019; Foltz et al., 2018; Glon et al., 2020; Glon et al., 2019a; Glon et al., 2019b; Johnson, 2018; Loughman et al., 2019; Loughman & Williams, 2018; Lyko, 2017; Pedraza-Lara et al., 2021; Perkins et al., 2019; Taylor, 2018; Álvarez et al., 2021), five in Parastacidae (Huber et al., 2018; Huber et al., 2020; Lukhaup et al., 2018; Patoka et al., 2017), and one in Astacidae (Pârvulescu, 2019). These species were therefore also imputed into the phylogeny. To capture the uncertainty in the placements of imputed species, we conducted the imputation 1000 times.

We then calculated EDGE2 scores for all 673 crayfish species using Red List data (Version 2019-3) (IUCN, 2019), which returned categories for 590 species. We calculated PD and expected PD loss for crayfish to determine the extent of threatened crayfish evolutionary history. The median terminal branch lengths (TBL) of threatened and non-threatened crayfish were contrasted using a Mann Whitney U test. For all Mann Whitney U tests, we used Cliff's delta d to quantify effect size (Cliff, 1993). We then determined the phylogenetic signal of extinction risk in crayfish using Pagel's λ by using species probability of extinction scores across the 1000 possible trees and taking the median value. Using Analysis of Variance (ANOVA) with Tukey's Honest Significant Difference tests, we then compared the TBL and EDGE2 scores of crayfish with tetrapod animal clades using data from Gumbs et al. (2024).

EDGE species

Species are identified as priority EDGE species if they are in a threatened IUCN Red List category (Vulnerable, Endangered, Critically Endangered) and have a median EDGE2 score that is above the median for all species in at least 95% of iterations (Gumbs et al., 2023b). We used IUCN Red List data to map EDGE crayfish by country. We further determined the extent of expected PD loss contributed solely by EDGE crayfish by pruning the extinction risk-weighted phylogenies to contain only EDGE species and summing the branch lengths. Similarly, we also estimated the proportion of expected PD loss contributed by EDGE species. To compare if the contribution of EDGE species to total expected PD loss is consistent across clades, we repeated this procedure using mammal EDGE data (Gumbs et al., 2023b).

In line with the EDGE2 framework, we categorized species into three additional groups: an EDGE2 Watch List, for species that have EDGE2 scores above the median in 95% of cases but are listed as either Least Concern or Near Threatened; an EDGE2 Research List, for species with EDGE2 scores above the median in 95% of cases and that are either Data Deficient or unassessed by the IUCN; and an EDGE2 Borderline List, for threatened species whose EDGE2 scores are above the median for the clade in 80% of iterations.

Data uncertainty

The EDGE2 methodology built on the original EDGE metric (Isaac et al., 2007) by incorporating uncertainty in both phylogenetic and extinction risk; the original EDGE metric (Isaac et al., 2007) could only compute scores for species with sufficient data. To test how this methodological improvement affects species priority lists, we compared the species ranks obtained via the EDGE2 metric and the

original EDGE metric for the 446 species with both phylogenetic and Red List data using a Spearman's correlation. The EDGE metric combines a species' evolutionary distinctiveness (ED) with its extinction risk (GE) and is calculated as $EDGE = \ln(1 + ED) + (GE \times \ln(2))$ (Isaac et al., 2007).

A phylogenetic prioritisation was previously applied to crayfish (Owen et al., 2015) using the software 'Mesquite' (Maddison, 2005; Maddison & Mooers, 2007), which differs from the original EDGE metric in that the ED score is multiplied by a probability of extinction rather than being summed with the GE index. We used the probabilities of extinction calculated in this study to run a Spearman's correlation test between Mesquite EDGE, Isaac et al. EDGE, and EDGE2 to determine the consistency of the methods.

We then explored the effect of species imputation on EDGE2 priority lists by comparing how the scores and ranks of the original set of species from the Stern et al. phylogeny differed with and without the imputation of missing species. To do this, we conducted a series of Spearman's correlations to compare EDGE2 scores, ED2 scores and species ranks of the original set of species before and after phylogenetic imputation. We also calculated the expected PD loss for both sets of species.

Climate change

To determine the extent of crayfish PD threatened by climate change, we compared the amount of PD lost between species vulnerable to climate change and a null model of random extinction. Climate change vulnerable (CCV) species were those identified by a recent traitbased vulnerability analysis for crayfish (Hossain et al., 2018) focusing on species traits relating to sensitivity, adaptive capacity, and exposure to climate change (Foden et al., 2013). In our current study, we considered two of the warming projections investigated in Hossain et al. (2018), intermediate and high (relating to RCP6.0 and RCP8.5, respectively), at two points, 2050 and 2070. Across scenarios, a total of 126 different species were listed as vulnerable

to climate change; under an intermediate warming scenario in 2050 and 2070 there were 87 and 94 CCV species, respectively, whilst there were 107 and 112 species CCV under a high warming scenario (Hossain et al., 2018). We assessed the phylogenetic signal of climate change vulnerability in crayfish using the D statistic (Fritz & Purvis, 2010).

To simulate all CCV species going extinct, we pruned the CCV species for each scenario from each of the 1000 trees and calculated the mean reduction in PD. We then repeated this procedure using a random sample of species of equal size to that of each set of CCV species and where the sets were drawn afresh for each of the 1000 iterations. We tested for differences in PD loss between observed CCV species and randomly drawn species using Mann Whitney U tests. To determine whether climate change is set to affect more evolutionarily distinct species in general, we compared the ED2 scores of CCV species to random sets of species of an equivalent size using Mann Whitney tests.

Finally, we compared the simulated PD loss and ED2 scores of crayfish listed as vulnerable to climate change by Hossain et al. (2018) to those recorded as threatened by climate change on the Red List using Mann Whitney U tests.

Results

Crayfish evolutionary history

Across 673 species, crayfish represent 14,934 million years (MY) of phylogenetic diversity, of which 18.5% (2,769 MY) is at risk of being lost. EDGE2 scores ranged from 0.02 MY to 63 MY (Fig. 1), with the median being 1.51 MY. The median TBL of a crayfish species is 8.46 MY (ranging from 0.157 – 126 MY) and is higher in threatened species (10.2 MY) than non-threatened species (8.11 MY; Mann Whitney U: 41720, $p = < 0.05$), though the effect size was small ($d = 0.118$). Extinction risk displayed a weak phylogenetic signal in crayfish ($\lambda = 0.161$).




Fig. 1

A crayfish phylogeny coloured by EDGE2 score, with higher values in brown and lower values in green, and with crayfish families labelled.

ANOVA with Tukey's Honest Significant Difference test showed that the median log TBL and EDGE2 score of crayfish is significantly greater than observed in amphibians, squamates, birds, and mammals ($p < 0.0001$ for all comparisons; Fig. 2). Turtles showed higher log TBL and EDGE2 scores than crayfish ($p < 0.0001$), whilst crocodilians showed greater log EDGE2 scores ($p < 0.01$) and non-significantly different log TBL.




Fig. 2

Crayfish log terminal branch length and EDGE2 scores compared with tetrapod animal clades.

There was a significant positive correlation between the original EDGE and EDGE2 ranks for the 446 species with data sufficient IUCN assessments ($p = 0.929, p < 0.0001, r^2 = .863$), meaning those species ranked as high priority in EDGE were also likely to be ranked highly in EDGE2 (Appendix S1). Furthermore, Mesquite EDGE ranks were strongly correlated with the original EDGE metric ranks ($p = 0.995, p < 0.0001, r^2 = .99$) and with EDGE2 ranks ($p = 0.922, p < 0.0001, r^2 = .85$).

EDGE Crayfish

Seventy species were in a threatened IUCN Red List category and had an EDGE2 score above the median with 95% certainty and were thus deemed EDGE species (Table 1; see Figshare repository).

267 These species had EDGE2 scores ranging from 2.75 to 63 MY (median = 13.8 MY), and span 14 genera,
 268 three families, and four main regions, with 44 species in Australia, 21 in USA, four in Mexico, and one,
 269 *Austropotamobius pallipes* (the white clawed crayfish), found in Europe (Fig. 3). There were 29 Critically
 270 Endangered species, 34 Endangered species, and 7 Vulnerable species. The top ranked EDGE species
 271 was the Critically Endangered Scottsdale burrowing crayfish, *Engaenus spinicaudatus*, a Tasmanian endemic.

Family	Species	Red List category	Terminal branch length (MY)	ED2 (MY)	EDGE2 (MY)	Native range	Climate change vulnerable
Parastacidae	<i>Engaenus spinicaudatus</i>	CR	65.4	65.5	63.0	Australia	NO
Parastacidae	<i>Tennibranchiurus glypticus</i>	EN	124.8	124.8	61.3	Australia	YES
Parastacidae	<i>Engaenus mallacoota</i>	CR	52.4	53.0	50.3	Australia	YES
Parastacidae	<i>Euastacus robertsi</i>	CR	14.2	49.9	45.1	Australia	YES
Parastacidae	<i>Euastacus bindal</i>	CR	39.6	42.3	39.9	Australia	YES
Parastacidae	<i>Euastacus setosus</i>	CR	37.2	42.0	39.0	Australia	YES
Parastacidae	<i>Euastacus eungella</i>	CR	37.2	42.0	38.8	Australia	YES
Parastacidae	<i>Euastacus fleckeri</i>	EN	14.2	83.2	38.4	Australia	YES
Parastacidae	<i>Engaenus martigener</i>	EN	75.3	77.5	36.3	Australia	NO
Parastacidae	<i>Euastacus monteithorum</i>	CR	37.3	37.8	35.2	Australia	NO
Parastacidae	<i>Engaenus phyllocercus</i>	EN	69.6	69.6	32.5	Australia	NO
Cambaridae	<i>Creaserinus bortoni</i>	CR	35.5	35.5	32.1	United States	NO
Parastacidae	<i>Euastacus jagara</i>	CR	27.7	29.7	27.7	Australia	NO
Parastacidae	<i>Euastacus gamilaroi</i>	CR	26.9	28.1	26.1	Australia	NO
Parastacidae	<i>Astacopsis gouldi</i>	EN	50.2	52.3	25.4	Australia	NO
Parastacidae	<i>Engaenus sternalis</i>	CR	23.4	27.6	25.2	Australia	NO

Parastacidae	<i>Euastacus mirangudjin</i>	CR	24.2	24.2	23.0	Australia	YES
Parastacidae	<i>Euastacus maida</i>	CR	22.6	24.7	22.8	Australia	YES
Parastacidae	<i>Euastacus yigara</i>	CR	21.4	24.7	22.7	Australia	YES
Parastacidae	<i>Engaeus granulatus</i>	CR	22.1	22.9	22.0	Australia	NO
Cambaridae	<i>Cambarus tartarus</i>	CR	17.6	23.7	21.6	United States	NO
Parastacidae	<i>Engaewa reducta</i>	EN	23.8	37.9	19.1	Australia	YES
Cambaridae	<i>Cambarus aculabrum</i>	CR	14.9	21.4	19.0	United States	NO
Parastacidae	<i>Euastacus balanensis</i>	EN	34.9	36.1	17.4	Australia	YES
Parastacidae	<i>Euastacus girumalayn</i>	CR	11.4	19.0	17.2	Australia	YES

TABLE 1

The top 25 EDGE crayfish species, where climate change vulnerability (marked as YES/NO) was taken from Hossain et al. (2018).




Fig. 3

The number of priority EDGE crayfish per country, where presence in a country was determined using IUCN data.

Of the 2,769 MY of crayfish PD that is expected to be lost given present day extinction risks, 40.8% (1,128 MY) is contributed by the 70 EDGE crayfish, despite EDGE species representing just 10.3% of species richness. This is comparable to that observed in mammals, where 36% of the expected loss of the clade is attributable to 10.3% of species. Together, the 140 threatened crayfish species represent 52% (1,440 MY) of expected crayfish phylogenetic diversity loss, of which 78.5% of this is attributable to the 70 that are EDGE listed.

Seven non-threatened species had EDGE2 scores above the median with 95% confidence; one, the monotypic *Samastacus spinifrons* (endemic to Argentina and Chile), is Data Deficient and was therefore placed on the EDGE Research List. The six others were all Near Threatened and were therefore placed on the EDGE Watch List. Thirty-three threatened species had EDGE2 scores above the median with 80% confidence and were therefore marked as Borderline EDGE species. EDGE2 scores, EDGE lists and expected phylogenetic diversity loss trees will be made available on Figshare on publication (10.6084/m9.figshare.27283635).

Data uncertainty

Overall, 227 species (33.7%) lacked extinction risk data (i.e., they were Data Deficient or not evaluated on the IUCN Red List) and 238 (35.4%) were without phylogenetic information. Not accounting for these

species would lead us to consider 20.7% (3,088 MY) less phylogenetic diversity and 28.6% (791 MY) less expected loss, thus dramatically underestimating the true extent of crayfish threatened evolutionary history.

A Spearman's correlation revealed that species ranks ($S_r = 0.979$, $p < .0001$, $r^2 = .959$), EDGE2 scores ($S_r = 0.979$, $p < .0001$, $r^2 = .959$), and ED2 scores ($S_r = 0.973$, $p < .0001$, $r^2 = .902$) for the 435 species featuring in the Stern et al. phylogeny were highly consistent before and after the imputation of missing species (Fig. 4).




Fig. 4

The effect of phylogenetic imputation on species ranks, EDGE2 scores, and ED2 scores. Points represent individual species, coloured by their IUCN Red List categories.

Climate change

For all four scenarios explored, the simulated extinction of climate change vulnerable species represents a greater loss of total PD than random groups of the same size (Fig. 5). Crayfish projected to be CCV under a 2070 high warming scenario (Hossain et al., 2018) showed the greatest departure from the random expectation, threatening 409 million more years of evolutionary history (Mann Whitney U = 860578, $p < 0.0001$, $d = 0.721$), followed by 315 MY of PD loss for crayfish that are CCV under a 2050 high warming scenario (Mann Whitney U = 804058, $p < 0.0001$, $d = 0.608$). Both groups of crayfish projected to be CCV under an intermediate warming scenario in 2050 and 2070 showed a similar mean difference in PD loss of 158 MY and 152 MY, respectively (Mann Whitney U = 657521, $p < 0.0001$, $d = 0.315$; Mann Whitney U = 652865, $p < 0.0001$, $d = 0.306$). The extinction of CCV crayfish species therefore represents between 11.9% and 26.3% more PD being lost than the random expectation.




Fig. 5

The effect of climate change on crayfish evolutionary history. Left: The median change in phylogenetic diversity resulting from the loss of all climate change vulnerable (CCV) species beyond that of the random expectation (denoted by the black dotted line) under each warming scenario, where negative values represent a greater loss. Right: The median ED2 of a CCV crayfish compared with the random expectation (dotted line).

The median ED2 of a crayfish species is 10 MY, whilst the median ED2 of a CCV species is 12.3 MY. The highest median ED2 for a CCV species was 125 MY for *Tenuibranchiurus glypticus*, the second highest ranked EDGE2 species and smallest crayfish in the world at 3 cm, whilst the lowest was the Critically Endangered and visually striking *Cherax tenuimanus* with 0.433 MY.

For all four scenarios of projected CCV crayfish, the observed distributions of ED2 differed significantly ($p < 0.0001$) from the random expectations in Mann-Whitney tests (Fig. 5), with all scenarios except a 2070 intermediate warming affecting crayfish with higher ED2 scores more than expected and with each having large effect sizes. The greatest departure from the random expectation was for species projected to be CCV under a 2050 high warming scenario, where vulnerable species were on average 3.22 MY older ($U = 962913$, $d = 0.926$). This was followed by 2.2 MY older for a 2070 high warming scenario ($U = 905521$, $d = 0.811$) and then 1.77 MY for species projected as CCV under a 2050 intermediate warming scenario ($U = 812544$, $d = 0.625$). Species projected to be CCV under a 2070 intermediate warming scenario, however, had a lower ED2 than the random expectation by 0.76 MY ($U = 331178$, $d = -0.338$).

We found that the simulated extinction of all 134 species listed as CCV across the four scenarios (median = 12,644 MY) resulted in 0.64% more PD being lost in comparison to the simulated extinction of the 86 crayfish that are listed as threatened by climate change using the Red List (median = 12,564 MY; $U = 581982$, $p < 0.0001$, $d = 0.164$). However, the crayfish threatened by climate change on the Red List had ED2 scores (median = 20.7 MY) that were 68.5% greater than those of the CCV crayfish (median = 12.3; $U = 7834.5$, $p < 0.0001$, $d = 0.36$).

360

361

362 Discussion

363

364 Our study draws attention to the underappreciated plight of crayfish, a group where almost one-fifth of
 365 evolutionary history is at risk of extinction, and that are disproportionately threatened by climate change.
 366 Given that roughly one-third of crayfish lack sufficient conservation assessments, and that Data Deficient
 367 species are likely to have elevated extinction risks (Borgelt et al., 2022), this predicted loss is likely an
 368 underestimate. Furthermore, we highlight that threatened crayfish tend to be more evolutionarily distinct
 369 than non-threatened crayfish and, in our comparison with tetrapod animals, we demonstrated that
 370 crayfish rank amongst the most evolutionarily distinct clades examined so far.

371

372 Together, the 70 EDGE crayfish species identified here represent 40% of expected PD loss for all
 373 crayfish, despite comprising just a tenth of crayfish species richness. Conserving the remaining 70
 374 threatened crayfish that do not qualify as EDGE species would safeguard just 10% more of the total
 375 expected loss. This demonstrates that safeguarding a relatively small number of priority species can
 376 protect a large portion of evolutionary history. If conservation efforts overlook evolutionary history as a
 377 key component of biodiversity, however, then we may face a situation where large and unique branches
 378 are lost from the Tree of Life (De Meester et al., 2024; Forest et al., 2015; Hendry et al., 2010).

379

380

381 Of the 70 EDGE priority species, 31 are within Australia's largest crayfish genus *Euastacus*. This is the
 382 most threatened crayfish genus with more than one species, where 74% of the group's 53
 383 morphologically and ecologically diverse species are threatened with extinction (Furse & Coughran,
 384 2011). The high proportion of EDGE species can be explained by the phylogenetic complementarity

embedded within the EDGE2 metric, which ensures that closely related species that are similarly threatened have similar conservation value. In the case of *Euastacus*, the high concentration of priority species therefore reflects that each species represents a possible route to securing a portion of the whole clade's shared evolutionary history. The EDGE priority list can then be used in accompaniment with known information on conservation feasibility to determine which species to protect and, in doing so, efficiently deliver huge gains for the preservation of the crayfish Tree of Life.

The nature of the EDGE metric explains the inclusion of and exclusion of other crayfish groups of interest. *Tenuibranchiurus glypticus* is a monotypic parastacid crayfish with the second highest ED2 score that is found in the swamps outside of Brisbane, a region of known value for its threatened tetrapod evolutionary history (Pipins et al., 2024). Given its extreme phylogenetic isolation and Endangered threat status, the species ranks as the second top scoring EDGE species on the list. Elsewhere, the Endangered *Austropotamobius pallipes* is Europe's only EDGE crayfish. It appears on the EDGE list due to the genus' notable evolutionary history, with the median genus-level ED2 score (29 MY) ranking sixth highest among all 34 genera, and the fact that it is the only member within its four species genus that has a data-sufficient Red List assessment. Some genera of interest do not feature on the list, like the genus *Paraneophrops*, which contains the only two endemic crayfish on the islands of New Zealand, but which do not constitute conservation priorities as both species are marked as Least Concern on the Red List. Elsewhere, no species from the three South American endemic genera (*Virilastacus*, *Samastacus*, and *Parastacus*) make it on the list. This is because of a lack of conservation data on these genera, with all 4 *Virilastacus* species and the only *Samastacus* species being Data Deficient, along with 12 of the 14 *Parastacus* species, despite their high median ED2 scores (16 MY, 70.1 MY, and 19.7 MY, respectively) and the suggested threatened status of many (Almerão et al., 2015). Given its especially large ED2 score, *Samastacus spinifrons* was placed on the EDGE2 Research List, highlighting its urgent need for a conservation assessment. Elsewhere, no Malagasy species (*Astacoides*) make it on the list, despite four being Vulnerable (Crandall et al., 2022; Jones et al., 2007). This is due to their phylogenetic uncertainty with three of seven species requiring phylogenetic imputation. Three *Astacoides* species did make it on the

EDGE Borderline list, reflecting their notable threatened evolutionary history which fails to meet the 95% certainty threshold.

Our study presents the first EDGE2 assessment for an invertebrate group, building on the previous phylogenetic crayfish prioritisation of Owen et al. (2015) but employing a standardised procedure that allows for comparisons to other EDGE2 assessed groups like tetrapod animals (Gumbs et al., 2024). Though the data deficiency observed in crayfish is beyond that of most tetrapod animal clades, we found that the metric still performed robustly, with species ranks remaining largely consistent before and after the imputation of the 35% of crayfish with absent phylogenetic data (Fig. 4). We therefore also highlight this study as being an important stepping stone for determining the validity of such methods for clades with even larger data gaps, such as in flowering plants, fungi, and insects.

Our results echo the concerns of previous studies on the impact that climate change will have on phylogenetic diversity (Faith & Richards, 2012; Gonzalez-Orozco et al., 2016). A high warming scenario could affect up to 409 million more years of crayfish evolutionary history than expected at random, whilst following an intermediate warming scenario could spare 251 MY in comparison. Furthermore, a high warming scenario threatens individual species that are more evolutionarily distinct than the random expectation by 3.22 MY, due to the greater proportion of parastacid crayfish that are considered CCV in high warming scenarios (Hossain et al., 2018). The parastacids, native to southern latitudes, are known to have lower diversification rates than crayfish of northern latitudes and exhibit longer terminal branch lengths (Owen et al., 2015; Toon et al., 2010). Australia is the centre of parastacid diversity and, owing to their extensive evolutionary history, parastacids are also disproportionately represented as EDGE species, making up 44 of the 70 EDGE priority crayfish, with 23 of these being climate change vulnerable (Hossain et al., 2018). Avoiding a high warming scenario will therefore be essential for the conservation of Australia's highly distinct crayfish.

We found that only 29 species were jointly marked as being at risk from climate change according to the climate change trait-based vulnerability analysis (CCVA) and the Red List threat classification. This is unsurprising since IUCN Red List assessments generally focus on a shorter time period than that evoked by CCVA. There may also be a bias inherent in Red List assessments, which often generalize associations, for example, between temperature tolerance and climate change risk to generate assessments for large species groups (Trull et al., 2018). Meanwhile, trait-data incompleteness introduces ambiguity; in the crayfish CCVA of Hossain et al. (2018), traits with unknown trait values were scored as *low* in relation to a species' overall vulnerability to climate change, leading to an average of 18% of recorded crayfish being listed as CCV. In contrast, when unknown trait values were instead scored as *high*, an average of 48% of species were CCV. Approximately one-third of the crayfish used in this study had no information relating to their risk from climate change. When these discrepancies are taken into consideration, it is possible that a much larger proportion of crayfish PD is threatened than our study suggests.

Phylogenetic diversity has been gaining traction within conservation, with recent developments including the EDGE index featuring as a complementarity indicator under Target 4 of the Kunming-Montreal Global Biodiversity Framework (Secretariat of the Convention on Biological Diversity, 2022) and the establishment of the IUCN SSC Phylogenetic Diversity Task Force (Gumbs et al., 2023a). We believe the creation of this crayfish EDGE list can capitalise on this momentum by feeding into the conservation activities of the EDGE of Existence programme, which to date has funded work on over 130 EDGE species. Furthermore, although an increasing number of crayfish are being recognised by the Endangered Species Act in the US (Taylor et al., 2019), crayfish are still rarely featured in freshwater conservation policy and action globally (Danilović et al., 2022; Taylor et al., 2019). As most EDGE crayfish and climate change vulnerable crayfish are located within Australia, we advise a reassessment of this country's threatened parastacids coupled with action to secure their long-term future, taking phylogenetic diversity into account across the bioregionalization of Australia (Whiting et al., 2000).

464 Given that a high warming scenario threatens a disproportionately large amount of crayfish PD and
465 evolutionarily distinct species, the absence of climate mitigation strategies will likely see the loss of some
466 of the world's most unique and irreplaceable crayfish. If action can be taken to curb carbon emissions and
467 avoid a worst-case scenario, however, we could safeguard hundreds of millions of years of crayfish
468 evolutionary history.

469

470 **Supporting Information**

471 Additional supporting information may be found online in the Supporting Information section at the end
472 of this article.

473

474 **Acknowledgements**

475 SP was funded by the NERC Science and Solutions for a Changing Planet Doctoral Training Programme
476 (grant number NE/S007415/1), the CASE component of which was funded by On the Edge. MB was
477 supported by a grant from the Rufford Foundation. KAC was supported as a VisitANTS Visiting Scholar
478 as part of the Biodiverse Anthropocenes program at the University of Oulu, Finland.

479

480 **Author contributions**

481 RG, JR, and MB conceived the study. SP, RG, and JR designed the analyses. SP conducted the analyses.
482 LB and MAH provided crayfish trait-based vulnerability analysis data. KAC provided crayfish
483 phylogenetic data. RG provided the EDGE2 code and tetrapod data. RG, JR, and MB supervised the
484 study. SP led the writing of the manuscript. All authors contributed critically to the drafts and gave final
485 approval for publication.

486

487

488 **References**

489

490 Albert, J. S., Destouni, G., Duke-Sylvester, S. M., Magurran, A. E., Oberdorff, T., Reis, R. E.,
491 Winemiller, K. O., Ripple, W. J. (2020). Scientists' warning to humanity on the freshwater
492 biodiversity crisis. *Ambio*. **50**, 1-10.

493 Almerão, M. P., Rudolph, E., Souty-Grosset, C., Crandall, K., Buckup, L., Amouret, J., Verdi, A.,
494 Santos, S., Araujo, P. B. D. (2015). The native South American crayfishes (Crustacea,
495 Parastacidae): state of knowledge and conservation status. *Aquat. Conserv.: Mar.*
496 *Freshw. Ecosyst*. **25**, 288-301.

497 Álvarez, F., Torres, E., & Villalobos, J. L. (2021). New species of crayfish of the genus
498 Procambarus, and notes on Procambarus pilosimanus (Decapoda: Cambaridae) from
499 Chiapas, Mexico. *Rev. Mex. Biodivers.* **92**, e923833.

500 Balian, E.V., Segers, H., Martens, K., Lévêque, C. (2007). The Freshwater Animal Diversity
501 Assessment: an overview of the results. In *Freshwater Animal Diversity Assessment*
502 627-637. Balian, E.V., Lévêque, C., Segers, H., Martens, K. Dordrecht: Springer.

503 Bland, L. M., Collen, B., Orme, C. D. L., & Bielby, J. (2015). Predicting the conservation status of
504 data-deficient species. *Conserv. Biol.* **29**, 250-259.

505 Bloom, E. T., McCall, B. L., Schuster, G. A., & Blanton, R. E. (2019). Systematics and description of
506 a new species of Faxonius Ortmann, 1905 (Decapoda: Astacidea: Cambaridae) from the
507 Red River system of Kentucky and Tennessee, USA. *J. Crustac. Biol*. **39**, 40-53.

508 Borgelt, J., Dorber, M., Høiberg, M. A., & Verones, F. (2022). More than half of data deficient
509 species predicted to be threatened by extinction. *Commun. Biol.* **5**, 679.

510 Cliff, N. (1993). Dominance statistics: Ordinal analyses to answer ordinal questions. *Psychol.*
511 *Bull.* **114**, 494.

512 Crandall, K., Jones, J., & Razanabolana, J. R. (2022). Parastacidae, Astacoides, Freshwater
513 Crayfishes, Orana, Orambanonga, Orambato. In *The New Natural History of Madagascar*.
514 908-912. Goodman, S. M. Princeton: Princeton University Press.

515 Crandall, K. A. (1998). Conservation phylogenetics of Ozark crayfishes: assigning priorities for
516 aquatic habitat protection. *Biol. Conserv.* **84**, 107-117.

517 Crandall, K. A., & De Grave, S. (2017). An updated classification of the freshwater crayfishes
518 (Decapoda: Astacidea) of the world, with a complete species list. *J. Crustac. Biol.* **37**, 615-
519 653.

520 Curnick, D., Head, C., Huang, D., Crabbe, M. J. C., Gollock, M., Hoeksema, B., Johnson, K., Jones, R.,
521 Koldewey, H., Obura, D. (2015). Setting evolutionary-based conservation priorities for a
522 phylogenetically data-poor taxonomic group (Scleractinia). *Anim. Conserv.* **18**, 303-312.

523 Danilović, M., Maguire, I., & Füreder, L. (2022). Overlooked keystone species in conservation
524 plans of fluvial ecosystems in Southeast Europe: a review of native freshwater crayfish
525 species. *Knowl. Manag. Aquat. Ecosyst* **423**, 21.

526 Darwall, W. R. T., & Freyhof, J. (2016). Lost fishes, who is counting? The extent of the threat to
527 freshwater fish biodiversity. In *Conservation of freshwater fishes*: 1-36. Closs, G.P.,
528 Krkosek, M. & Olden, J. D. Cambridge: Cambridge University Press.

529 De Meester, L., Vázquez-Domínguez, E., Kassen, R., Forest, F., Bellon, M. R., Koskella, B., Scherson,
530 R. A., Colli, L., Hendry, A. P., & Crandall, K. A. (2024). A link between evolution and
531 society fostering the UN sustainable development goals. *Evol. Appl.* **17**, e13728.

532 DecaNet. (2024). *DecaNet*, Accessed 18 February 24. <https://www.decanet.info>

533 Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z.-I., Knowler, D. J., Lévêque, C., Naiman,
534 R. J., Prieur-Richard, A.-H., Soto, D., Stiassny, M. L. J., & Sullivan, C. A. (2006). Freshwater
535 biodiversity: importance, threats, status and conservation challenges. *Biol. Rev.* **81**, 163-
536 182.

537 Edsman, L., Füreder, L., Gherardi, F., & Souty-Grosset, C. (2010). *Astacus astacus*. The IUCN Red
538 List of Threatened Species 2010, e.T2191A9338388.

539 Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1-10.

540 Faith, D. P. (2008). Threatened species and the potential loss of phylogenetic diversity:
541 conservation scenarios based on estimated extinction probabilities and phylogenetic
542 risk analysis. *Conserv. Biol.* **22**, 1461-1470.

543 Faith, D. P., & Richards, Z. T. (2012). Climate change impacts on the tree of life: changes in
544 phylogenetic diversity illustrated for *Acropora* corals. *Biology*, **1**, 906-932.

545 Foden, W. B., Butchart, S. H., Stuart, S. N., Vié, J.-C., Akçakaya, H. R., Angulo, A., DeVantier, L. M.,
546 Gutsche, A., Turak, E. & Cao, L. (2013). Identifying the world's most climate change
547 vulnerable species: a systematic trait-based assessment of all birds, amphibians and
548 corals. *Plos One*, **8**, e65427.

549 Foltz, D. A., Sadecky, N. M., Myers, G. A., Fetzner Jr, J. W., Welsh, S. A., Stocker, G. W., Glon, M. G. &
550 Thoma, R. F. (2018). *Cambarus loughmani*, a new species of crayfish (Decapoda:
551 Cambaridae) endemic to the pre-glacial Teays River Valley in West Virginia, USA. *J. Nat.*
552 *Hist.*, **52**, 2875-2897.

553 Forest, F., Crandall, K. A., Chase, M. W., & Faith, D. P. (2015). Phylogeny, extinction and
554 conservation: embracing uncertainties in a time of urgency. *Philos. Trans. R. Soc. Lond. B*
555 *Biol. Sci.* **370**, 20140002.

- 556 Forest, F., Moat, J., Baloch, E., Brummitt, N. A., Bachman, S. P., Ickert-Bond, S., Hollingsworth, P.
557 M., Liston, A., Little, D. P. & Mathews, S. (2018). Gymnosperms on the EDGE. *Sci. Rep.* **8**,
558 1-11.
- 559 Fritz, S. A., & Purvis, A. (2010). Selectivity in mammalian extinction risk and threat types: a new
560 measure of phylogenetic signal strength in binary traits. *Conserv. Biol.* **24**, 1042-1051.
- 561 Furse, J. M., & Coughran, J. (2011). An assessment of the distribution, biology, threatening
562 processes and conservation status of the freshwater crayfish, genus *Euastacus*
563 (Decapoda, Parastacidae), in continental Australia. I. Biological background and current
564 status. *Crustac. Monogr.* **15**, 241-252.
- 565 Füreder, L., Gherardi, F., Holdich, D., Reynolds, J., Sibley, P., & Souty-Grosset, C. (2010).
566 *Austropotamobius pallipes*. *The IUCN Red List of Threatened Species 2010*,
567 e.T2430A9438817.
- 568 Glon, M. G., Adams, S. B., Loughman, Z. J., Myers, G. A., Taylor, C. A., & Schuster, G. A. (2020). Two
569 new species of burrowing crayfish in the genus *Lacunicambarus* (Decapoda:
570 *Cambaridae*) from Alabama and Mississippi. *Zootaxa*. **4802**, 401-439.
- 571 Glon, M. G., Thoma, R. F., Daly, M., & Freudenstein, J. V. (2019a). *Lacunicambarus chimera*: a new
572 species of burrowing crayfish (Decapoda: *Cambaridae*) from Illinois, Indiana, Kentucky,
573 and Tennessee. *Zootaxa*. **4544**, 451-478.
- 574 Glon, M. G., Williams, B. W., & Loughman, Z. J. (2019b). *Lacunicambarus dalyae*: a new species of
575 burrowing crayfish (Decapoda: *Cambaridae*) from the southeastern United States.
576 *Zootaxa*. **4683**, 361-380.
- 577 Gonzalez-Orozco, C., Pollock, L., Thornhill, A., Mishler, B., Knerr, N., Laffan, S., Miller, J., Rosauer,
578 D., Faith, D., Nipperess, D., Kujala, H., Linke, S., Butt, N., Külheim, C., Crisp, M. & Gruber, B.

579 (2016). Phylogenetic approaches reveal biodiversity threats under climate change. *Nat*
580 *Clim. Change*. **6**, 1110–1114.

581 Gumbs, R., Chaudhary, A., Daru, B. H., Faith, D. P., Forest, F., Gray, C. L., Kowalska, A., Lee, W. S.,
582 Pellens, R. & Pipins, S., Pollock, L. J., Rosindell, J., Scherson, R. A., Owen, N. (2023a).
583 Indicators to monitor the status of the tree of life. *Conserv. Biol.* **37**, e14138.

584 Gumbs, R., Gray, C. L., Böhm, M., Burfield, I. J., Couchman, O. R., Faith, D. P., Forest, F., Hoffmann,
585 M., Isaac, N. J. B., Jetz, W., Mace, G. M., Mooers, A. O., Safi, K., Scott, O., Steel, M., Tucker, C.
586 M., Pearse, W. D., Owen, N. R. & Rosindell, J. (2023b). The EDGE2 protocol: Advancing
587 the prioritisation of Evolutionarily Distinct and Globally Endangered species for
588 practical conservation action. *PLoS Biol.* **21**, e3001991.

589 Gumbs, R., Gray, C. L., Wearn, O. R., & Owen, N. R. (2018). Tetrapods on the EDGE: Overcoming
590 data limitations to identify phylogenetic conservation priorities. *Plos One*. **13**, e0194680.

591 Gumbs, R., Scott, O., Bates, R., Böhm, M., Forest, F., Gray, C. L., Hoffmann, M., Kane, D., Low, C.,
592 Pearse, W. D., Pipins, S., Tapley, B., Turvey, S. T., Jetz, W., Owen, N. R., & Rosindell, J.,
593 (2024). Global conservation status of the jawed vertebrate Tree of Life. *Nat. Commun.*
594 **15**, 1101.

595 Harvey, G. L., Moorhouse, T. P., Clifford, N. J., Henshaw, A. J., Johnson, M. F., Macdonald, D. W.,
596 Reid, I. & Rice, S. P. (2011). Evaluating the role of invasive aquatic species as drivers of
597 fine sediment-related river management problems: the case of the signal crayfish
598 (*Pacifastacus leniusculus*). *Prog. Phys. Geogr.*, **35**, 517-533.

599 Hendry, A. P., Lohmann, L. G., Conti, E., Cracraft, J., Crandall, K. A., Faith, D. P., Häuser, C., Joly, C.
600 A., Kogure, K., Larigauderie, A., Magallón, S., Moritz, C., Tillier, S., Zardoya, R., Prieur-
601 Richard, A. H., Walther, B. A., Yahara, T., Donoghue, M. J. (2010). Evolutionary biology in
602 biodiversity science, conservation, and policy: a call to action. *Evolution*. **64**, 1517-1528.

603 Hossain, M. A., Lahoz-Monfort, J. J., Burgman, M. A., Böhm, M., Kujala, H., & Bland, L. M. (2018).
604 Assessing the vulnerability of freshwater crayfish to climate change. *Divers. Distrib.* **24**,
605 1830-1843.

606 Howard, S. D., & Bickford, D. P. (2014). Amphibians over the edge: silent extinction risk of Data
607 Deficient species. *Divers. Distrib.* **20**, 837-846.

608 Huber, A. F., Ribeiro, F. B., & Araujo, P. B. (2018). New endemic species of freshwater crayfish
609 *Parastacus* Huxley, 1879 (Crustacea: Decapoda: Parastacidae) from the Atlantic forest in
610 southern Brazil. *Nauplius*. **26**, 2018015.

611 Huber, A. F., Rockhill, E. R., Araujo, P. B., & Ribeiro, F. B. (2020). A new species of burrowing
612 crayfish in genus *Parastacus* Huxley, 1879 (Decapoda, Parastacidae) from the Sinos
613 River Basin, Southern Brazil. *Zool. Stud.* **59**, 1-13.

614 International Union for Conservation of Nature (IUCN). (2020). IUCN Red List of Threatened
615 Species: Version 2019–3. www.iucnredlist.org

616 Isaac, N. J., Redding, D. W., Meredith, H. M., & Safi, K. (2012). Phylogenetically-informed
617 priorities for amphibian conservation. *PLoS One*. **7**, e43912.

618 Isaac, N. J., Turvey, S. T., Collen, B., Waterman, C., & Baillie, J. E. (2007). Mammals on the EDGE:
619 conservation priorities based on threat and phylogeny. *PLoS One*, **2**, e296.

620 Jetz, W., Thomas, G. H., Joy, J. B., Redding, D. W., Hartmann, K., & Mooers, A. O. (2014). Global
621 distribution and conservation of evolutionary distinctness in birds. *Curr. Biol.* **24**, 919-
622 930.

623 Johnson, D. P. (2018). The subgenus *Ortmannicus* (Decapoda: Cambaridae) in Texas, with
624 descriptions of new species. *Zootaxa*. **4468**, 1-63.

- 625 Jones, J. P., Andriahajaina, F. B., Hockley, N. J., Crandall, K. A., & Ravoahangimalala, O. R. (2007).
626 The ecology and conservation status of Madagascar's endemic freshwater crayfish
627 (Parastacidae; Astacoides). *Freshw. Biol.* **52**, 1820-1833.
- 628 Loughman, Z. J., Welsh, S. A., & Thoma, R. F. (2019). *Cambarus fetzneri* sp. nov., a new species of
629 burrowing crayfish (Decapoda: Cambaridae) from the Allegheny Mountains of Virginia
630 and West Virginia, USA. *Zootaxa*. **4651**, 38-50.
- 631 Loughman, Z. J., & Williams, B. W. (2018). *Cambarus polypilosus*, a new species of stream-
632 dwelling crayfish (Decapoda: Cambaridae) from the Western Highland Rim of
633 Tennessee, USA. *Zootaxa*. **4403**, 171-185.
- 634 Lukhaup, C., Eprilurahman, R., & von Rintelen, T. (2018). Two new species of crayfish of the
635 genus *Cherax* from Indonesian New Guinea (Crustacea, Decapoda, Parastacidae).
636 *ZooKeys*. **769**, 89-116.
- 637 Lyko, F. (2017). The marbled crayfish (Decapoda: Cambaridae) represents an independent new
638 species. *Zootaxa*. **4363**, 544–552.
- 639 Maddison, W. (2005). Mesquite: a modular system for evolutionary analysis. Version 1.06.
640 <https://www.mesquiteproject.org/>.
- 641 Maddison, W. P., & Mooers, A. Ø. (2007). Tuatara: Conservation priority in a phylogenetic
642 context. Version 1.0. <http://mesquiteproject.org/packages/tuatara>
- 643 Molina-Venegas, R., Rodríguez, M. Á., Pardo-de-Santayana, M., Ronquillo, C., & Mabberley, D. J.
644 (2021). Maximum levels of global phylogenetic diversity efficiently capture plant
645 services for humankind. *Nat. Ecol. Evol.* **5**, 583-588.
- 646 Mooers, A. Ø., Faith, D. P., & Maddison, W. P. (2008). Converting endangered species categories
647 to probabilities of extinction for phylogenetic conservation prioritization. *Plos One*. **3**,
648 e3700.

- 649 Owen, C. L., Bracken-Grissom, H., Stern, D., & Crandall, K. A. (2015). A synthetic phylogeny of
650 freshwater crayfish: insights for conservation. *Philos. Trans. R. Soc. B: Biol. Sci.* **370**,
651 20140009.
- 652 Patoka, J., Blaha, M., & Kouba, A. (2017). *Cherax acherontis* (Decapoda: parastacidae), the first
653 cave crayfish from the Southern Hemisphere (Papua Province, Indonesia). *Zootaxa*.
654 **4363**, 137-144.
- 655 Pearse, W. D., Cadotte, M. W., Cavender-Bares, J., Ives, A. R., Tucker, C. M., Walker, S. C., & Helmus,
656 M. R. (2015). pez: phylogenetics for the environmental sciences. *Bioinformatics*. **31**,
657 2888-2890.
- 658 Pedraza-Lara, C., Ortiz-Herrera, H. S., & Jones, R. W. (2021). A new species of crayfish of the
659 genus *Cambarellus* (Decapoda: Cambaridae) from central Mexico. *Rev. Mex. Biodiv.* **92**,
660 e923150.
- 661 Perkins, M. A., Williams, B. W., & Russ, W. T. (2019). *Cambarus franklini*, a new crayfish
662 (Decapoda: Cambaridae) from the Catawba River Basin in western North Carolina, USA.
663 *Zootaxa*. **4568**, 520-532.
- 664 Pipins, S., Baillie, J. E., Bowmer, A., Pollock, L. J., Owen, N., & Gumbs, R. (2024). Advancing EDGE
665 Zones to identify spatial conservation priorities of tetrapod evolutionary history. *Nat.*
666 *Commun.* **15**, 7672.
- 667 Pârvulescu, L. (2019). Introducing a new *Austropotamobius* crayfish species (Crustacea,
668 Decapoda, Astacidae): A miocene endemism of the Apuseni Mountains, Romania. *Zool.*
669 *Anz.* **279**, 94-102.
- 670 Pârvulescu, L., Pacioglu, O., & Hamchevici, C. (2011). The assessment of the habitat and water
671 quality requirements of the stone crayfish (*Austropotamobius torrentium*) and noble

672 crayfish (*Astacus astacus*) species in the rivers from the Anina Mountains (SW
673 Romania). *Knowl. Manag. Aquat. Ecosyst* **401**, 03.

674 Reid, A. J., Carlson, A. K., Creed, I. F., Eliason, E. J., Gell, P. A., Johnson, P. T., Kidd, K. A.,
675 MacCormack, T. J., Olden, J. D., Ormerod, S. J., John P. Smol, J. P., William W. Taylor, W. W.,
676 Tockner, K., Vermaire, J. C., Dudgeon, D., Cooke, S. J. (2019). Emerging threats and
677 persistent conservation challenges for freshwater biodiversity. *Biol. Rev.* **94**, 849-873.

678 Reynolds, J., Souty-Grosset, C., & Richardson, A. (2013). Ecological roles of crayfish in freshwater
679 and terrestrial habitats. *Freshw. Crayfish*. **19**, 197-218.

680 Richman, N. I., Böhm, M., Adams, S. B., Alvarez, F., Bergey, E. A., Bunn, J. J. S., Burnham, Q.,
681 Cordeiro, J., Coughran, J., Crandall, K. A., Dawkins, K. L., DiStefano, R. J., Doran, N. E.,
682 Edsman, L., Eversole, A. G., Füreder, L., Furse, J. M., Gherardi, F., Hamr, P., ... & Collen, B.
683 (2015). Multiple drivers of decline in the global status of freshwater crayfish (Decapoda:
684 Astacidea). *Philos. Trans. R. Soc. B: Biol. Sci.* **370**, 20140060.

685 Sayer, C. et al. (2024). Under waster, under threat: Extinction risk of the world's freshwater
686 fauna. Accepted in Nature.

687 Secretariat of the Convention on Biological Diversity. (2022). Kunming-Montreal Global
688 Biodiversity Framework. [https://www.cbd.int/doc/decisions/cop-15/cop-15-dec-04-](https://www.cbd.int/doc/decisions/cop-15/cop-15-dec-04-en.pdf)
689 [en.pdf](https://www.cbd.int/doc/decisions/cop-15/cop-15-dec-04-en.pdf).

690 Steel, M., Mimoto, A., & Mooers, A. Ø. (2007). Hedging our bets: the expected contribution of
691 species to future phylogenetic diversity. *Evol. Bioinform.* **3**, 117693430700300024.

692 Stein, R. W., Mull, C. G., Kuhn, T. S., Aschliman, N. C., Davidson, L. N., Joy, J. B., Smith, G. J., Dulvy, N.
693 K. & Mooers, A. O. (2018). Global priorities for conserving the evolutionary history of
694 sharks, rays and chimaeras. *Nat. Ecol. Evol.* **2**, 288-298.

695 Stern, D. B., Breinholt, J., Pedraza-Lara, C., López-Mejía, M., Owen, C. L., Bracken-Grissom, H.,
696 Fetzner Jr, J. W. , Crandall, K. A. (2017). Phylogenetic evidence from freshwater
697 crayfishes that cave adaptation is not an evolutionary dead-end. *Evolution*, **71**, 2522-
698 2532.

699 Taylor, C. A. (2018). Two new species of freshwater crayfish of the genus *Faxonius* (Decapoda:
700 Cambaridae) from the Ozark Highlands of Arkansas and Missouri. *Zootaxa*. **4399**, 491-
701 520.

702 Taylor, C. A., DiStefano, R. J., Larson, E. R., & Stoeckel, J. (2019). Towards a cohesive strategy for
703 the conservation of the United States' diverse and highly endemic crayfish fauna.
704 *Hydrobiologia*. **846**, 39-58.

705 Toon, A., Pérez-Losada, M., Schweitzer, C. E., Feldmann, R. M., Carlson, M., & Crandall, K. A.
706 (2010). Gondwanan radiation of the Southern Hemisphere crayfishes (Decapoda:
707 Parastacidae): evidence from fossils and molecules. *J. Biogeogr.* **37**, 2275-2290.

708 Trull, N., Böhm, M., & Carr, J. (2018). Patterns and biases of climate change threats in the IUCN
709 Red List. *Conserv. Biol.* **32**, 135-147.

710 Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., Glidden, S.,
711 Bunn, S. E., Sullivan, C. A., Liermann, C. R. & Davies, P. M. (2010). Global threats to human
712 water security and river biodiversity. *Nature*, **467**, 555-561.

713 Whiting, A. S., Lawler, S. H., Horwitz, P., & Crandall, K. A. (2000). Biogeographic regionalization
714 of Australia: assigning conservation priorities based on endemic freshwater crayfish
715 phylogenetics. *Anim. Conserv.* **3**, 155-163.

716 WWF. (2024). Living Planet Report 2024 – A System in Peril. WWF, Gland, Switzerland.

717