"Essential oils" AND ("Phytochemical profile" OR "Phytochemical composition" OR "Chemical characterization" OR "Chemical components" OR "Chemical composition" OR "Chemical constituents") AND TABLE

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T1/ | TABLE 1<br>Major chemical constituents of clove plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Amelia et al. (2017) / From clove bud / Clove from Java, Indonesia: eugenol 55.60%, eugenyl acetate 20.54%, caryophyllene 14.84%, α-humelene 2.75%, β-elemene 0.04%, α-cadinene 0.05%, ledol 0.06%; from Manodo, Indonesia: eugenol 74.64%, caryophylene 12.79%, eugenyl acetate 8.70%, α-humelene 1.53%, β-gurjunene 0.04%, γ-cadinene 0.03%, humelene oxide 0.05%; comparison with other papers, eugenol, 47.60–89.20%, caryophyllene, trace level to 35.40%, eugenyl acetate, 1.20–20.54%, α-humulene, trace level to 2.75%<br>Chaieb et al. (2007) / From clove flower bud essential oil / Eugenol 88.59%, eugenyl acetate 5.62%, β-caryophyllene 1.39%, 2-heptanone 0.93%, ethyl hexanoate 0.66%, α-humulene 0.20%, calacorene 0.11%, calamenene 0.11%<br>Jirowetz et al. (2006) / From clove leaf essential oil / Eugenol 76.8%, β-caryophyllene 17.4%, α-humulene 2.1%, eugenyl acetate 1.2%, caryophyllene oxide 0.4%, methylchavicol 0.2%<br>Uddin et al. (2017) / From clove buds oil in Bangladesh / m-eugenol 69.44%, eugenyl acetate 10.79%, caryophyllene 6.8%, tyranton 7.78%, trace amounts of other constituents <1%<br>Open in a separate window | **Data source**<br>**Source of samples**<br>**Major chemical constituents**<br>Amelia et al. (2017)<br>From clove bud<br>Clove from Java, Indonesia: eugenol 55.60%, eugenyl acetate 20.54%, caryophyllene 14.84%, α-humelene 2.75%, β-elemene 0.04%, α-cadinene 0.05%, ledol 0.06%; from Manodo, Indonesia: eugenol 74.64%, caryophylene 12.79%, eugenyl acetate 8.70%, α-humelene 1.53%, β-gurjunene 0.04%, γ-cadinene 0.03%, humelene oxide 0.05%; comparison with other papers, eugenol, 47.60–89.20%, caryophyllene, trace level to 35.40%, eugenyl acetate, 1.20–20.54%, α-humulene, trace level to 2.75%<br>Chaieb et al. (2007)<br>From clove flower bud essential oil<br>Eugenol 88.59%, eugenyl acetate 5.62%, β-caryophyllene 1.39%, 2-heptanone 0.93%, ethyl hexanoate 0.66%, α-humulene 0.20%, calacorene 0.11%, calamenene 0.11%<br>Jirowetz et al. (2006)<br>From clove leaf essential oil<br>Eugenol 76.8%, β-caryophyllene 17.4%, α-humulene 2.1%, eugenyl acetate 1.2%, caryophyllene oxide 0.4%, methylchavicol 0.2%<br>Uddin et al. (2017)<br>From clove buds oil in Bangladesh<br>m-eugenol 69.44%, eugenyl acetate 10.79%, caryophyllene 6.8%, tyranton 7.78%, trace amounts of other constituents <1% |
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T2/ | TABLE 2<br>Major chemical constituents of eucalyptus plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Barbosa et al. (2016) / Review paper / Major chemical constituents in the extracts from leaves are terpenes and terpenoids at various concentrations. The concentrations vary among the species of *Eucalyptus* and the location they were harvested. 1,8-Cineole most often is the most major chemical constituent in *E. camaldulensis*, *E. cinerea*, and *E. globulus* (percentages vary from as low as almost 10% to as high as over 90.0%), whereas citronellal is the major chemical constituent in *E. citriodora* (percentages varied from as low as almost 20% to as high as over 90%). α-Pinene is the major chemical constituent in *E. saligna* (from 24.4 to 45.1%), depending on the location the plants are harvested<br>Dogan et al. (2017) / Leaf Fruit / 1,8-cineole 14.1%, p-cymene 42.1%, α-pinene 12.7%, α-terpineol 10.7%, limonene 5.5%, borneol L 5.5%, spathulenol 3.2% 1,8-cineole 34.5%, p-cymene 30.0%, α-terpineol 15.1%, α-pinene 9.0%, borneol L 5.3%, γ-terpinene 5.1%, spathulenol none, limonene none<br>Salehi et al. (2019b) / Review paper / Percentages are not shown<br>Sebei et al. (2015) / Review paper / Comparing the percentages among species, 1,8-cineole (49.07–83.50%) and α-pinene (1.27–26.35%) were the two major chemical constituents<br>Open in a separate window | **Data source**<br>**Source of samples**<br>**Major chemical constituents**<br>Barbosa et al. (2016)<br>Review paper<br>Major chemical constituents in the extracts from leaves are terpenes and terpenoids at various concentrations. The concentrations vary among the species of *Eucalyptus* and the location they were harvested. 1,8-Cineole most often is the most major chemical constituent in *E. camaldulensis*, *E. cinerea*, and *E. globulus* (percentages vary from as low as almost 10% to as high as over 90.0%), whereas citronellal is the major chemical constituent in *E. citriodora* (percentages varied from as low as almost 20% to as high as over 90%). α-Pinene is the major chemical constituent in *E. saligna* (from 24.4 to 45.1%), depending on the location the plants are harvested<br>Dogan et al. (2017)<br>Leaf Fruit<br>1,8-cineole 14.1%, p-cymene 42.1%, α-pinene 12.7%, α-terpineol 10.7%, limonene 5.5%, borneol L 5.5%, spathulenol 3.2% 1,8-cineole 34.5%, p-cymene 30.0%, α-terpineol 15.1%, α-pinene 9.0%, borneol L 5.3%, γ-terpinene 5.1%, spathulenol none, limonene none<br>Salehi et al. (2019b)<br>Review paper<br>Percentages are not shown<br>Sebei et al. (2015)<br>Review paper<br>Comparing the percentages among species |
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T3/ | TABLE 3<br>Major chemical constituents of lemon plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Klimek-Szczykutowicz et al. (2020) / Essential oil of pericarp Essential oil of leaf / Limonene 69.9%, p-menta-3,8-diene 18.0%, β-pinene 11.2%, γ-terpinene 8.21%, myrcene 4.4%, sabinene 3.9%, geranial 2.9%, neral 1.5%, linalool 1.41%, α-pinene 1.1%, α-thujene 1.1%, β-bisabolene 0.5%, (E)-β-ocimene 0.4%, geraniol 0.2%, β-caryophyllene 0.2% limonene 31.5%, sabinene 15.9%, citronellal 11.6%, linalool 4.6%, neral 4.5%, geranial 4.5%, (E)-β-ocimene 3.9%, myrcene 2.9%, citronellol 2.3%, β-caryophyllene 1.7%, terpinen-4-ol 1.4%<br>Luciardi et al. (2021) / Essential oils / Limonene 59.14%, γ-terpinene 10.48%, β-pinene 15.41%, sabinene 1.76%, β-myrcene 1.65%, α-pinene 1.64%<br>Pucci et al. (2020) / The whole fractions / Limonene 67.1%, β-pinene 10.88%, γ-terpinene 9.32%, α-pinene 1.81%, geranial 1.72%, sabinene 1.83%, myrcene 1.57%<br>Open in a separate window | **Data source**<br>**Source of samples**<br>**Major chemical constituents**<br>Klimek-Szczykutowicz et al. (2020)<br>Essential oil of pericarp Essential oil of leaf<br>Limonene 69.9%, p-menta-3,8-diene 18.0%, β-pinene 11.2%, γ-terpinene 8.21%, myrcene 4.4%, sabinene 3.9%, geranial 2.9%, neral 1.5%, linalool 1.41%, α-pinene 1.1%, α-thujene 1.1%, β-bisabolene 0.5%, (E)-β-ocimene 0.4%, geraniol 0.2%, β-caryophyllene 0.2% limonene 31.5%, sabinene 15.9%, citronellal 11.6%, linalool 4.6%, neral 4.5%, geranial 4.5%, (E)-β-ocimene 3.9%, myrcene 2.9%, citronellol 2.3%, β-caryophyllene 1.7%, terpinen-4-ol 1.4%<br>Luciardi et al. (2021)<br>Essential oils<br>Limonene 59.14%, γ-terpinene 10.48%, β-pinene 15.41%, sabinene 1.76%, β-myrcene 1.65%, α-pinene 1.64%<br>Pucci et al. (2020)<br>The whole fractions<br>Limonene 67.1%, β-pinene 10.88%, γ-terpinene 9.32%, α-pinene 1.81%, geranial 1.72%, sabinene 1.83%, myrcene 1.57% |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T4/ | **TABLE 4**<br>Major chemical constituents of rose plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Akram et al. (2019) / Oil Absolute rose Hydrosol / Geraniols (5.5–18%), β-citronellol (14.5–47.5%), nonadecane (10.5–40.5%) heneicosane, ethanol (0–13.43%), geraniol (3.71%), citronellol (9.91%), nonadecane (4.35%), phenylethylalcohol (78.38%) nerol (16.12%), phenylethyl alcohol (23.74%), citronellol (29.44%), geraniol (30.74%)<br>Mileva et al. (2021) / Oils / There are large differences among geological differences and species differences; geraniol (17.60–30.98%), nerol (4.36–10.10%), citronellol (9.22–28.72%), n-nonadecane (8.10–22.67%), n-heneicosane (5.00–10.21%)<br>Ryu et al. (2020) / Oils / There are large differences among breeds (mutants) of roses<br>Verma et al. (2010) / From bud, half bloom, full bloomEssential oils / Phenyl ethyl alcohol (PEA) 66.2–80.7%, other major chemical constituent, although much less in the percentages, were citronellol (1.8–5.5%) and geraniol (4.4–7.9%). Major chemical constituents were citronellol (15.9–35.3%), geraniol (8.3–30.2%), nerol (4.0–9.6%), nonadecane (4.5–16.0%), and<br>Open in a separate window | TABLE 4<br>Major chemical constituents of rose plants and essential oils. | **Data source**<br>**Source of samples**<br>**Major chemical constituents**<br>Akram et al. (2019)<br>Oil Absolute rose Hydrosol<br>Geraniols (5.5–18%), β-citronellol (14.5–47.5%), nonadecane (10.5–40.5%) heneicosane, ethanol (0–13.43%), geraniol (3.71%), citronellol (9.91%), nonadecane (4.35%), phenylethylalcohol (78.38%) nerol (16.12%), phenylethyl alcohol (23.74%), citronellol (29.44%), geraniol (30.74%)<br>Mileva et al. (2021)<br>Oils<br>There are large differences among geological differences and species differences; geraniol (17.60–30.98%), nerol (4.36–10.10%), citronellol (9.22–28.72%), n-nonadecane (8.10–22.67%), n-heneicosane (5.00–10.21%)<br>Ryu et al. (2020)<br>Oils<br>There are large differences among breeds (mutants) of roses<br>Verma et al. (2010)<br>From bud, half bloom, full bloomEssential oils<br>Phenyl ethyl alcohol (PEA) 66.2–80.7%, other major chemical constituent, although much less in the percentages, were citronellol (1.8–5.5%) and geraniol (4.4–7.9%). Major chemical constituents were citronellol (15.9–35.3%), geraniol (8.3–30.2%), nerol (4.0–9.6%), nonadecane (4.5–16.0%), and not PEA (0.6–2.9%) |
| https://europepmc.org/article/MED/35840624 | Ovicidal and repellent activities of several plant essential oils against Periplaneta americana L. and enhanced activities from their combined formulation. | https://europepmc.org/articles/PMC9287551/table/Tab1/ | Item / Compound / R[a] / K[b] / Peak area (%), (average of 3 runs ± SD) C. citratus / C. verum / E. globulus / I. verum / Z. limonella / I[c]<br>1. Acetoin 680 680 – – – 0.41 ± 0.02 – M,R<br>2. α-Thujene 928 930 – – – 0.21 ± 0.01 2.31 ± 0.54 M,R<br>3. α-Pinene 932 933 3.52 ± 0.11 0.91 ± 0.02 5.01 ± 0.03 0.32 ± 0.02 1.83 ± 0.08 M,R<br>4. Camphene 946 946 – 0.52 ± 0.01 0.32 ± 0.02 – – M,R<br>5. Sabinene 977 977 – – 3.75 ± 0.06 – 4.52 ± 0.05 M,R<br>6. B-Mycene 990 991 – 0.43 ± 0.03 0.31 ± 0.01 – 0.32 ± 0.02 M,R<br>7. α-Phellandrene 1002 1003 – 0.32 ± 0.01 1.67 ± 0.10 – 1.31 ± 0.07 M,R<br>8. α-Terpinene 1002 1002 0.22 ± 0.08 – – 0.31 ± 0.01 4.82 ± 0.18 M,R<br>9. Benzene 1011 1009 – 14.15 ± 0.25 8.32 ± 0.38 – 8.21 ± 0.76 M,R<br>10. 1,8-Cineole 1024 1025 10.93 ± 1.01 0.61 ± 0.02 45.82 ± 1.01 0.52 ± 0.03 – M,R<br>11. Limonene 1029 1029 – – – 1.85 ± 0.09 28.13 ± 1.37 M,R<br>12. ϒ-Terpinene 1051 1050 0.26 ± 0.14 – 20.13 ± 1.44 – 7.82 ± 0.82 M,R<br>13. Butanoic acid 1085 1086 – – 0.52 ± 0.06 – – M,R<br>14. Linalool 1086 1086 0.86 ± 0.07 – – – 1.15 ± 0.07 M,R<br>15. Terpinolene 1089 1088 – – 1.12 ± 0.61 0.23 ± 0.01 2.23 ± 0.83 M,R<br>16. D-Fenchyl alcohol 1110 1110 – – 0.55 ± 0.04 – – M,R<br>17. *trans*-Pinocaveol 1139 1139 – – 0.72 ± 0.05 – – M,R<br>18. Borneol 1146 1147 – 1.23 ± 0.08 0.81 ± 0.01 – – M,R<br>19. Terpinen-4-ol 1165 1166 – – 3.21 ± 0.23 – 24.13 ± 1.65 M,R<br>20. *trans*-Carveol 1204 1204 – – 0.23 ± 0.01 – 1.73 ± 0.82 M,R<br>21. Myrtenol 1214 1214 – – 0.71 ± 0.02 – – M,R<br>22. Neral 1217 1217 25.72 ± 1.97 – – – – M,R<br>23. Cinnamaldehyde 1222 1222 – 75.23 ± 2.17 – – – M,R<br>24. *p*-Anisaldehyde 1223 1223 – – – 1.63 ± 0.02 – M,R<br>25. Carvone 1230 1231 – – – – 2.47 ± 0.08 M,R<br>26. Geraniol 1237 1237 5.71 ± 0.82 – – – – M,R<br>27 Geranial 1247 1247 46.33 ± 1.32 – – – – M,R<br>28. α-Terpineol 1280 1279 – – 2.85 ± 0.49 – 6.23 ± 1.12 M,R<br>29. *trans*-Anethole 1283 1285 – – – 92.24 ± 1.83 – M,R<br>30. Eugenol 1356 1355 – – – 0.84 ± 0.03 – M,R<br>31. Geranyl acetate 1379 1380 3.87 ± 0.33 – – – – M,R<br>32. Copaene 1380 1381 – 2.02 ± 0.15 – – – M,R<br>33. Cinnamic acid 1460 1462 – 0.33 ± 0.09 – – – M,R<br>34. α-Gurjunene 1527 1529 – – 0.86 ± 0.02 – – M,R<br>35. Epiglobulol 1563 1564 – – 0.44 ± 0.01 – – M,R<br>36. Caryophyllene 1578 1580 1.17 ± 0.87 – – – 0.54 ± 0.02 M,R<br>37. Cadalene 1659 1659 – 0.32 ± 0.08 – – – M,R<br>Total (%) 97.40 98.60 97.35 98.56 97.75 –<br>Yield (%) 1.15 1.10 1.21 3.23 9.27 – | **Table 1**<br>Chemical constituents of *Cymbopogon citratus*, *Cinnamomum verum*, *Eucalyptus globulus*, *Illicium verum*, and *Zanthoxylum limonella* and their percentage yield in the total chemical profile. [a]R = Retention index of a chemical constituent determined with an HP-5 MS column and compared with the retention indices of standard alkanes ($C_7$-$C_{30}$) for identity verification. [b]K = Kovat retention index from NIST17[62]. [c]I = Identified and confirmed by mass spectrum (M) matching with chemicals in the computer mass library of Adams[61] and by retention index (R) matching with those reported in NIST17[62]. | |
| https://europepmc.org/article/MED/35807419 | Effects of Frankincense Compounds on Infection, Inflammation, and Oral Health. | https://europepmc.org/articles/PMC9268443/table/molecules-27-04174-t002/ | **Table 2**<br>Prominent features of compounds in frankincense resin and oils.<br>Frankincense Type / Country of Origin / Source / Chemical/Active Component / References<br>*Boswellia dalzielii* / Chad, Mali, Nigeria / Hydro-distilled leaf essential oil / δ-3-carene (27.7%), α-pinene (15.2%), *p*-cymene (9.5%), β-phellandrene (8.5%), isolongifolene (6.2%), and myrcene (5.7%). / [65]<br>*Boswellia dalzielii* / Nigeria / Hydro-distilled leaf essential oil / α-pinene (45.7%) and α-terpinene (11.5%), trans-sabinene hydrate (4.6%), cis-p-menth-2-en1-ol (2.9%), α-campholenal (2.7%), caryophyllene oxide, and α-phellandrene (2.3%) / [64]<br>*B. carterii* / Somalia / Gum resin / Esters (62.1%), 1-octyl acetate being predominant (60.0%). Alcohols amount to 15.4%, 1-octanol being the major component (12.7%), and diterpene constituents amount to 7.1%, including cembrene (1.4%), isocembrene (1.8%), incensole (2.7%), and isoincensole (0.8) and a mixture of monoterpene hydrocarbons which amounts to 9.9%. / [58]<br>*B. rivae* / Ethiopia / Essential oil from frankincense / α-Pinene (36.1–67.7%), δ -3-carene (12.2%), and limonene (12.0%). / [66]<br>*B. neglecta* / Ethiopia / Essential oil from frankincense / α-Pinene (36.1–67.7%), terpinen-4-ol (11.3%). / [66]<br>Open in a separate window | **Table 2**<br>Prominent features of compounds in frankincense resin and oils. | **Frankincense Type**<br>**Country of Origin**<br>**Source**<br>**Chemical/Active Component**<br>**References**<br>*Boswellia dalzielii*<br>Chad, Mali, Nigeria<br>Hydro-distilled leaf essential oil<br>δ-3-carene (27.7%), α-pinene (15.2%), *p*-cymene (9.5%), β-phellandrene (8.5%), isolongifolene (6.2%), and myrcene (5.7%).<br>[65]<br>*Boswellia dalzielii*<br>Nigeria<br>Hydro-distilled leaf essential oil<br>α-pinene (45.7%) and α-terpinene (11.5%), trans-sabinene hydrate (4.6%), cis-p-menth-2-en1-ol (2.9%), α-campholenal (2.7%), caryophyllene oxide, and α-phellandrene (2.3%)<br>[64]<br>*B. carterii*<br>Somalia<br>Gum resin<br>Esters (62.1%), 1-octyl acetate being predominant (60.0%). Alcohols amount to 15.4%, 1-octanol being the major component (12.7%), and diterpene constituents amount to 7.1%, including cembrene (1.4%), isocembrene (1.8%), incensole (2.7%), and isoincensole (0.8) and a mixture of monoterpene hydrocarbons which amounts to 9.9%.<br>[58]<br>*B. rivae*<br>Ethiopia |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35631552 | Essential Oils of Gardenia jasminoides J. Ellis and Gardenia jasminoides f. longicarpa Z.W. Xie & M. Okada Flowers: Chemical Characterization and Assessment of Anti-Inflammatory Effects in Alveolar Macrophage. | https://europepmc.org/articles/PMC9145545/table/pharmaceutics-14-00966-t001/ | **Table 1**<br>The main constituents of the essential oils of the two gardenia flowers.<br>The table lists those components whose peak area is >0.5% of the total peak area; "-" means not detected or the peak area <0.5%. | **Table 1**<br>The main constituents of the essential oils of the two gardenia flowers. | The table lists those components whose peak area is >0.5% of the total peak area; "-" means not detected or the peak area <0.5%. |
| https://europepmc.org/article/MED/35011355 | Phytochemical Profile, Antimicrobial, Cytotoxic, and Antioxidant Activities of Fresh and Air-Dried Satureja nabateorum Essential Oils. | https://europepmc.org/articles/PMC8746579/table/molecules-27-00125-t001/ | **Table 1**<br>Chemical constituents of the essential oils from fresh and dry *Satureja nabateorum*. | **Table 1**<br>Chemical constituents of the essential oils from fresh and dry *Satureja nabateorum*. | |
| https://europepmc.org/article/MED/35893636 | Analysis of Volatile Constituents in Curcuma Species, viz. C. aeruginosa, C. zedoaria, and C. longa, from Nepal. | https://europepmc.org/articles/PMC9332366/table/plants-11-01932-t001/ | **Table 1**<br>Individual constituents of *Curcuma longa* essential oil. | **Table 1**<br>Individual constituents of *Curcuma longa* essential oil. | $RI_{calc}$ = Retention index determined with respect to homologous series of n-alkanes on a ZB-5 column. $RI_{db}$ = Retention index from the database [33,34]. 't' indicate trace (< 0.05%). |

**Table 1**

The main constituents of the essential oils of the two gardenia flowers.

| No. | Retention Time (min) | Name | Peak Area (%) | |
|---|---|---|---|---|
| | | | GJE | GJLE |
| 1 | 14.35 | Linalool | 34.65 | 6.56 |
| 2 | 17.19 | α-Terpineol | 6.27 | 2.40 |
| 3 | 18.41 | Nerol | 1.96 | 0.64 |
| 4 | 19.66 | Geraniol | 5.79 | 1.93 |
| 5 | 21.78 | cis-3-Hexenyl tiglate | 3.13 | 8.02 |
| 6 | 21.99 | Hexyl tiglate | 2.39 | 1.23 |
| 7 | 28.00 | α-Farnesene | 10.24 | 32.45 |
| 8 | 28.34 | Octyl (E)-2-methylbut-2-enoate | 0.70 | 0.52 |
| 9 | 29.50 | (+)-trans-Nerolidol | - | 0.53 |
| 10 | 29.66 | 3-Hexen-1-ol, benzoate, (Z)- | - | 1.76 |
| 11 | 29.82 | Benzoic acid, hexyl ester | - | 0.66 |
| 12 | 29.92 | (3E,7E)-4,8,12-Trimethyltrideca-1,3,7,11-tetraene | - | 0.90 |
| 13 | 31.79 | tau-Cadinol | 1.77 | 2.40 |
| 14 | 32.17 | α-Cadinol | 0.68 | 0.74 |
| 15 | 32.30 | α-Terpinyl acetate | 1.19 | - |
| 16 | 33.51 | Geranyl angelate | 1.89 | 0.55 |
| 17 | 36.57 | 8-Hydroxylinalool | 1.67 | 1.23 |
| 18 | 40.67 | Cembrene A | 5.77 | 0.97 |
| 19 | 40.86 | Verticillol | 1.01 | 0.97 |
| 20 | 41.55 | n-Hexadecanoic acid | 1.33 | 5.70 |
| 21 | 42.10 | 3,7,11,15-Tetramethylhexadeca-1,6,10,14-tetraen-3-ol | - | 1.17 |
| 22 | 45.10 | Benzoic acid, [(E,E)-3,7,11-trimethyl-2,6,10-dodecatrien-1-yl] ester | - | 1.11 |
| 23 | 45.87 | trans-Geranylgeraniol | - | 1.03 |
| 24 | 45.69 | 9,12-Octadecadienoic acid (Z,Z)- | - | 1.44 |
| 25 | 48.44 | Tricosane | 1.93 | 0.90 |
| 26 | 52.66 | Pentacosane | 2.96 | 3.68 |
| 27 | 56.37 | Heptacosane | - | 0.53 |
| 28 | 58.77 | Squalene | - | 0.64 |
| Sum | | | 85.33 | 72.64 |

The table lists those components whose peak area is >0.5% of the total peak area; "-" means not detected or the peak area <0.5%.

**Table 1**

Chemical constituents of the essential oils from fresh and dry *Satureja nabateorum*.

| | Component | RI | Dry, % | Fresh % |
|---|---|---|---|---|
| 1 | α-Thujene | 918 | 1.24 ± 0.01 | 1.06 ± 0.12 |
| 2 | α-Pinene | 930 | 0.65 ± 0.002 | 0.59 ± 0.008 |
| 3 | Camphene | 952 | 0.23 ± 0.006 | 0.37 ± 0.005 |
| 4 | Sabinene | 972 | 0.08 ± 0.001 | 0.15 ± 0.001 |
| 5 | β-Pinene | 979 | 0.18 ± 0.001 | 0.20 ± 0.02 |
| 6 | Isolimonene | 984 | - | 0.41 ± 0.002 |
| 7 | Myrcene | 991 | 1.47 ± 0.01 | 0.69 ± 0.001 |
| 8 | δ-2-Carene | 1003 | 0.23 ± 0.002 | 0.25 ± 0.02 |
| 9 | α-Terpinene | 1019 | 2.81 ± 0.01 | 2.68 ± 0.98 |
| 10 | p-Cymene | 1023 | 15.02 ± 1.02 | 11.51 ± 0.97 |
| 11 | Sylvesterine | 1029 | 0.68 ± 0.009 | 0.60 ± 0.001 |
| 12 | Ocimene | 1033 | 0.60 ± 0.008 | 0.51 ± 0.006 |
| 13 | γ-Terpinene | 1061 | 21.15 ± 1.05 | 20.65 ± 1.12 |
| 14 | Terpinolene | 1087 | - | 0.09 ± 0.002 |
| 15 | Linalool | 1098 | 1.69 ± 0.05 | 0.78 ± 0.006 |
| 16 | cis-Thujone | 1103 | - | 0.17 ± 0.003 |
| 17 | 3-Terpinen-1-ol | 1133 | - | 0.13 ± 0.001 |
| 18 | (-)-Camphor | 1144 | - | 0.69 ± 0.006 |
| 19 | Not identified | 1148 | - | 0.19 ± 0.003 |
| 20 | Terpinen-4-ol | 1173 | 0.69 ± 0.006 | 0.17 ± 0.002 |
| 21 | Nerol | 1228 | - | 0.15 ± 0.001 |
| 22 | Thymol methylether | 1235 | 1.21 ± 0.01 | 0.76 ± 0.005 |
| 23 | Carvacrol methylether | 1242 | 1.04 ± 0.01 | 3.44 ± 0.01 |
| 24 | Linalool acetate | 1256 | - | 0.15 ± 0.001 |
| 25 | Geranial | 1262 | 0.24 ± 0.002 | 2.91 ± 0.01 |
| 26 | Neomenthyl acetate | 1266 | 0.24 ± 001 | - |
| 27 | Isobornyl acetate | 1278 | - | 1.80 ± 0.01 |
| 28 | Thymol | 1292 | 46.07 ± 1.1 | 40.64 ± 1.21 |
| 29 | α-Cubebene | 1345 | - | 0.89 ± 0.002 |
| 30 | α-Cupaene | 1377 | - | 0.25 ± 0.004 |
| 31 | β-Cubebene | 1383 | - | 0.25 ± 0.002 |
| 32 | β-Caryophyllene | 1424 | 2.78 ± 0.006 | 2.85 ± 0.02 |
| 33 | α-Caryophyllene | 1457 | 0.14 ± 0.003 | 0.16 ± 0.001 |
| 34 | Geranyl propanoate | 1471 | 0.38 ± 0.001 | 0.30 ± 0.01 |
| 35 | Muurolene | 1476 | 0.11 ± 0.006 | - |
| 36 | Germacrene D | 1482 | 0.54 ± 0.003 | 1.14 ± 0.07 |
| 37 | α-Muurolene | 1497 | 0.07 ± 0.001 | 0.13 ± 0.06 |
| 38 | Germacrene A | 1508 | 0.07 ± 0.007 | 0.36 ± 0.21 |
| 39 | γ-Cadinene | 1514 | 0.16 ± 0.006 | 0.12 ± 0.03 |
| 40 | δ-Cadinene | 1528 | 0.22 ± 0.001 | 0.09 ± 0.001 |
| 41 | Germacrene B | 1552 | 0.04 ± 0.001 | 0.26 ± 0.004 |
| 42 | Caryophellene oxide | 1580 | 0.07 ± 0.001 | 0.10 ± 0.002 |
| | Total | | 99.56 | 98.64 |
| | **Grouped Compounds** | | | |
| | Monoterpene hydrocarbons | | 47.58 | 47.20 |
| | Phenolic monoterpenes | | 48.32 | 44.84 |
| | Sesquiterpene hydrocarbon | | 3.66 | 6.60 |

**Table 1**

Individual constituents of *Curcuma longa* essential oil.

| $RI_{calc}$ | $RI_{db}$ | Compound Name | Area% of Constituents in *Curcuma longa* |
|---|---|---|---|
| 925 | 924 | α-Thujene | t |
| 933 | 931 | α-Pinene | 0.1 |
| 972 | 972 | Sabinene | t |
| 978 | 978 | β-Pinene | t |
| 989 | 989 | Myrcene | 0.1 |
| 1001 | 1000 | δ-2-Carene | t |
| 1007 | 1008 | α-Phellandrene | 0.2 |
| 1009 | 1009 | δ-3-Carene | 0.1 |
| 1017 | 1016 | α-Terpinene | 0.4 |
| 1020 | 1024 | *p*-Cymene | 0.1 |
| 1024 | 1028 | Limonene | 0.1 |
| 1025 | 1029 | β-Phellandrene | t |
| 1033 | 1031 | 1,8-Cineole | 1.3 |
| 1035 | 1034 | (*Z*)-β-Ocimene | t |
| 1045 | 1044 | (*E*)-β-Ocimene | t |
| 1057 | 1058 | γ-Terpinene | t |
| 1069 | 1071 | *cis*-Sabinene hydrate | t |
| 1085 | 1086 | Terpinolene | 7.2 |
| 1090 | 1091 | *p*-Cymenene | 0.1 |
| 1108 | 1111 | 1,3,8-*p*-Menthatriene | t |
| 1130 | 1132 | 1(7),3,8-*o*-Menthatriene | t |
| 1138 | 1143 | Epoxyterpinolene | t |
| 1146 | 1145 | Myrcenone | t |
| 1180 | 1180 | Terpinen-4-ol | t |
| 1182 | 1184 | Anethofuran | t |
| 1187 | 1187 | *p*-Cymen-8-ol | 0.2 |
| 1195 | 1195 | α-Terpineol | t |
| 1234 | 1237 | Turmeric dione | t |
| 1291 | 1292 | 2-Undecanone | t |
| 1365 | 1368 | (Z,Z)-Megastigma-4,6,8-triene | t |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35893636 | Analysis of Volatile Constituents in Curcuma Species, viz. C. aeruginosa, C. zedoaria, and C. longa, from Nepal. | https://europepmc.org/articles/PMC9332366/table/plants-11-01932-t002/ | **Table 2**<br>Individual constituents of *Curcuma aeruginosa* and *Curcuma zedoaria* essential oil.<br>$RI_{calc}$ = Retention index determined with respect to homologous series of n-alkanes on a ZB-5 column. $RI_{db}$ = Retention index from the database [33,34]. 't' indicate trace (< 0.05%). | **Table 2**<br>Individual constituents of *Curcuma aeruginosa* and *Curcuma zedoaria* essential oil. | $RI_{calc}$ = Retention index determined with respect to homologous series of n-alkanes on a ZB-5 column. $RI_{db}$ = Retention index from the database [33,34]. 't' indicate trace (< 0.05%). |
| https://europepmc.org/article/MED/35893636 | Analysis of Volatile Constituents in Curcuma Species, viz. C. aeruginosa, C. zedoaria, and C. longa, from Nepal. | https://europepmc.org/articles/PMC9332366/table/plants-11-01932-t003/ | **Table 3**<br>Enantiomeric distributions of chiral terpenoids of *Curcuma aeruginosa*, *Curcuma zedoaria*, and *Curcuma longa* essential oil. | **Table 3**<br>Enantiomeric distributions of chiral terpenoids of *Curcuma aeruginosa*, *Curcuma zedoaria*, and *Curcuma longa* essential oil. | |
| https://europepmc.org/article/MED/35755940 | Antioxidant, Antidiabetic, and Antibacterial Potentials and Chemical Composition of Salvia officinalis and Mentha suaveolens Grown Wild in Morocco. | https://europepmc.org/articles/PMC9217590/table/tab5/ | **Table 5**<br>Chemical composition of the essential oils (EOs) of *Salvia officinalis* and *Mentha suaveolens*. | **Table 5**<br>Chemical composition of the essential oils (EOs) of *Salvia officinalis* and *Mentha suaveolens*. | |

**Table 2**

Individual constituents of *Curcuma aeruginosa* and *Curcuma zedoaria* essential oil.

| $RI_{calc}$ | $RI_{db}$ | Compound Name | Area % of Constituents in | |
|---|---|---|---|---|
| | | | *Curcuma zedoaria* | *Curcuma aeruginosa* |
| 882 | 885 | 3-Hepten-6-ol | 0.1 | - |
| 889 | 888 | 2-Heptanone | t | - |
| 894 | 894 | 2-Heptanol | 0.3 | 0.2 |
| 918 | 921 | Tricyclene | t | t |
| 925 | 924 | α-Thujene | t | t |
| 933 | 931 | α-Pinene | 0.8 | 0.2 |
| 945 | 945 | α-Fenchene | t | - |
| 949 | 948 | Camphene | 1.2 | 0.8 |
| 953 | 954 | Thuja-2,4(10)-diene | t | - |
| 972 | 972 | Sabinene | 0.2 | t |
| 978 | 978 | β-Pinene | 1.7 | 1.3 |
| 989 | 989 | Myrcene | 0.3 | 0.1 |
| 998 | 994 | 2-Octanol | t | t |
| 1007 | 1008 | α-Phellandrene | t | t |
| 1017 | 1016 | α-Terpinene | t | t |
| 1020 | 1024 | *p*-Cymene | t | t |
| 1024 | 1028 | Limonene | 1.1 | 0.2 |
| 1033 | 1031 | 1,8-Cineole | 19.6 | 1.2 |
| 1035 | 1034 | (*Z*)-β-Ocimene | t | - |
| 1045 | 1044 | (*E*)-β-Ocimene | t | - |
| 1057 | 1058 | γ-Terpinene | 0.1 | t |
| 1069 | 1071 | *cis*-Sabinene hydrate | t | - |
| 1085 | 1086 | Terpinolene | t | t |
| 1087 | 1090 | 2-Nonanone | 0.6 | t |
| 1090 | 1090 | (3*Z*)-Hexenyl methyl carbonate | t | - |
| 1099 | 1099 | Linalool | 0.2 | 0.3 |
| 1105 | 1103 | 2-Nonanol | 0.6 | 0.4 |
| 1112 | 1118 | *trans*-Thujone | - | t |
| 1119 | 1122 | *trans*-*p*-Mentha-2,8-dien-1-ol | t | - |
| 1122 | 1124 | *cis*-*p*-Menth-2-en-1-ol | t | - |

**Table 3**

Enantiomeric distributions of chiral terpenoids of *Curcuma aeruginosa*, *Curcuma zedoaria*, and *Curcuma longa* essential oil.

| Chiral Terpenoid Compound | *Curcuma aeruginosa* | *Curcuma zedoaria* | *Curcuma longa* |
|---|---|---|---|
| α-Pinene | (+)24.7: (–)75.3 | (+)43.5: (–)56.5 | (+)70.6: (–)29.4 |
| Camphene | (+)90.7: (–)9.3 | (+)92.3: (–)7.7 | - |
| β-Pinene | (+)59.1: (–)40.9 | (+)47.8: (–)52.2 | - |
| Sabinene | (+)0: (–)100 | - | - |
| α-Phellandrene | - | - | (+)100: (–)0 |
| *δ*-3-Carene | - | - | (+)100: (–)0 |
| Limonene | (+)0: (–)100 | - | (+)45.6: (–)54.4 |
| β-Phellandrene | - | - | (+)91.4: (–)8.6 |
| Linalool | (+)62.4: (–)37.6 | (+)55.3: (–)44.7 | - |
| Camphor | (+)99.8: (–)0.2 | (+)92.3: (–)7.7 | - |
| Bornyl acetate | (+)0: (–)100 | (+)0: (–)100 | - |
| Terpinen-4-ol | (+)0: (–)100 | - | - |
| *δ*-Elemene | (+)45.0: (–)55.0 | (+)33.7: (–)66.3 | - |
| α-Terpineol | (+)29.1: (–)70.9 | | (+)51.5: (–)48.5 |
| Borneol | (+)100: (–)0 | | - |
| β-Elemene | (+)0: (–)100 | (+)0: (–)100 | - |
| β-Caryophyllene | (+)0: (–)100 | (+)0: (–)100 | (+)0: (–)100 |
| Germacrene D | (+)100: (–)0 | (+)100: (–)0 | - |
| β-Bisabolene | - | - | (+)92.9: (–)7.1 |
| (*E*)-Nerolidol | - | - | (+)0: (–)100 |

**Table 5**

Chemical composition of the essential oils (EOs) of *Salvia officinalis* and *Mentha suaveolens*.

| No. | *Salvia officinalis* EO | | *Mentha suaveolens* EO | |
|---|---|---|---|---|
| | Compound | % | Compound | % |
| 1 | Thujone | 33.77 | Pulegone | 37.16 |
| 2 | Caryophyllene | 12.28 | Pyrazines | 33.81 |
| 3 | Humulene | 12.19 | Limonene | 11.19 |
| 4 | Camphor | 11.52 | Umbellulone | 6.09 |
| 5 | Naphthalene | 9.94 | Camphor | 4.27 |
| 6 | Eucalyptol | 8.11 | 3-Carene | 1.34 |
| 7 | *α*-Pinene | 3.31 | 2-Bornanone | 1.2 |
| 8 | *β*-Pinene | 1.8 | Menthone | 0.98 |
| 9 | *β*-Myrcene | 1.49 | 1-Octen-3-ol | 0.97 |
| 10 | Germacrene D | 1.36 | *α*-Pinene | 0.88 |
| 11 | Borneol | 1.18 | Thujone | 0.60 |
| 12 | Cyclopentane-3 | 0.96 | o-Menth-8-ene | 0.39 |
| 13 | *trans*-*β*-Ocimene | 0.60 | *trans*-Calamenene | 0.17 |
| 14 | *γ*-Terpinene | 0.51 | | |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35744919 | Chemical Characterization and Biological Activity of the Essential Oil from Araucaria brasiliensis Collected in Ecuador. | https://europepmc.org/articles/PMC9230380/table/molecules-27-03793-t001/ | **Table 1**<br>Chemical composition of *Araucaria brasiliensis* leaf essential oil (EO).<br>DB5-ms / HP-INNOWax<br>N° Compound LRI [a] LRI [b] $\bar{X}$ ± SD Ref. LRI [a] LRI [b] $\bar{X}$ ± SD Ref.<br>1 α-Pinene 931 932 9.99 ± 2.81 [27] 1062 1066 6.32 ± 1.01 [22]<br>2 Camphene 948 946 0.08 ± 0.02 [27] - - -<br>3 Sabinene 971 969 0.12 ± 0.04 [27] 1122 1121 0.20 ± 0.05 [23]<br>4 β-Pinene 976 974 3.34 ± 0.73 [27] 1111 1112 4.68 ± 1.02 [23]<br>5 Myrcene 989 988 11.02 ± 1.97 [27] 1168 1170 13.41 ± 0.98 [28]<br>6 α-Phellandrene 1006 1002 0.16 ± 0.05 [27] - - -<br>7 α-Terpinene 1016 1014 0.07 ± 0.02 [27] 1176 1185 0.06 ± 0.01 [29]<br>8 β-Phellandrene 1030 1025 1.90 ± 0.39 [27] 1209 1210 1.25 ± 0.27 [30]<br>9 γ-Terpinene 1057 1054 0.08 ± 0.01 [27] 1244 1245 0.11 ± 0.02 [31]<br>10 Terpinolene 1084 1086 0.07 ± 0.02 [27] 1281 1280 0.11 ± 0.03 [30]<br>11 Aromadendrene 1374 1374 0.07 ± 0.03 [27] 1594 1604 0.13 ± 0.03 [32]<br>12 Guaia-6,9-diene 1436 1439 0.07 ± 0.03 [27] 1597 1594 0.15 ± 0.03 [33]<br>13 γ-Muurolene 1440 1442 0.07 ± 0.02 [27] 1679 1676 0.18 ± 0.04 [34]<br>14 Germacrene D 1473 1478 1.11 ± 0.13 [27] 1698 1700 1.51 ± 0.12 [30]<br>15 Viridiflorene 1479 1480 0.11 ± 0.05 [27] 1685 1679 0.14 ± 0.03 [35]<br>16 Bicyclogermacrene 1489 1496 3.14 ± 0.21 [27] 1725 1722 4.08 ± 0.38 [36]<br>17 δ-Cadinene 1493 1500 0.15 ± 0.02 [27] 1750 1750 0.29 ± 0.04 [30]<br>18 β-Copaen-4-α-ol 1511 1513 0.18 ± 0.16 [27] - - -<br>19 Viridiflorol 1516 1522 0.19 ± 0.04 [27] 2075 2069 0.16 ± 0.01 [37]<br>20 Globulol 1556 1559 0.14 ± 0.04 [27] 2066 2057 0.27 ± 0.09 [37]<br>21 α-Muurolol 1560 1575 0.12 ± 0.04 [27] - - -<br>22 *epi*-α-Cadinol 1567 1567 0.15 ± 0.02 [27] -<br>23 Octadec-1-ene 1574 1590 0.10 ± 0.03 [27] - - -<br>24 Rosa-5,15-diene (Rimuene) 1592 1592 5.87 ± 1.01 [27] 2240 2255 8.00 ± 0.24 [38]<br>25 Beyerene 1642 1644 26.08 ± 3.65 [27] 2186 - 25.78 ± 1.55<br>26 Pimaradiene 1653 1638 0.24 ± 0.04 [27] - - -<br>27 Sandaracopimara-8(14),15-diene 1795 1789 4.47 ± 0.84 [27] - - -<br>28 Sclarene 1902 1900 0.54 ± 0.15 [27] 2262 - 0.52 ± 0.04<br>29 Kaur-15-ene 1933 1933 0.60 ± 0.11 [27] 2280 - 0.76 ± 0.03<br>30 13-*epi*-Manool oxide 1939 1931 2.31 ± 1.07 [27] 2360 2376 2.17 ± 0.16 [32]<br>31 Kaurene 1950 1948 24.86 ± 2.21 [27] 2386 2399 29.24 ± 0.97 [34]<br>32 Abietadiene 1970 1968 0.14 ± 0.02 [27] 2450 2450 0.14 ± 0.01 [32]<br>33 Phyllocladanol 2000 1997 0.22 ± 0.01 [27] - - -<br>Monoterpene hydrocarbons (%) 12.55 21.05<br>Sesquiterpene hydrocarbons (%) 6.10 6.72<br>Oxygenated sesquiterpenes (%) 0.93 0.87<br>Diterpenes hydrocarbons (%) 65.41 68.16<br>Oxygenated diterpenes (%) 3.12 2.17<br>Others (%) 0.05 0.10<br>Total (%) 88.16 99.07<br>LRI [a] = calculated linear retention index; LRI [b] = linear retention index from reference; Ref. = reference; $\bar{X}$ ± SD = percentage in the oil and standard deviation; both values were calculated as the means of three determinations; (-) unidentified compound on HP-INNOWax column. | **Table 1**<br>Chemical composition of *Araucaria brasiliensis* leaf essential oil (EO). | LRI [a] = calculated linear retention index; LRI [b] = linear retention index from reference; Ref. = reference; $X$ ± SD = percentage in the oil and standard deviation; both values were calculated as the means of three determinations; (-) unidentified compound on HP-INNOWax column. |
| https://europepmc.org/article/MED/35889245 | Essential Oil of the Plants Growing in the Brazilian Amazon: Chemical Composition, Antioxidants, and Biological Applications. | https://europepmc.org/articles/PMC9318482/table/molecules-27-04373-t001/ | **Table 1**<br>Major chemical constituents (≥3.00%) found in the essential oils of the Amazon.<br>Species / Family / Extraction Method / Compounds / References<br>*Anaxagorea brevipes* (leaves) — Annonaceae — HD — β-eudesmol (13.16%), α-eudesmol (13.05%), γ-eudesmol (7.54%), guaiol (5.12%), caryophyllene oxide (4.18%) and β-bisabolene (4.10%) — [15]<br>*Aniba duckei* (Synonym: *A. rosaeodora*) (leaves and thin branches) — Lauraceae — HD — linalool (89.34%) — [16]<br>*A. parviflora* (Aerial parts) — Lauraceae — HD — linalool (45.0%) — [17]<br>*A. parviflora* (branches) — Lauraceae — HD — γ-eudesmol (16.80%), (*E*)-caryophyllene (15.70%), linalool (12.40%), β-phellandrene (6.7%), and bicyclogermacrene (6.00%) — [18]<br>*A. parviflora* (leaves) — Lauraceae — HD — β-phellandrene (15.10%), linalool (14.10%) and γ-eudesmol (12.90%). — [18]<br>*A. rosaeodora* (Aerial parts) — Lauraceae — HD — linalool (88.60%) — [17]<br>*A. rosaeodora* (Aerial parts) — Lauraceae — HD — linalool (93.60%) — [19]<br>*Annona exsucca* (Dry leaves) — Annonaceae — HD — (*E*)-caryophyllene (31.26%), linalool (10.80%), β-elemene (10.30%), germacrene D (10.28%), bicyclogermacrene (9.84%) — [20]<br>*Bauhinia ungulata* (leaves) — Fabaceae — HD — (*E*)-caryophyllene (15.9%), caryophyllene oxide (9.2%) α-humulene (8.1%) and *epi*-γ-eudesmol (7.5%) — [21]<br>*Bocageopsis pleiosperma* (Barks) — Annonaceae — HD — β-bisabolene (38.53%), δ-cadinene (7.55%), β-selinene (6.46%) and α-selinene (5.18%) — [22]<br>*B. pleiosperma* (leaves) — Annonaceae — HD — β-bisabolene (55.77%), (*E*)-α-bergamotene (6.94%) and β-farnesene (*E*) (6.05%) — [22]<br>*B. pleiosperma* (twigs) — Annonaceae — HD — β-bisabolene (34.37%), cryptomerione (9,60%) and (2*Z*, 6*Z*)-farnesol (7,20%), — [22]<br>*B. multiflora* (Leaves) — Annonaceae — HD — spathulenol (20.30%) and β-bisabolene (11.90%) — [23]<br>*B. multiflora* (Aerial parts) — Annonaceae — HD — *cis*-linalool oxide (33.10%) and 1-epi-cubenol (16.60%). — [24]<br>*B. multiflora* (fresh leaves) — Annonaceae — HD — spathulenol (13.00–16.20%), β-bisabolene (13.20–13.80%) and caryophyllene oxide — [25]<br>Open in a separate window<br>HD: Hydrodistillation; SD: steam distillation; MAE: microwave-assisted extraction. | **Table 1**<br>Major chemical constituents (≥3.00%) found in the essential oils of the Amazon. | HD: Hydrodistillation; SD: steam distillation; MAE: microwave-assisted extraction. |
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T1/ | **TABLE 1**<br>Major chemical constituents of clove plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Amelia et al. (2017) — From clove bud — Clove from Java, Indonesia: eugenol 55.60%, eugenyl acetate 20.54%, caryophyllene 14.84%, *α*-humelene 2.75%, *β*-elemene 0.04%, *α*-cadinene 0.05%, ledol 0.06%; from Manodo, Indonesia: eugenol 74.64%, caryophylene 12.79%, eugenyl acetate 8.70%, *α*-humelene 1.53%, *β*-gurjunene 0.04%, *γ*-cadinene 0.03%, humelene oxide 0.05%; comparison with other papers, eugenol, 47.60–89.20%, caryophyllene, trace level to 35.40%, eugenyl acetate, 1.20–20.54%, *α*-humulene, trace level to 2.75%<br>Chaieb et al. (2007) — From clove flower bud essential oil — Eugenol 88.59%, eugenyl acetate 5.62%, *β*-caryophyllene 1.39%, 2-heptanone 0.93%, ethyl hexanoate 0.66%, *α*-humulene 0.20%, calacorene 0.11%, calamenene 0.11%<br>Jirowetz et al. (2006) — From clove leaf essential oil — Eugenol 76.8%, *β*-caryophyllene 17.4%, *α*-humulene 2.1%, eugenyl acetate 1.2%, caryophyllene oxide 0.4%, methylchavicol 0.2%<br>Uddin et al. (2017) — From clove buds oil in Bangladesh — m-eugenol 69.44%, eugenyl acetate 10.79%, caryophyllene 6.8%, tyranton 7.78%, trace amounts of other constituents <1% | **TABLE 1**<br>Major chemical constituents of clove plants and essential oils. | |
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T2/ | **TABLE 2**<br>Major chemical constituents of eucalyptus plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Barbosa et al. (2016) — Review paper — Major chemical constituents in the extracts from leaves are terpenes and terpenoids at various concentrations. The concentrations vary among the species of *Eucalyptus* and the location they were harvested. 1,8-Cineole most often is the most major chemical constituent in *E. camaldulensis*, *E. cinerea*, and *E. globulus* (percentages vary from as low as almost 10% to as high as over 90.0%), whereas citronellal is the major chemical constituent in *E. citriodora* (percentages varied from as low as almost 20% to as high as over 90%). *α*-Pinene is the major chemical constituent in *E. saligna* (from 24.4 to 45.1%), depending on the location the plants are harvested<br>Dogan et al. (2017) — Leaf Fruit — 1,8-cineole 14.1%, p-cymene 42.1%, *α*-pinene 12.7%, *α*-terpineol 10.7%, limonene 5.5%, borneol L 5.5%, spathulenol 3.2% 1,8-cineole 34.5%, p-cymene 30.0%, *α*-terpineol 15.1%, *α*-pinene 9.0%, borneol L 5.3%, *γ*-terpinene 5.1%, spathulenol none, limonene none<br>Salehi et al. (2019b) — Review paper — Percentages are not shown<br>Sebei et al. (2015) — Review paper — Comparing the percentages among species, 1,8-cineole (49.07–83.50%) and *α*-pinene (1.27–26.35%) were the two major chemical constituents | **TABLE 2**<br>Major chemical constituents of eucalyptus plants and essential oils. | |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T3/ | **TABLE 3**<br>Major chemical constituents of lemon plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Klimek-Szczykutowicz et al. (2020) — Essential oil of pericarp: Limonene 69.9%, p-menta-3,8-diene 18.0%, β-pinene 11.2%, γ-terpinene 8.21%, myrcene 4.4%, sabinene 3.9%, geranial 2.9%, neral 1.5%, linalool 1.41%, α-pinene 1.1%, α-thujene 1.1%, β-bisabolene 0.5%, (E)-β-ocimene 0.4%<br>Essential oil of leaf: geraniol 0.2%, β-caryophyllene 0.2% limonene 31.5%, sabinene 15.9%, citronellal 11.6%, linalool 4.6%, neral 4.5%, geranial 4.5%, (E)-β-ocimene 3.9%, myrcene 2.9%, citronellol 2.3%, β-caryophyllene 1.7%, terpinen-4-ol 1.4%<br>Luciardi et al. (2021) — Essential oils: Limonene 59.14%, γ-terpinene 10.48%, β-pinene 15.41%, sabinene 1.76%, β-myrcene 1.65%, α-pinene 1.64%<br>Pucci et al. (2020) — The whole fractions: Limonene 67.1%, β-pinene 10.88%, γ-terpinene 9.32%, α-pinene 1.81%, geranial 1.72%, sabinene 1.83%, myrcene 1.57% | **TABLE 3**<br>Major chemical constituents of lemon plants and essential oils. | |
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T4/ | **TABLE 4**<br>Major chemical constituents of rose plants and essential oils.<br>Data source / Source of samples / Major chemical constituents<br>Akram et al. (2019) — Oil Absolute rose Hydrosol: Geraniols (5.5–18%), β-citronellol (14.5–47.5%), nonadecane (10.5–40.5%) heneicosane, ethanol (0–13.43%), geraniol (3.71%), citronellol (9.91%), nonadecane (4.35%), phenylethylalcohol (78.38%) nerol (16.12%), phenylethyl alcohol (23.74%), citronellol (29.44%), geraniol (30.74%)<br>Mileva et al. (2021) — Oils: There are large differences among geological differences and species differences; geraniol (17.60–30.98%), nerol (4.36–10.10%), citronellol (9.22–28.72%), n-nonadecane (8.10–22.67%), n-heneicosane (5.00–10.21%)<br>Ryu et al. (2020) — Oils: There are large differences among breeds (mutants) of roses<br>Verma et al. (2010) — From bud, half bloom, full bloomEssential oils: Phenyl ethyl alcohol (PEA) 66.2–80.7%, other major chemical constituent, although much less in the percentages, were citronellol (1.8–5.5%) and geraniol (4.4–7.9%). Major chemical constituents were citronellol (15.9–35.3%), geraniol (8.3–30.2%), nerol (4.0–9.6%), nonadecane (4.5–16.0%), and heneicosane (2.6–7.9%), and not PEA (0.6–2.9%)<br>Open in a separate window | **TABLE 4**<br>Major chemical constituents of rose plants and essential oils. | |
| https://europepmc.org/article/MED/35721200 | Chemical Constituents of Essential Oils Used in Olfactory Training: Focus on COVID-19 Induced Olfactory Dysfunction. | https://europepmc.org/articles/PMC9201274/table/T5/ | **TABLE 5**<br>Chemicals compounds in the four essential oils with anti-inflammatory effects and binding affinity with SARS-CoV-2.<br>Chemical constituents with anti-inflammatory effect / Effects other than anti-inflammation / References<br>Borneol — Anti-viral (HSV-1 virus) — Armaka et al. (1999)<br>β-caryophyllene — Facilitate regeneration, enhance cell proliferation/migration; analgesic — Koyama et al. (2019)<br>1,8-cineole (eucalyptol) — — —<br>Citronellol — Activate peroxisome proliferator-activated receptor (PPAR) alpha and gamma — Katsukawa et al. (2011)<br>p-cymene — — —<br>Eugenol — — —<br>Eugenyl acetate — — —<br>Geraniol — Anti-angiogenic, anti-cell proliferative, apoptosis-inducing effects Vinothkumar et al., (2012); anti-ulcerogenic effects de Carvalho et al. (2014); Binding affinity to SARS-CoV-2 RBD of S-glycoprotein Kulkarni et al. (2021); activate peroxisome proliferator-activated receptor (PPAR) alpha and gamma Katsukawa et al. (2011) — Vinothkumar et al. (2012), de Carvalho et al. (2014), Kulkarni et al. (2021), Katsukawa et al. (2011)<br>α-humulene — — —<br>Limonene — Anti-tumorigenesis — —<br>Linalool — Analgesic — —<br>Myrcene — Analgesic — —<br>(E)-β-ocimene — — —<br>α-pheliandrene — — —<br>α-pinene — — —<br>Sabinene — — —<br>Spathulenol — Dnti-nociceptive effects — Dos Santos et al. (2020)<br>γ-terpinene — — — | **TABLE 5**<br>Chemicals compounds in the four essential oils with anti-inflammatory effects and binding affinity with SARS-CoV-2. | |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35747375 | Determination of Volatile Compounds of Mentha piperita and Lavandula multifida and Investigation of Their Antibacterial, Antioxidant, and Antidiabetic Properties. | https://europepmc.org/articles/PMC9213120/table/tab1/ | **Table 1**<br>Chemical composition of MPEO and LMEO.<br>Number \| Compounds \| RT \| MPEO \| LMEO<br>1 \| α-Pinene \| 2.07 \| 0.41 \| 0.30<br>2 \| Camphene \| 2.2 \| 0.03 \| 0.74<br>3 \| β-Terpinene \| 2.72 \| 0.95 \| 2.98<br>4 \| 2,3-Dehydro-1,8-cineole \| 3.01 \| — \| 0.16<br>5 \| α-Phellandrene \| 3.12 \| 0.06 \| 0.34<br>6 \| 3-Carene \| 3.4 \| — \| 1.86<br>7 \| Eucalyptol \| 3.918 \| 6.75 \| 28.11<br>8 \| Trans-β-ocimene \| 4.31 \| — \| 4.93<br>9 \| Terpinen-4-ol \| 4.69 \| 0.27 \| 3.0<br>10 \| Linalool \| 6.08 \| 0.23 \| 2.89<br>11 \| 2-Bornanone \| 6.71 \| — \| 11.57<br>12 \| Menthone \| 7.15 \| 29.24 \| —<br>13 \| Endo-borneol \| 7.2 \| — \| 7.82<br>14 \| Menthol \| 7.35 \| 2.71 \| 1.90<br>15 \| Terpinen-4-ol \| 7.46 \| — \| 7.65<br>16 \| Levomenthol \| 8.22 \| 38.73 \| 3.26<br>17 \| Caren-2-ol \| 8.89 \| 0.15 \| 3.24<br>18 \| Pulegone \| 9.09 \| 1.07 \| —<br>119 \| D-Carvone \| 9.21 \| 0.11 \| 0.15<br>20 \| Linalyl acetate \| 9.77 \| — \| 5.22<br>21 \| Bornyl acetate \| 10.15 \| — \| 0.07<br>22 \| Lavandulyl acetate \| 10.67 \| — \| 1.35<br>23 \| Elemene \| 11.52 \| 0.11 \| —<br>24 \| α-Copaene \| 12.07 \| 0.20 \| —<br>25 \| β-Bourbonene \| 12.89 \| 0.38 \| —<br>26 \| Geranyl acetate \| 13.50 \| — \| 0.35<br>27 \| Caryophyllene \| 13.71 \| — \| 1.45<br>28 \| β-Copaene \| 13.98 \| 0.20 \| —<br>29 \| Germacrene D \| 14.28 \| 0.24 \| 0.15<br>30 \| α-Bergamotene \| 14.32 \| — \| 0.18<br>31 \| β-Famesene \| 14.91 \| 0.16 \| 4.91<br>32 \| γ-Muurolene \| 14.98 \| 0.39 \|<br>33 \| Lavandulyl butyrate \| 15.3 \| — \| 1.02<br>34 \| β-Bisabolene \| 15.6 \| — \| 0.14<br>35 \| α-Curcumene \| 15.99 \| 0.86 \|<br>36 \| Caryophyllene oxide \| 16.06 \| 3.42 \| 0.22<br>37 \| (+)-β-Himachalene oxide \| 16.9 \| 0.06 \| —<br>38 \| Caryophylla-4(12),8(13)-dien- \| 17.20 \| 0.13 \| —<br>39 \| Cadinol \| 17.31 \| — \| 0.21<br>40 \| Alloaromadendrene \| 17.35 \| 0.17 \| —<br>41 \| Aromadendrene oxide \| 17.44 \| 0.14 \| —<br>42 \| α-Bisabolol \| 17.91 \| — \| 0.53<br>43 \| p-Menth-8-en-3-ol, acetate \| 18.03 \| 0.07 \| —<br>44 \| Culmorin \| 18.69 \| 0. 41 \| —<br>Total \| \| 96.04 \| 97.7<br>Others \| \| 3.96 \| 2.3 | **Table 1**<br>Chemical composition of MPEO and LMEO. | |
| https://europepmc.org/article/MED/35630653 | Investigation and Biological Assessment of Rumex vesicarius L. Extract: Characterization of the Chemical Components and Antioxidant, Antimicrobial, Cytotoxic, and Anti-Dengue Vector Activity | https://europepmc.org/articles/PMC9147116/table/molecules-27-03177-t001/ | **Table 1**<br>The characterized chemical components isolated from the extracted shoots of *Rumex vesicarius*.<br>Entry \| Chemical Name \| Classification \| Rt \| MF \| Composition %<br>Oxygenated Hydrocarbon<br>1 \| (*Z*)-2-Ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine \| Aryl substituted cyclic amine \| 4.21 \| $C_{20}H_{23}N$ \| 0.92<br>2 \| 3-(2-Oxocyclohexyl)propanenitrile \| Oxygenated hydrocarbon \| 5.54 \| $C_9H_{13}NO$ \| 0.41<br>3 \| Ethyl 2-hydroxycyclohexane-1-carboxylate \| Oxygenated hydrocarbon \| 9.22 \| $C_9H_{16}O_3$ \| 18.96<br>4 \| 2-Propyltetrahydro-2*H*-pyran-3-ol \| Oxygenated hydrocarbon \| 9.38 \| $C_8H_{16}O_2$ \| 11.18<br>5 \| Ascaridole epoxide \| Oxygenated hydrocarbon \| 12.99 \| $C_{10}H_{16}O_3$ \| 0.67<br>6 \| 3,5-Heptadienal, 2-ethylidene-6-methyl-"(2*Z*,3*E*)-2-ethylidene-6-methylhepta-3,5-dienal" \| Oxygenated hydrocarbon \| 13.07 \| $C_{10}H_{14}O$ \| 0.28<br>7 \| 1,25-Dihydroxyvitamin D3, TMS derivative \| Oxygenated hydrocarbon \| 34.16 \| $C_{30}H_{52}O_3Si$ \| 0.20<br>8 \| Deoxyspergualin \| Polyamine spermidine \| 8.74 \| $C_{17}H_{37}N_7O_3$ \| 0.36<br>9 \| Methyl 2,2,3,3,4,4,4-heptafluorobutanoate \| Ester \| 12.87 \| $C_5H_3F_7O_2$ \| 0.20<br>10 \| Tetraacetyl-D-xylonic nitrile \| Polyester \| 14.63 \| $C_{14}H_{17}NO_9$ \| 0.41<br>Fatty Acids and Lipids<br>11 \| High oleic safflower oil \| Vegetable oil "80% oleic acid" \| 4.45 \| $C_{21}H_{22}O_{11}$ \| 2.93<br>12 \| Methyl octadeca-8,11-diynoate \| Fatty acid \| 6.13 \| $C_{19}H_{30}O_2$ \| 3.82<br>13 \| (2-Phenyl-1,3-dioxolan-4-yl)methyl oleate \| Fatty-acid derivative \| 7.40 \| $C_{28}H_{44}O_4$ \| 5.75<br>14 \| bis(2-Ethylhexyl) adipate \| Ester of fatty acid \| 7.94 \| $C_{22}H_{42}O_4$ \| 0.80<br>15 \| D-Lyxo-D-manno-nononic-1,4-lactone \| Lactone of tetronic acid \| 11.31 \| $C_9H_{16}O_9$ \| 0.56<br>16 \| *tert*-Butyl palmitate \| Ester of fatty acid \| 14.52 \| $C_{20}H_{40}O_2$ \| 0.26<br>17 \| Oleic acid \| Fatty acid \| 14.67 \| $C_{18}H_{34}O_2$ \| 0.57<br>18 \| 9-Hexadecenoic acid \| Fatty acid \| 15.20 \| $C_{16}H_{30}O_2$ \| 0.17<br>19 \| *trans*-2-Dodecenoic acid \| Fatty acid \| 15.52 \| $C_{12}H_{22}O_2$ \| 0.24<br>20 \| 2-Hydroxypropane-1,3-diyl (9*E*,9'*E*)-bis(octadec-9-enoate) \| Diester derivative of fatty acid \| 16.64 \| $C_{39}H_{72}O_5$ \| 0.94<br>21 \| (*E*)-Octadec-13-enoic acid \| Fatty acid \| 17.13 \| $C_{18}H_{34}O_2$ \| 2.74<br>22 \| 2-Bromotetradecanoic acid \| Fatty-acid derivative \| 19.89 \| $C_{14}H_{27}BrO_2$ \| 1.45<br>23 \| [1,1'-Bicyclopropyl]-2-octanoic acid, 2'-hexyl-, methyl ester \| Fatty-acid derivative \| 19.96 \| $C_{21}H_{38}O_2$ \| 0.90<br>24 \| 8-[(2*R*,3*S*)-3-Octyloxiran-2-yl]octanoic acid \| Fatty acid \| 21.41 \| $C_{18}H_{34}O_3$ \| 1.09<br>25 \| 2,3-Dihydroxypropyl stearate \| Fatty acid \| 21.69 \| $C_{21}H_{42}O_4$ \| 1.23<br>26 \| Methyl 5-[(1*R*,2*R*)-2-undecylcyclopropyl)-pentanoate \| Fatty-acid derivative \| 25.88 \| $C_{20}H_{38}O_2$ \| 4.50<br>27 \| 2-Hydroxypropane-1,3-diyl dipalmitate \| Ester of fatty acid \| 28.31 \| $C_{35}H_{68}O_5$ \| 2.87<br>28 \| 1,3-Dihydroxypropan-2-yl oleate \| Ester of fatty acid \| 29.18 \| $C_{21}H_{40}O_4$ \| 17.56<br>29 \| Methyl 11-[(2*R*,3*R*)-3-pentyloxiran-2-yl]undecanoate \| Ester of fatty acid \| 29.70 \| $C_{19}H_{36}O_3$ \| 1.75<br>30 \| 9-Octadecenoic acid,1,2,3-propanetriyl ester, (*E*,*E*,*E*)- \| Ester of fatty acid \| 31.35 \| $C_{57}H_{104}O_6$ \| 1.23<br>Carbohydrates<br>31 \| D-Gala-L-ido-octonic amide "2,3,4,5,6,7,8-heptahydroxyoctanamide" \| Carbohydrate amide \| 6.50 \| $C_8H_{17}NO_8$ \| 0.26<br>32 \| Desulfosinigrin "1-*S*-[(1*E*)-*N*-hydroxy-3-butenimidoyl]-1-thiohexopyranose" \| Glycoside \| 6.64 \| $C_{10}H_{17}NO_6S$ \| 0.27<br>33 \| L-Gala-L-ido-octose \| Carbohydrate \| 6.85 \| $C_8H_{16}O_8$ \| 1.09<br>34 \| Melezitose \| Trisaccharide sugar \| 8.79 \| $C_{18}H_{32}O_{16}$ \| 0.34<br>35 \| D-Ribo-hexos-3-ulose "(2*S*,4*R*,5*R*)-2,4,5,6-tetrahydroxy-3-oxohexanal" \| Dicarbonyl sugar \| 9.84 \| $C_6H_{10}O_6$ \| 0.30<br>36 \| 2,3-Dihydroxypropyl palmitate \| 1-Monoacylglycerols \| 4.89 \| $C_{19}H_{38}O_4$ \| 3.80<br>Amines<br>37 \| *N*2,*N*4-Diisopropyl-6-(methylsulfonyl)-1,3,5-triazine-2,4-diamine \| Hetryl amine \| 4.14 \| $C_{10}H_{19}N_5O_2S$ \| 2.44<br>38 \| (*E*)-[2-(Chloroimino)-3-methylbutanoyl]-L-valine \| Amino acid \| 4.94 \| $C_{10}H_{17}ClN_2O_3$ \| 3.84<br>39 \| *S*-(2-Aminoethyl) *O*-hydrogen sulfurothioate \| Amino-thioester \| 11.39 \| $C_2H_7NO_3S_2$ \| 0.59<br>40 \| Glutamic acid \| Amino acid \| 11.46 \| $C_5H_9NO_4$ \| 0.34<br>41 \| Methyl *N*-acetyl-D-glucosamide \| *N*-Acetyl-D-glucosamine \| 10.35 \| $C_9H_{17}NO_6$ \| 0.14<br>Steroids<br>42 \| Estra-1,3,5(10)-trien-17β-ol \| Steroid \| 20.54 \| $C_{18}H_{24}O$ \| 0.34<br>43 \| Ethyl iso-allocholate \| Steroidal ester \| 34.94 \| $C_{26}H_{44}O_5$ \| 0.87<br>RT: retention time, MF: molecular formula. | **Table 1**<br>The characterized chemical components isolated from the extracted shoots of *Rumex vesicarius*. | RT: retention time, MF: molecular formula. |
| https://europepmc.org/article/MED/34073176 | Impact of Altitudinal Variation on the Phytochemical Profile, Anthelmintic and Antimicrobial Activity of Two Pinus Species. | https://europepmc.org/articles/PMC8198815/table/molecules-26-03170-t001/ | **Table 1**<br>Chemical composition of the essential oils of *P. halepensis* (Ph-1 and Ph-2) and *P. pinea* (Pp-1 and Pp-2).<br>\# \| \| AI \| RI \| Ph-1 \| Ph-2 \| Pp-1 \| Pp-2<br>***Monoterpene***<br>1. \| α-Pinene \| 932 \| 939 \| **8.54** ± 0.50 a \| **13.33** ± 0.21 b \| **13.82** ± 0.03 b \| **20.97** ± 0.16 c<br>2. \| Sabinene \| 970 \| 976 \| 3.15 ± 0.20 \| – \| – \| –<br>3. \| β-pinene \| 974 \| 980 \| 4.08 ± 0.31 \| – \| – \| –<br>4. \| β-Myrcene \| 990 \| 990 \| **5.80** ± 0.05 a \| – \| – \| 1.37 ± 0.06 b<br>5. \| δ-3-carene \| 1008 \| 1011 \| 4.07 ± 0.19 \| – \| – \| –<br>6. \| p-Cymene \| 1022 \| 1022 \| 1.39 ± 1.10 a \| - \| 0.82 ± 0.21 a \| 0.69 ± 0.11 a<br>7. \| Limonene \| 1024 \| 1029 \| 1.49 ± 0.02 a \| **6.77** ± 0.02 b \| 1.13 ± 0.14 a \| **8.49** ± 0.01 b<br>8. \| γ-terpinene \| 1054 \| 1055 \| 3.38 ± 0.19 a \| – \| 3.18 ± 0.09 a \| –<br>9. \| α-Terpinolene \| 1018 \| 1088 \| 1.08 ± 0.05 a \| – \| **5.67** ± 0.12 b \| 4.75 ± 0.21 b<br>27.18 \| 20.1 \| 24.62 \| 36.27<br>***Oxygenated monoterpenes***<br>10. \| α-Campholenal \| 1130 \| 1128 \| – \| – \| 1.85 ± 0.02 a \| 0.63 ± 0.02 a<br>11. \| *trans*-pinocarveol \| 1135 \| 1136 \| – \| – \| 4.89 ± 0.25 a \| 1.08 ± 0.12 b<br>12. \| Pinocarvone \| 1160 \| 1164 \| – \| – \| 1.77 ± 0.31 a \| 1.35 ± 0.05 a<br>13. \| Borneol \| 1165 \| 1168 \| – \| – \| 2.47 ± 0.42 a \| 0.91 ± 0.14 a<br>14. \| Myrtanol \| 1194 \| 1195 \| – \| – \| 1.20 ± 0.29 \| -<br>15. \| Verbenone \| 1204 \| 1206 \| – \| – \| 0.56 ± 0.00 \| -<br>16. \| *trans*-(+)-carveol \| 1215 \| 1219 \| – \| – \| 1.04 ± 0.17 \| -<br>17. \| Carveol methyl ether \| 1229 \| 1229 \| – \| – \| 1.09 ± 0.07 \| -<br>18. \| Carvone \| 1239 \| 1236 \| – \| – \| 0.57 ± 0.12 \| -<br>19. \| Bornyl acetate \| 1287 \| 1285 \| – \| – \| 1.49 ± 0.06 a \| 1.29 ± 0.07 a<br>20. \| Thymol \| 1232 \| 1232 \| – \| – \| **5.53** ± 0.05 a \| 0.65 ± 0.01 b<br>21. \| Carvacrol \| 1298 \| 1298 \| – \| – \| 0.90 ± 0.23 a \| 2.08 ± 0.00 a<br>Zero \| Zero \| 23.36 \| 7.99<br>***Sesquiterpene hydrocarbons***<br>22. \| α-Cubebene \| 1345 \| 1348 \| – \| – \| 1.42 ± 0.09 a \| 2.64 ± 0.02 a<br>23. \| α-longipinene \| 1350 \| 1350 \| 3.47 ± 0.19 a \| – \| 2.80 ± 0.01 a \| 2.64 ± 0.19 a<br>24. \| β-Elemene \| 1389 \| 1390 \| – \| 2.06 ± 0.07 a \| 1.26 ± 0.05 a \| –<br>25. \| β-Caryophyllene \| 1417 \| 1417 \| 5.29 ± 0.00 a \| 6.20 ± 0.19 a \| 1.65 ± 0.12 b \| **5.39** ± 0.01 a<br>26. \| Aromadendrene \| 1439 \| 1430 \| 1.61 ± 0.12 a \| 1.38 ± 0.12 a \| 1.01 ± 0.42 a \| –<br>27. \| α-Humulene \| 1452 \| 1454 \| 3.12 ± 0.05 a \| 2.0 ± 0.29 a \| 1.19 ± 0.41 b \| 3.39 ± 0.20 a<br>28. \| Germacrene D \| 1484 \| 1485 \| – \| 0.68 ± 0.02 a \| – \| **7.81** ± 0.03 b<br>29. \| δ-amorphene \| 1511 \| 1507 \| – \| – \| 1.82 ± 0.57 a \| 4.54 ± 0.11 b<br>30. \| Cis-α-bisabolene \| 1508 \| 1508 \| – \| – \| 1.38 ± 0.67 a \| 0.65 ± 0.09 a<br>31. \| γ-cadinene \| 1514 \| 1514 \| – \| – \| 0.21 ± 0.03 \| –<br>32. \| δ-cadinene \| 1523 \| 1522 \| 2.63 ± 0.01 a \| 2.03 ± 0.09 a \| 4.54 ± 0.07 b \| 2.62 ± 0.02 a<br>16.12 \| 14.35 \| 19.95 \| 33.16<br>***Oxygenated sesquiterpenes***<br>33. \| Caryophyllene oxide \| 1582 \| 1581 \| 1.39 ± 0.03 a \| 4.45 ± 0.03 b \| 1.70 ± 0.54 a \| 3.55 ± 0.07 b<br>34. \| Aromadendrene oxide \| 1595 \| 1595 \| 2.37 ± 0.10 a \| **12.13** ± 0.36 b \| 3.63 ± 0.21 a \| 1.57 ± 0.08 a<br>35. \| Cubenol \| 1618 \| 1616 \| – \| – \| 0.89 ± 0.15 a \| 0.92 ± 0.15 a<br>36. \| Muurolol \| 1640 \| 1642 \| – \| – \| 2.59 ± 0.00 a \| 2.95 ± 0.09 a<br>37. \| Vulgarol B \| 1688 \| 1688 \| – \| – \| 2.55 ± 0.02 a \| 1.74 ± 0.01 a<br>3.76 \| 16.58 \| 11.36 \| 10.73<br>***Diterpene hydrocarcons***<br>38. \| Cembrene \| 1937 \| 1932 \| 2.54 ± 0.05 a \| **8.23** ± 0.07 b \| 2.10 ± 0.02 a \| –<br>2.54 \| 8.23 \| 2.1 \| –<br>***Oxygenated diterpenes***<br>39. \| Manool oxide \| 1987 \| 1994 \| 3.94 ± 0.02 a \| 2.55 ± 0.25 b \| 5.06 ± 0.00 a \| 2.25 ± 0.15 b<br>40. \| Thunbergol \| 2047 \| 2047 \| **5.76** ± 0.58 a \| **18.85** ± 0.17 b \| **6.16** ± 0.02 a \| –<br>41. \| 1,4-menthano-azulene \| 2110 \| 2110 \| – \| – \| 2.67 ± 0.09 a \| 3.48 ± 0.07 a<br>42. \| Dehydro abietinal \| 2279 \| 2279 \| 4.14 ± 0.10 \| – \| – \| –<br>43. \| Kaur-16-en-19-ol \| 2346 \| 2346 \| 3.28 ± 0.21 \| – \| – \| –<br>44. \| Methyl dehydroabietate \| 2359 \| 2354 \| **7.78** ± 0.03 a \| – \| 0.71 ± 0.19 b \| –<br>45. \| Dehydro abietic acid \| 2380 \| 2380 \| 5.74 ± 0.31 a \| 1.38 ± 0.27 b \| 0.71 ± 0.30 b \| –<br>46. \| Abietic acid methyl ester \| 2377 \| 2380 \| 3.35 ± 0.06 a \| – \| 0.84 ± 0.09 b \| –<br>33.99 \| 22.78 \| 12.64 \| 2.25<br>***Miscellaneous compounds***<br>47. \| 2-pentadecanone \| 1694 \| 1698 \| – \| – \| 1.59 ± 0.01 a \| 0.70 ± 0.21 a<br>48. \| Pentadecanal \| 1716 \| 1716 \| – \| – \| 1.91 ± 0.09 \| –<br>– \| – \| 4.34 \| 0.7<br>Total % \| \| \| \| 89.39 ± 4.10 \| 82.04 ± 2.16 \| 94.03 ± 6.75 \| 91.1 ± 2.46<br>Yield% (v/w) \| \| \| \| 0.22 \| 0.59 \| 0.12 \| 0.22<br>(Ph-1): *P. halepensis* at altitude 830 m; (Ph-2): *P. halepensis* at altitude 75 m; (Pp-1): *P. pinea* at altitude 625 m; (Pp-2): *P. pinea* at alt<br>*n*-alkanes. RI, Published Kovats retention indices. The main compounds are in bold. The values shown in this table were the aver<br>range test were used. Values with different superscripts (a–c) were significantly different at $p < 0.05$. Values followed by a commi | **Table 1**<br>Chemical composition of the essential oils of *P. halepensis* (Ph-1 and Ph-2) and *P. pinea* (Pp-1 and Pp-2). | (Ph-1): *P. halepensis* at altitude 830 m; (Ph-2): *P. halepensis* at altitude 75 m; (Pp-1): *P. pinea* at altitude 625 m; (Pp-2): *P. pinea* at altitude 408 m. AI, Kovats index determined experimentally on HP-5MS column relative to C8–C28 *n*-alkanes. RI, Published Kovats retention indices. The main compounds are in bold. The values shown in this table were the average of three replicates and given as mean ± SD ($n = 3$). One-way ANOVA followed by Duncan's multiple range test were used. Values with different superscripts (a–c) were significantly different at $p < 0.05$. Values followed by a common letter in columns were not significant ($p > 0.05$). |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35745024 | Phytochemical Composition of Commiphora Oleogum Resins and Their Cytotoxicity against Skin Cancer Cells. | https://europepmc.org/articles/PMC9229828/table/molecules-27-03903-t004/ | **Table 4**<br>Chemical composition of *Commiphora* oleogum resin essential oils. Relative quantification by gas chromatography and flame ionization detection (GC-FID) using internal normalization, areas in uncorrected %. List of main components (area > 1%). Hydrodistillation of *Commiphora* oleogum resin from Tarraxo (Somalia) yielded no essential oil and consequently, no GC analysis is shown. Traces tr < 0.05%. Complete data are shown in Supplementary Materials, Table S3.<br>Compound \| *C. myrrha* (Somalia) \| *C. erythraea* (Somalia) \| *C. mukul* (Nepal) \| *C. kataf* (Kenya) \| *C. holtziana* (Kenya) \| *C. confuse* (Kenya) \| *C. kua* (Socotra, Yemen) \| *C.* from Ogaden (Ethiopia)<br>α-Thujene - tr 0.2 0.2 0.1 9.0 - tr<br>α-Pinene - 0.5 8.5 5.8 0.8 39.5 tr 0.5<br>3,7,7-Trimethylcyclohepta-1,3,5-triene - - 2.2 - - tr - -<br>β-Pinene - 0.1 2.0 3.5 0.1 8.9 - 0.1<br>Sabinene - tr 0.2 0.3 tr 1.1 - tr<br>$\Delta^3$-Carene - tr 29.8 tr - - - tr<br>*para*-Cymene - 0.1 2.1 0.2 0.1 10.0 - 0.1<br>Limonene 0.1 0.1 1.2 0.2 0.1 1.1 0.1 0.1<br>β-Thujone - - - - tr 1.0 - -<br>*trans*-Pinocarveol - - 0.3 0.3 0.1 2.0 tr tr<br>α-Phellandren-8-ol - tr 0.3 0.2 - 1.6 - tr<br>Terpinen-4-ol - tr 1.0 0.1 tr 4.5 tr tr<br>*meta*-Cymen-8-ol - - 1.2 - - - - -<br>α-Terpineol - tr 1.3 0.1 - 1.0 - tr<br>Myrtenol - - 0.2 0.2 - 1.0 tr -<br>Verbenone - tr 1.3 0.1 tr 1.6 tr tr<br>δ-Elemene 2.0 1.1 - 25.1 1.8 - 0.9 1.8<br>α-Terpinyl acetate - - 1.1 - - 0.1 - -<br>α-Copaene 0.4 1.3 0.1 1.4 2.2 tr 1.5 0.6<br>β-Bourbonene 1.2 5.2 - 3.7 1.3 - 0.9 2.1<br>β-Elemene 9.1 11.1 0.1 4.7 9.6 tr 0.9 12.8<br>Longifolene - - 24.3 - - - - -<br>*cis*-α-Bergamotene 0.2 - - 1.2 0.5 - - 0.1<br>β-Caryophyllene 0.7 1.1 0.7 1.2 0.9 - 7.7 1.0<br>β-Copaene 0.4 1.1 - 1.0 0.4 - 0.3 0.7<br>γ-Elemene 2.7 1.1 - 0.5 1.8 - 0.2 3.8<br>*trans*-α-Bergamotene 0.1 1.8 - - 0.3 - 0.4 0.9<br>α-Humulene 0.5 0.7 0.1 1.0 0.5 - 3.9 0.7<br>Allo-Aromadendrene 0.1 0.3 - 0.3 0.2 - 3.8 0.2<br>*cis*-Cadina-1(6),4-diene - 0.1 - 0.1 - - 1.7 0.2<br>γ-Muurolene 0.2 1.7 - 1.6 0.6 - 2.0 1.0<br>Germacrene D 3.3 1.2 - 13.7 1.8 - 1.5 8.3<br>β-Selinene 1.1 1.9 - 2.1 1.8 - 1.2 1.1<br>α-Selinene 1.2 2.2 - 0.9 1.8 - 1.8 1.3<br>Curzerene 29.7 37.8 - - 18.7 - tr 32.6<br>(*Z*)-α-Bisabolene - 4.8 - - - - - 0.2<br>β-Bisabolene - 1.2 0.1 - - - - 0.1<br>γ-Cadinene 0.1 0.9 - 1.2 0.4 - 24.0 2.0<br>δ-Cadinene 0.4 2.3 tr 1.7 0.9 - 8.9 1.4<br>α-Cadinene - 0.2 - 0.2 - - 1.9 0.1<br>α-Elemol 0.2 1.1 - 0.5 0.4 - 0.1 1.1<br>Germacrene B 4.6 1.3 - 0.7 2.6 - 0.3 4.9<br>Curzerenone 0.6 0.3 - 2.6 2.2 - 0.1 2.7<br>10-*epi*-Cubenol - 0.1 - 0.3 - - 3.3 0.2<br>Furanoeudesma-1,3-diene 17.4 - - - 23.9 - tr -<br>Lindestrene 8.7 - - - 6.5 - tr -<br>τ-Cadinol 0.2 0.2 - 0.6 0.2 - 16.9 0.6<br>Furanodiene 0.4 1.1 - - 0.4 - - -<br>Germacrone 0.4 0.3 - 0.2 0.3 - - 1.8<br>2-Methoxyfuranodiene 1.1 - - - 0.4 - - - | **Table 4**<br>Chemical composition of *Commiphora* oleogum resin essential oils. Relative quantification by gas chromatography and flame ionization detection (GC-FID) using internal normalization, areas in uncorrected %. List of main components (area > 1%). Hydrodistillation of *Commiphora* oleogum resin from Tarraxo (Somalia) yielded no essential oil and consequently, no GC analysis is shown. Traces tr < 0.05%. Complete data are shown in Supplementary Materials, Table S3. | |
| https://europepmc.org/article/MED/35351963 | Repellency and insecticidal activity of seven Mugwort (Artemisia argyi) essential oils against the malaria vector Anopheles sinensis. | https://europepmc.org/articles/PMC8964668/table/Tab5/ | **Table 5**<br>GC–MS analyses of compositional compounds of the essential oils extracted from seven regions of *Ar. argyi*.<br>Constituents \| CAS[a] \| RI–A[b] \| RI–B[c] \| Constituent percentage: CQ \| SC \| YN<br>Santolina triene 2153-66-4 908 907 – – 12.36<br>1R-α-Pinene 7785-70-8 929 932 2.58 1.38 2.49<br>α-Pinene 80-56-8 937 932 – – –<br>Camphene 79-92-5 952 947 – 1.24 –<br>Sabinen 3387-41-5 974 973 – 1.81 4.48<br>β-Pinene 127-91-3 979 977 1.62 2.02 –<br>( +)-2-Carene 149,946 991 1015 – – –<br>Yomogi alchool 26,127-98-0 1000 1000 – – –<br>α-Phellandrene 99-83-2 1005 1003 – – 1.91<br>o-Cymene 527-84-4 1022 1022 1.74 1.78 4.39<br>m-Cymene 535-77-3 1023 1022 – – –<br>p-Cymene 99-87-6 1025 1023 – – –<br>**Eucalyptol** 470-82-6 1032 1031 **11.44** **12.62** **5.49**<br>Artemisia ketone 546-49-6 1062 1061 – – –<br>cis-Sabinene hydroxide 17,699-16-0 1070 1068 – – –<br>Artemisia alcohol 27,644-04-8 1084 1084 2.26 – –<br>Thujone 546-80-5 1103 1107 – 16.66 –<br>Isothujone 471-15-8 1114 1117 – – –<br>Chrysanthone 1125-12-8 1119 1108 – – –<br>Isochrysanthenone 473-06-3 1123 1125 4.08 1.8 –<br>(-)-Camphor 464-48-2 1142 1143 – 1.49 2.15<br>( +)-2-Bornanone 471-16-9 1143 1143 – – –<br>p-Menth-8-en-1-ol, cis 7299-41-4 1144 1066 – 1.35 –<br>Open in a separate window<br>[a]CAS: Chemical Abstracts Service Registry Number. [b]Retention index from NIST references (qualitative index of gas chromatography). [c]Retention index | **Table 5**<br>GC–MS analyses of compositional compounds of the essential oils extracted from seven regions of *Ar. argyi*. | [a]CAS: Chemical Abstracts Service Registry Number. [b]Retention index from NIST references (qualitative index of gas chromatography). [c]Retention index calculated with N alkanes ($C_5$–$C_{25}$Hc) as standard. The chemical constituents with less than 1% composition in each essential oil are not listed in the table. CQ, SC, YN, HB, HN, SD and GS: essential oils from Chongqing, Sichuan, Yunnan, Hubei, Henan, Shandong and Gansu province/municipality, respectively. |
| https://europepmc.org/article/MED/35523830 | Application of direct peptide reactivity assay for assessing the skin sensitization potential of essential oils. | https://europepmc.org/articles/PMC9076902/table/Tab1/ | **Table 1**<br>Chemical composition of selected essential oils.<br>Compounds \| \| RI[a] \| *Thymi aetheroleum* \| *Rosmarini aetheroleum* \| *Salivae aetherolum* \| *Melissae aetheroleum* \| *Lavandulae aetheroleum*<br>Monoterpene hydrocarbons: α-pinene 939 1.54 11.18 7.59 0.18<br>α-terpinene 1020 1.89 0.45 0.24<br>α-thujene 930 1.03 0.12 0.33 0.11<br>β-pinene 981 0.72 1.97 1.23<br>Camphene 956 1.02 4.22 7.39<br>*cis*-β-ocimene 1050 3.30<br>γ-terpinene 1020 9.04 0.20 0.20 0.09<br>Limonene 1033 0.67 2.14 2.37 0.93<br>Myrcene 990 1.53 1.51 0.98 0.15 0.91<br>*p*-cymene 1029 29.93 3.58 1.57 0.53<br>Sabinene 974<br>Terpinolene 1008<br>*trans*-β-ocimene 1049 4.41<br>Other – 0.33 0.60 1.20 0.30 0.81<br>Monoterpene alcohols: α-terpinole 1196 0.37 5.55 0.34 2.80<br>Borneol 1176 1.73 4.99 3.11 2.13 1.95<br>Carvacrol 1298 3.60 0.08 0.07<br>Citronellol 1228 1.48<br>Geraniol 1252 6.56<br>Lavandulol 1166 3.78<br>Linalool 1100 2.38 1.39 0.46 37.87<br>Nerol 1226 1.50<br>Terpinene-4-ol 1183 0.87 1.18 4.92<br>Thymol 1292 36.69 0.09 0.90<br>Other – 1.40 0.40 0.17 2.70 0.70<br>Monoterpene oxides: 1,8-cineole 1037 1.39 41.94 13.41 0.45 0.35<br>α-thujone 1109 25.23 0.97<br>β-thujone 1121 7.42 0.27<br>Camphor 1152 0.65 16.31 19.27<br>*Cis*-linalool oxide 1038 1.56<br>Citronellal 1155 5.42<br>Geranial 1270 27.71<br>Neral 1241 19.54<br>[a]Retention index calculated relative to elution time of C9-C24 n-alkane from HP 5MS column. | **Table 1**<br>Chemical composition of selected essential oils. | [a]Retention index calculated relative to elution time of C9-C24 n-alkane from HP 5MS column. |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35336701 | Wild-Grown and Cultivated Glechoma hederacea L.: Chemical Composition and Potential for Cultivation in Organic Farming Conditions. | https://europepmc.org/articles/PMC8949430/table/plants-11-00819-t002/ | **Table 2**<br>Phytocomponents identified in the 70% ethanol extracts of *G. hederacea*.<br>Peak \| RT, min \| Compound \| Characteristic Ions [1], *m/z* [M + H]+ \| [M − H]− \| Calculated Elemental Composition [2] \| HRMS-MS/MS Fragments (ESI+), *m/z* \| Parent Scan of Aglycone Fragment [3] (ESI+), Ion, *m/z*<br>1 \| 6, 2 \| Chlorogenic acid isomer \| 355.103 \| 353.085 \| $C_{16}H_{18}O_9$ \| - \| -<br>2 \| 9, 0 \| **Chlorogenic acid** \| 355.103 \| 353.085 \| $C_{16}H_{18}O_9$ \| - \| -<br>3 \| 10, 0 \| Chlorogenic acid isomer \| 355.103 \| 353.085 \| $C_{16}H_{18}O_9$ \| - \| -<br>4 \| 11, 1 \| Quercetin 3-O-diglucoside \| 627.153 \| 625.138 \| $C_{27}H_{30}O_{17}$ \| 465.100; 303.049 \| 627, 465 (303)<br>5 \| 15, 1 \| Quercetin 3-O-galactosyl-rhamnoside \| 611.159 \| 609.142 \| $C_{27}H_{30}O_{16}$ \| 449.108; 303.050 \| 611, 449 (303)<br>6 \| 16, 5 \| Acylated quercetin diglycoside \| 755.202 \| 753.185 \| $C_{33}H_{38}O_{20}$ \| 449.106; 303.049 \| 755, 449, (303)<br>7 \| 16, 9 \| **Rutin** \| 611.157 \| 609.143 \| $C_{27}H_{30}O_{16}$ \| 465.102; 303.049 \| 611, 465, (303)<br>8 \| 17, 2 \| **Hyperoside** \| 465.102 \| 463.085 \| $C_{21}H_{20}O_{12}$ \| 303.049 \| 465 (303)<br>9 \| 17, 6 \| **Luteolin 7-O-glucoside** \| 449.107 \| 447.089 \| $C_{21}H_{20}O_{11}$ \| 287.054 \| 449 (287)<br>10 \| 18, 0 \| Acylated kaempferol diglycoside \| 739.207 \| 737.190 \| $C_{33}H_{38}O_{19}$ \| 433.113; 287.054 \| 739, 433, (287)<br>11 \| 18, 5 \| Acylated quercetin glycoside \| 609.145 \| 607.127 \| $C_{27}H_{28}O_{16}$ \| 303.049 \| 609 (303)<br>12 \| 18, 5 \| **Kaempferol 3-O-rutinoside** \| 595.163 \| 593.331 \| $C_{27}H_{30}O_{15}$ \| - \| 595 (287)<br>13 \| 19, 5 \| **Apigenin 7-O-glucoside** \| 433.113 \| 431.096 \| $C_{21}H_{20}O_{10}$ \| 271.059 \| 433 (271)<br>14 \| 20, 1 \| **Rosmarinic acid** \| - \| 359.075 \| $C_{18}H_{16}O_8$ \| - \| -<br>15 \| 20, 3 \| Acylated kaempferol glycoside \| 593.146 \| 591.133 \| $C_{27}H_{28}O_{15}$ \| 287.056 \| (287)<br>16 \| 22, 7 \| **Luteolin** \| 287.054 \| 285.040 \| $C_{15}H_{10}O_6$ \| - \| -<br>17 \| 25, 2 \| **Apigenin** \| 271.058 \| 269.045 \| $C_{15}H_{10}O_5$ \| - \| -<br>[1] HRMS data, [2] mass difference within ± 5 mDa, [3] *m/z* of aglycone fragment in brackets, **bold**—identified by comparison to reference compounds. | **Table 2**<br>Phytocomponents identified in the 70% ethanol extracts of *G. hederacea*. | [1] HRMS data, [2] mass difference within ± 5 mDa, [3] *m/z* of aglycone fragment in brackets, **bold**—identified by comparison to reference compounds. |
| https://europepmc.org/article/MED/35448753#free-full-text | Essential Oils and Extracts of Juniperus macrocarpa Sm. and Juniperus oxycedrus L.: Comparative Phytochemical Composition and Anti-Proliferative and Antioxidant Activities | https://europepmc.org/articles/PMC9031627/table/plants-11-01025-t001/ | **Table 1**<br>The main identified constituents of *J. macrocarpa* and *J. oxycedrus* essential oils<br>(−)-Carvone \| om \| 1242 \| 0.2 ± 0.03 \| *n.d.* \| 1,2 \| *<br>Piperitone \| om \| 1254 \| 0.2 ± 0.02 \| *n.d.* \| 1,2 \| *<br>Phellandral \| om \| 1281 \| 0.3 ± 0.02 \| 0.2 ± 0.01 \| 1,2 \| ns<br>(−)-Bornyl acetate \| om \| 1286 \| 0.2 ± 0.01 \| *n.d.* \| 1,2 \| *<br>α-Cubebene \| sh \| 1352 \| 0.2 ± 0.01 \| 0.5 ± 0.01 \| 1,2 \| ns<br>α-Copaene \| sh \| 1377 \| 0.2 ± 0.02 \| 0.3 ± 0.02 \| 1,2 \| ns<br>β-Bourbonene \| sh \| 1385 \| 0.3 ± 0.02 \| 0.9 ± 0.02 \| 1,2 \| **<br>β-Cubebene \| sh \| 1387 \| 1.8 ± 0.6 \| *n.d.* \| 1,2 \| **<br>α-Gurjunene \| sh \| 1407 \| *n.d.* \| 1.1 ± 0.04 \| 1,2 \| **<br>*trans*-Caryophyllene \| sh \| 1415 \| 0.2 ± 0.01 \| 2.0 ± 0.2 \| 1,2,3 \| **<br>*trans*-α-Bergamotene \| sh \| 1438 \| *n.d.* \| 0.4 ± 0.01 \| 1,2 \| **<br>α-Humulene \| sh \| 1455 \| 0.2 ± 0.01 \| 1.5 ± 0.08 \| 1,2 \| **<br>Germacrene D \| sh \| 1477 \| 4.5 ± 0.03 \| 2.0 ± 0.07 \| 1,2 \| **<br>γ-Cadinene \| sh \| 1515 \| 0.2 ± 0.01 \| 5.4 ± 0.6 \| 1,2 \| **<br>δ-Cadinene \| sh \| 1526 \| 0.4 ± 0.01 \| 2.7 ± 0.6 \| 1,2 \| **<br>(*E*)-β-Farnesene \| sh \| 1452 \| 0.4 ± 0.02 \| 0.7 ± 0.01 \| 1,2 \| **<br>α-Muurolene \| sh \| 1500 \| 0.3 ± 0.01 \| 0.4 ± 0.01 \| 1,2 \| ns<br>Caryophyllene oxide \| os \| 1580 \| 0.9 ± 0.04 \| 3.8 ± 0.5 \| 1,2 \| **<br>(*Z*,*E*)-Farnesol \| os \| 1722 \| 2.8 ± 0.2 \| 6.5 ± 0.8 \| 1,2,3 \| **<br>Manoyl oxide \| di \| 1989 \| 6.6 ± 0.6 \| 2.4 ± 0.4 \| 1,2 \| **<br>13-*epi*-Manoyl oxide \| di \| 1994 \| 0.6 ± 0.2 \| 0.6 ± 0.02 \| 1,2 \| ns<br>(*Z*)-Phytol \| di \| 1950 \| *n.d.* \| 0.5 ± 0.01 \| 1,2 \| **<br>Abietatriene \| di \| 2054 \| 2.9 ± 0.7 \| 2.6 ± 0.3 \| 1,2 \| *<br>Abietadiene \| di \| 2080 \| 1.8 ± 0.6 \| 1.7 ± 0.3 \| 1,2 \| ns<br>Heneicosane \| oc \| 2100 \| *n.d.* \| 0.2 ± 0.02 \| 1,2,3 \| *<br>Tricosane \| oc \| 2300 \| 0.3 ± 0.02 \| 0.2 ± 0.01 \| 1,2,3 \| ns<br>Pentacosane \| oc \| 2500 \| 0.2 ± 0.01 \| 0.5 ± 0.01 \| 1,2,3 \| *<br>Heptacosane \| oc \| 2700 \| 0.4 ± 0.01 \| 0.5 ± 0.02 \| 1,2,3 \| ns<br>Nonacosane \| oc \| 2900 \| 0.4 ± 0.03 \| 0.3 ± 0.01 \| 1,2,3 \| ns<br>Monoterpene hydrocarbons \| mh \| 57.8 \| 57.5<br>Oxygenated monoterpens \| om \| 6.5 \| 1.2<br>Sesquiterpene hydrocarbons \| sh \| 8.7 \| 17.9<br>Oxygenated sesquiterpenes \| os \| 3.7 \| 10.3<br>Diterpenes \| di \| 11.9 \| 7.8<br>Other constituents \| oc \| 1.6 \| 2.3<br>Total \| 90.2 \| 97.0 | **Table 1**<br>The main identified constituents of *J. macrocarpa* and *J. oxycedrus* essential oils. | Data are expressed as mean ± standard deviation (S.D.) ($n = 3$). *n.d.*: not detected. tr: trace (<0.1%). [a] Retention Index (RI) on HP-5 MS column. [b] IM: identification methods: 1—comparison of retention times; 2—comparison of mass spectra with MS libraries; 3—comparison with authentic compounds. Differences were evaluated by one-way analysis of variance (ANOVA) completed with a multiple comparison Tukey's test (** $p < 0.01$, * $p < 0.05$). ns: not significant. |
| https://europepmc.org/article/MED/35721323#free-full-text | Chemical Composition, Antioxidant, and Mosquito Larvicidal Activity of Essential Oils from Hyptis capitata Jacq. | https://europepmc.org/articles/PMC9205432/table/t0001/ | **Table 1**<br>Composition of Essential Oils of Inflorescence and Leaf of *H. capitata*<br>S. No. \| Compounds \| RI \| RI[L] \| Content % Leaf Oil \| Inflorescence Oil<br>1. α-Pinene 931.86 932 – 0.4<br>2. 1-Octen-3-ol 975.60 974 34.08 6.46<br>3. 3-Octanol 995.74 988 – 0.19<br>4. Benzene acetaldehyde 1,039.20 1,036 – 0.54<br>5. n-Octanol 1,067.89 1,063 – 0.65<br>6. Linalool 1,098.58 1,095 – 6.99<br>7. n-Nonanal 1,103.17 1,100 – 0.87<br>8. Hexenylbutanoate 1,190 1,193 – 0.24<br>9. n-Decanal 1,203.97 1,201 – 3.53<br>10. Hexenyl 2-methyl butanoate 1,228.50 1,232 – 0.4<br>11. Hexyl-2 methyl butanoate 1,233.87 1,233 – 1.29<br>12. Hexenylisovalerate 1,240.18 1,241 – 0.35<br>13. Methyl citronellate 1,265.65 1,257 2.57 0.89<br>14. n-Decanol 1,270.56 1,266 – 0.3<br>15. Longicyclene 1,365.80 1,371 – 0.88<br>16. α-Copaene 1,372.17 1,374 2.07 3.05<br>17. β-Bourbonene 1,379.71 1,387 0.88 –<br>18. β-Caryophyllene 1,414.56 1,417 5.24 3.48<br>19. n-Octyl2-methylbutanoate 1,431.06 1,431 – 4.53<br>20. Prenyl limonene 1,444.41 1,443 0.97 3.16<br>21. α-Humulene 1,449.75 1,452 0.74 1.33<br>22. Muurola-4 (14), 5-diene 1,467.23 1,465 – 1.91<br>23. β-Acoradiene 1,471.35 1,474 2.63 5.21<br>24. α-Neocallitropsene 1,474.51 1,474 – 2.96<br>25. Germacrene D 1,475.48 1,480 11.16 –<br>26. γ-Himachalene 1,482.76 1,481 – 1.15<br>27. β-Selinene 1,487.37 1,489 1.08 1.18<br>28. β-Guaiene 1,490.04 1,492 1.34 0.41<br>29. δ-Selinene 1,493.68 1,493 1.9 3.21<br>30. α-Muurolene 1,497.81 1,500 1.19 1.62<br>31. δ-Amorphene 1,507.41 1,511 2.48 3.61<br>32. δ-Cadinene 1,513.55 1,522 7.24 14.68<br>33. Cubebol 1,517.39 1,514 – 1.57<br>34. Zonarene 1,526.85 1,528 – 1.6<br>**Abbreviations**: RI, retention index; RI[L], RI from literature; –, not detected. | **Table 1**<br>Composition of Essential Oils of Inflorescence and Leaf of *H. capitata* | Abbreviations: RI, retention index; RIL, RI from literature; –, not detected. |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35807578#free-full-text | Chemical Diversity and Anti-Insect Activity Evaluation of Essential Oils Extracted from Five Artemisia Species. | https://europepmc.org/articles/PMC9269011/table/plants-11-01627-t001/ | **Table 1**<br>Chemical composition of essential oils extracted from the five *Artemisia* species.<br>[1] RI, retention index of the chromatography determined on a HP-5MS column using the homologous series of *n*-hydrocarbons |
| https://europepmc.org/article/MED/35048794#free-full-text | Chemical composition, enantiomeric analysis and anticholinesterase activity of Lepechinia betonicifolia essential oil from Ecuador. | https://europepmc.org/articles/PMC8786249/table/t0001/ | **Table 1.**<br>Chemical composition of the essential oil of *L. betonicifolia* in DB-5MS.<br>[a]Rt: Retention time; [b]LRI: calculated linear retention indices obtained on DB-5ms column using a series of *n*-alkanes (C9–C24) from [19]; [c]LRI: reference linear retention indices from [20]; %: relative percentage amount. | **Table 1.**<br>Chemical composition of the essential oil of *L. betonicifolia* in DB-5MS. | aRt: Retention time; bLRI: calculated linear retention indices obtained on DB-5ms column using a series of n-alkanes (C9–C24) from [19]; cLRI: reference linear retention indices from [20]; %: relative percentage amount. |
| https://europepmc.org/article/MED/35684405#free-full-text | Chemical Constituents, Antioxidant, Anti-Tyrosinase, Cytotoxicity, and Anti-Melanogenesis Activities of Etlingera elatior (Jack) Leaf Essential Oils. | https://europepmc.org/articles/PMC9182332/table/molecules-27-03469-t001/ | **Table 1**<br>Volatile constituents of *E. elatior* leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD).<br>Open in a separate window<br>The results are presented as area sum, %. | **Table 1**<br>Volatile constituents of *E. elatior* leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD). | The results are presented as area sum, %. |

**Table 1**

Chemical composition of essential oils extracted from the five *Artemisia* species.

| No. | RI [1] | Compound | Relative Content (%) [2] | | | | |
|---|---|---|---|---|---|---|---|
| | | | AD | AG | AS | AC | AO [3] |
| 1 | 800 | Octane | 1.24 | - | - | - | - |
| 2 | 908 | Santolina triene | 14.45 | - | - | - | - |
| 3 | 925 | *α*-Thujene | - | - | - | 2.53 | - |
| 4 | 966 | *β*-Thujene | - | - | - | 2.63 | - |
| 5 | 980 | *β*-Pinene | - | - | - | 15.13 | - |
| 6 | 1010 | 3-Carene | - | - | - | 45.98 | - |
| 7 | 1021 | *α*-Cymene | 1.51 | - | - | - | - |
| 8 | 1027 | Sylvestrene | - | - | - | 5.92 | - |
| 9 | 1046 | 1,8-Cineole | 32.62 | - | 0.36 | 2.56 | - |
| 10 | 1074 | 3,5-Dimethylethylbenzene | 7.48 | - | - | - | - |
| 11 | 1106 | Linalool | - | - | 1.34 | 1.27 | 0.56 |
| 12 | 1108 | 3,7-Dimethyl-1,5,7-octatriene-3-ol | 15.85 | - | - | - | - |
| 13 | 1145 | Camphor | - | 51.07 | 1.32 | - | 1.38 |
| 14 | 1160 | Borneol | - | - | 6.97 | - | - |
| 15 | 1164 | Pinocarvone | 0.46 | - | - | - | - |
| 16 | 1175 | 4-Terpineol | 2.02 | 11.97 | 1.57 | 1.12 | - |
| 17 | 1182 | Isocitral | - | 9.20 | - | - | - |
| 18 | 1190 | *α*-Terpineol | - | 1.47 | 3.69 | 1.27 | 0.92 |
| 19 | 1232 | *β*-Nerol | - | - | 11.01 | - | - |
| 20 | 1250 | *γ*-Pironene | - | - | - | - | 2.41 |
| 21 | 1267 | Geraniol | - | - | 1.44 | 0.31 | - |
| 22 | 1372 | Copaene | - | - | - | - | 1.76 |
| 23 | 1388 | *β*-Elemene | - | - | - | 1.54 | 5.56 |
| 24 | 1390 | *β*-Cubebene | - | - | 7.50 | - | 0.76 |
| 25 | 1396 | 3-Methyl-2-pent-2-enyl-cyclopent-2-enone | - | - | 1.09 | - | - |
| 26 | 1417 | Caryophyllene | 0.91 | 3.76 | 2.02 | 0.98 | 17.81 |
| 27 | 1425 | 1-Methyl-4-(1-methylethylidene)-2-(1-methylvinyl)-1-vinylcyclohexane | - | - | - | 1.29 | - |
| 28 | 1430 | Neryl propanoate | - | - | 22.88 | - | - |
| 29 | 1432 | Elixene | - | - | 4.21 | - | 0.19 |
| 30 | 1441 | *β*-Farnesene | - | - | 2.73 | - | 1.23 |
| 31 | 1464 | *β*-Humulene | - | - | - | - | 1.74 |
| 32 | 1465 | *γ*-Muurolene | - | - | - | - | 1.33 |
| 33 | 1480 | Germacrene D | - | - | - | 8.79 | 3.36 |
| 34 | 1489 | Viridiflorene | 1.14 | - | - | - | - |

[1] RI, retention index of the chromatography determined on a HP-5MS column using the homologous series of *n*-hydrocarbons

**Table 1.**

Chemical composition of the essential oil of *L. betonicifolia* in DB-5MS.

| Rt[a] | Compound | LRI[b] | LRI[c] | % | Σ |
|---|---|---|---|---|---|
| 8.08 | α-Thujene | 924 | 924 | 0.35 | 0.12 |
| 8.39 | α-Pinene | 932 | 932 | 4.97 | 1.32 |
| 9.08 | Camphene | 948 | 946 | 0.30 | 0.04 |
| 10.08 | Sabinene | 971 | 969 | 27.98 | 2.94 |
| 10.30 | *β*-Pinene | 976 | 974 | 30.45 | 2.25 |
| 10.79 | Myrcene | 988 | 988 | 0.34 | 0.10 |
| 11.46 | *p*-Mentha-1(7),8-diene | 1003 | 1003 | 0.48 | 0.03 |
| 11.57 | α-Phellandrene | 1006 | 1002 | 0.28 | 0.31 |
| 11.68 | delta-3-Carene | 1008 | 1008 | 1.08 | 1.20 |
| 12.07 | α-Terpinene | 1016 | 1014 | 0.37 | 0.48 |
| 12.43 | *p*-Cymene | 1023 | 1020 | 0.36 | 0.39 |
| 12.66 | Limonene | 1028 | 1024 | 3.84 | 0.72 |
| 12.72 | β-Phellandrene | 1029 | 1025 | 4.79 | 2.16 |
| 12.83 | 1,8-Cineole | 1032 | 1026 | 0.79 | 0.27 |
| 13.48 | (*E*)-*β*-Ocimene | 1045 | 1044 | 0.18 | 0.17 |
| 14.04 | γ-Terpinene | 1057 | 1054 | 0.22 | 0.19 |
| 14.66 | *p*-Mentha-2,4,8 diene | 1070 | 1085 | 0.38 | 0.52 |
| 15.34 | Terpinolene | 1084 | 1086 | 0.13 | 0.04 |
| 18.35 | Camphor | 1146 | 1141 | 0.20 | 0.06 |
| 19.07 | Pinocarvone | 1161 | 1160 | 0.47 | 0.53 |
| 19.96 | Terpinen-4-ol | 1179 | 1174 | 0.39 | 0.14 |
| 20.66 | Myrtenal | 1194 | 1195 | 0.73 | 0.80 |
| 28.84 | α-Copaene | 1374 | 1374 | 0.67 | 0.66 |
| 29.18 | *β*-Bourbonene | 1381 | 1387 | 0.19 | 0.23 |
| 30.70 | (*E*)-Caryophyllene | 1417 | 1417 | 4.44 | 0.91 |
| 31.15 | *β*-Gurjunene | 1427 | 1431 | 0.62 | 0.11 |
| 31.91 | Aromadendrene | 1445 | 1439 | 0.26 | 0.21 |
| 32.20 | α-Humulen | 1452 | 1452 | 0.85 | 0.44 |
| 32.37 | allo-Aromadendrene | 1456 | 1458 | 0.41 | 0.30 |
| 33.26 | γ-Muurolene | 1477 | 1478 | 2.84 | 2.68 |
| 33.71 | γ-Curcumene | 1488 | 1481 | 2.95 | 2.41 |
| 33.85 | Bicyclogermacrene | 1491 | 1500 | 0.50 | 0.36 |

[a]Rt: Retention time; [b]LRI: calculated linear retention indices obtained on DB-5ms column using a series of *n*-alkanes (C9–C24) from [19]; [c]LRI: reference linear retention indices from [20]; %: relative percentage amount.

**Table 1**


Volatile constituents of *E. elatior* leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD).

| Compounds | | CAS No. | HD | MAHD |
|---|---|---|---|---|
| **Aldehyde** | Decanal | 112-31-2 | 0.556 | 0.705 |
| | Undecanal | 112-44-7 | 0.203 | - |
| | Dodecanal | 112-54-9 | 10.024 | 12.422 |
| **Alkene** | 9-Octadecyne | 35365-59-4 | - | 0.725 |
| **Carboxylic ester** | 9-Tetradecen-1-ol, acetate, (E)- | 23192-82-7 | 0.249 | 0.919 |
| | 1-Tetradecyl acetate | 638-59-5 | 0.334 | 0.669 |
| | Lauryl acetate | 112-66-3 | 2.245 | 6.599 |
| **Dialkyl ketone** | 2-Undecanone | 112-12-9 | 0.274 | - |
| **Fatty acids** | Dodecanoic acid | 143-07-7 | 0.542 | - |
| | Hexadecanoic acid | 57-10-3 | 0.334 | 1.136 |
| **Fatty aldehyde** | cis,cis-7,10,-Hexadecadienal | 56829-23-3 | 0.351 | - |
| | 7-Tetradecenal, (Z)- | 65128-96-3 | - | 1.084 |
| | Pentadecanal- | 2765-11-9 | 0.264 | - |
| **Fatty alcohols** | 1-Decanol | 112-30-1 | 0.425 | 0.802 |
| | 1-Dodecanol | 112-53-8 | 13.484 | 30.335 |
| | 11-Hexadecen-1-ol, (Z)- | 56683-54-6 | - | 1.094 |
| | cis-9-Tetradecen-1-ol | 35153-15-2 | 1.989 | 6.522 |
| | 1-Tetradecanol | 112-72-1 | 1.224 | 5.711 |
| **Monoterpenes** | α-Pinene | 80-56-8 | 3.055 | 0.443 |
| | (±)-β-Pinene | 127-91-3 | 7.354 | 0.707 |
| | β-Myrcene | 123-35-3 | 1.524 | - |
| | (+)-m-Mentha-1(6),8-diene | 1461-27-4 | 0.291 | - |
| | Eucalyptol | 470-82-6 | 0.312 | - |
| | trans-β-Ocimene | 3779-61-1 | 0.154 | - |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35889501 | Preliminary Study on Phytochemical Constituents and Biological Activities of Essential Oil from Myriactis nepalensis Less. | https://europepmc.org/articles/PMC9324352/table/molecules-27-04631-t001/ | **Table 1**<br>Chemical composition of *M. nepalensis* essential oil.<br>Peak No. / Compound / RI [a] / RI [b] / % Area<br>1 (*E*)-2-Octenal 1050 1049 1.1<br>2 Linalool 1098 1095 0.6<br>3 *cis*-Verbenol 1141 1137 0.1<br>4 *α*-Terpineol 1190 1186 0.1<br>5 *cis*-Myrtanol 1252 1250 0.2<br>6 (*E*)-2-Decenal 1259 1260 0.1<br>7 Bornyl acetate 1285 1284 0.1<br>8 Dihydroedulan 1295 1293 0.2<br>9 Menthyl acetate 1297 1294 0.1<br>10 Isomenthyl acetate 1306 1304 0.2<br>11 *δ*-Elemene 1338 1335 0.1<br>12 7-epi-Silphiperfol-5-ene 1346 1345 0.1<br>13 *α*-Cubebene 1350 1348 0.1<br>14 (*E*)-2-Undecenal 1361 1357 0.1<br>15 Cyclosativene 1369 1369 0.1<br>16 *α*-Ylangene 1373 1373 0.1<br>17 *α*-Copaene 1378 1374 0.3<br>18 *β*-Bourbonene 1387 1387 0.7<br>19 *β*-Elemene 1393 1389 6.1<br>20 *cis*-*α*-Bergamotene 1415 1411 0.8<br>21 *β*-Caryophyllene 1420 1417 4.1<br>22 *γ*-Elemene 1435 1434 1.0<br>23 Aromadendrene 1442 1439 0.1<br>24 *α*-Himachalene 1448 1449 0.1<br>25 *α*-Humulene 1455 1452 3.2<br>26 Alloaromadendrene 1460 1458 0.1<br>27 *γ*-Muurolene 1478 1478 1.5<br>28 Valencene 1491 1496 1.4<br>29 Viridiflorene 1493 1496 2.4<br>30 *α*-Selinene 1499 1498 2.0<br>31 *α*-Muurolene 1502 1500 0.8<br>32 *β*-Bisabolene 1508 1505 0.5<br>33 *γ*-Cadinene 1517 1513 0.7<br>34 Cubebol 1519 1514 0.1<br>[a] Retention index calculated from n-alkanes ($C_7$-$C_{30}$) on an HP-5MS column; [b] Retention index data from the literature. | **Table 1**<br>Chemical composition of *M. nepalensis* essential oil. | a Retention index calculated from n-alkanes (C7-C30) on an HP-5MS column; b Retention index data from the literature. |
| https://europepmc.org/article/MED/35684405 | Chemical Constituents, Antioxidant, Anti-Tyrosinase, Cytotoxicity, and Anti-Melanogenesis Activities of Etlingera elatior (Jack) Leaf Essential Oils. | https://europepmc.org/articles/PMC9182332/table/molecules-27-03469-t001/ | **Table 1**<br>Volatile constituents of *E. elatior* leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD).<br>Compounds / CAS No. / HD / MAHD<br>Aldehyde Decanal 112-31-2 0.556 0.705<br>Undecanal 112-44-7 0.203 -<br>Dodecanal 112-54-9 10.024 12.422<br>Alkene 9-Octadecyne 35365-59-4 - 0.725<br>Carboxylic ester 9-Tetradecen-1-ol, acetate, (E)- 23192-82-7 0.249 0.919<br>1-Tetradecyl acetate 638-59-5 0.334 0.669<br>Lauryl acetate 112-66-3 2.245 6.599<br>Dialkyl ketone 2-Undecanone 112-12-9 0.274 -<br>Fatty acids Dodecanoic acid 143-07-7 0.542 -<br>Hexadecanoic acid 57-10-3 0.334 1.136<br>Fatty aldehyde cis,cis-7,10,-Hexadecadienal 56829-23-3 0.351 -<br>7-Tetradecenal, (Z)- 65128-96-3 - 1.084<br>Pentadecanal- 2765-11-9 0.264 -<br>Fatty alcohols 1-Decanol 112-30-1 0.425 0.802<br>1-Dodecanol 112-53-8 13.484 30.335<br>11-Hexadecen-1-ol, (Z)- 56683-54-6 - 1.094<br>cis-9-Tetradecen-1-ol 35153-15-2 1.989 6.522<br>1-Tetradecanol 112-72-1 1.224 5.711<br>Monoterpenes α-Pinene 80-56-8 3.055 0.443<br>(±)-β-Pinene 127-91-3 7.354 0.707<br>β-Myrcene 123-35-3 1.524 -<br>(+)-m-Mentha-1(6),8-diene 1461-27-4 0.291 -<br>Eucalyptol 470-82-6 0.312 -<br>trans-β-Ocimene 3779-61-1 0.154 -<br>Pinocamphone 547-60-4 0.162 -<br>3-Pinanone 15358-88-0 1.985 0.358<br>Terpinen-4-ol 562-74-3 0.152 -<br>α-Terpineol 98-55-5 0.692 -<br>Methyl myrtenate 30649-97-9 2.674 0.711<br>(+)-3-Carene, 10-(acetylmethyl)- 163886-28-0 1.064 0.438<br>Sesquiterpenes β-Elemene, (−)- 515-13-9 0.245 -<br>Caryophyllene 87-44-5 12.575 7.857<br>Humulene 6753-98-6 15.207 9.852<br>The results are presented as area sum, %. | **Table 1**<br>Volatile constituents of *E. elatior* leaf essential oils from hydrodistillation (HD) and microwave-assisted hydrodistillation (MAHD). | The results are presented as area sum, %. |
| https://europepmc.org/article/MED/35406974#free-full-text | The Phytochemical Profile and Anticancer Activity of Anthemis tinctoria and Angelica sylvestris Used in Estonian Ethnomedicine. | https://europepmc.org/articles/PMC9003001/table/plants-11-00994-t001/ | **Table 1**<br>Essential oil content of aerial parts of *Anthemis tinctoria*.<br>Compound / RI * (DB-5) / Content in Essential Oil (%)<br>Sabinene 972 0.7<br>β-Pinene 975 2.7<br>Myrcene 983 1.0<br>p-Cymene 1019 12.6<br>(Z)-β-Ocimene 1040 4.3<br>(E)-β-Ocimene 1048 2.6<br>Isoborneol 1147 2.1<br>Terpinen-4-ol 1174 4.0<br>Crysanthenyl acetate 1260 3.6<br>δ-Cadinene 1520 1.6<br>Caryophyllene oxide 1572 2.8<br>Isocaryophyllene oxide/caryophyllenol 1577 3.9<br>Humulene epoxide 1603 2.9<br>δ-Cadinol 1638 6.6<br>α-Muurolene 1648 12.5<br>2-Pentadecanone 1680 2.5<br>Nerolidol acetate 1720 2.5<br>n-Hexadecanal 1814 2.5<br>Palmitic acid 1967 15.3<br>Unknown 2534 12.6<br>Total 99.3<br>Monoterpenes 15.5<br>Cyclic monoterpenes 5.5<br>Bicyclic sesquiterpenes 6.7<br>Sesquiterpenes 24.5<br>Other compounds 47.1<br>* RI, retention index. | **Table 1**<br>Essential oil content of aerial parts of *Anthemis tinctoria*. | * RI, retention index. |


| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35406974#free-full-text | The Phytochemical Profile and Anticancer Activity of Anthemis tinctoria and Angelica sylvestris Used in Estonian Ethnomedicine. | https://europepmc.org/articles/PMC9003001/table/plants-11-00994-t002/ | **Table 2**<br>Essential oils content of aerial parts and roots of *Angelica sylvestris*.<br>Compound \| RI * (DB-5) \| Content in Essential Oils (%): Aerial Parts \| Roots<br>α-Pinene 933 45.4 12.4<br>Camphene 946 4.6 nf<br>Sabinene 971 1.1 nf<br>β-Pinene 974 2.0 nf<br>β-Myrcene 990 13.3 nf<br>α-Terpinen 1014 0.4 0.7<br>p-Cymene 1018 15.5 8.2<br>β-Phellandrene 1030 1.3 nf<br>(Z)-β-Ocimene 1040 1.1 0.9<br>(E)-β-Ocimene 1048 0.7 nf<br>Terpinolene 1085 1.2 nf<br>n-Nonanal 1108 nf 1.3<br>α-Terpineol 1188 nf 1.2<br>(E)-Verbenyl acetate 1301 1.4 nf<br>β-Elemene 1391 nf 1.2<br>(E)-β-Caryophyllene 1416 0.8 nf<br>β-Copaene 1424 nf 2.7<br>α-Humulene 1450 1.0 nf<br>β-Farnesene 1456 nf 2.2<br>Germacrene D 1477 3.2 nf<br>β-Bisabolene 1501 nf 1.5<br>γ-Cadinene 1507 0.8 nf<br>Cadina-1,4-diene 1536 nf 2.3<br>Elemol 1546 nf 3.4<br>Caryophyllene oxide 1573 nf 2.4<br>Isocaryophyllene oxide/caryophyllenol 1582 nf 31.9<br>Epiglobulol/humulene epoxide 1614 nf 1.4<br>α-Muurolene 1648 nf 4.2<br>α-Cadinol 1663 nf 1.7<br>α-Bisabolol 1680 1.4 17.5<br>Nerolidol acetate 1721 nf 1.3<br>Unknown 1857 nf 1.5<br>Palmitic acid 1965 3.3 nf<br>Total 99.9 99.9<br>* RI, retention index; nf, not found. | **Table 2**<br>Essential oils content of aerial parts and roots of *Angelica sylvestris*. | * RI, retention index; nf, not found. |
| https://europepmc.org/article/MED/35406974#free-full-text | The Phytochemical Profile and Anticancer Activity of Anthemis tinctoria and Angelica sylvestris Used in Estonian Ethnomedicine. | https://europepmc.org/articles/PMC9003001/table/plants-11-00994-t003/ | **Table 3**<br>Polyphenolic compounds identified in the methanolic extracts of *A. tinctoria* and *A. sylvestris* by HPLC-ion trap MS/MS.<br>$t_R$ (min) \| $[M-H]^-$ \| MS/MS \| Plant Material/Substance: Aerial Parts of *A. tinctoria* \| Aerial Parts of *A. sylvestris* \| Roots of *A. sylvestris*<br>0.5 341 179, 161 Caffeic acid glucosides Caffeic acid glucosides Caffeic acid glucosides<br>1.7 315 225, 153, 109 Protocatechuic or gentisic acid glucoside Protocatechuic or gentisic acid glucoside nf<br>4.3 325 163; 119 nf nf Coumaric acid glucoside<br>5.1 299 137 4-Hydroxybenzoic acid glucoside 4-Hydroxybenzoic acid glucoside nf<br>10.3 339 281, 251, 177, 135 Daphnin = daphnetin glucoside nf nf<br>12.5 353 191, 179, 173, 135 Neochlorogenic acid Neochlorogenic acid Neochlorogenic acid<br>15.2 385 223, 179, 163 Sinapinic acid glucoside nf nf<br>15.5 353 306, 191, 135 nf Chlorogenic acid nf<br>16.5 639 463, 301, 535 Quercetin glucoside glucuronide nf nf<br>16.7 337 191, 163, 173 nf Coumaroylquinic acid Coumaroylquinic acid<br>17.3 625 463, 301 Quercetin diglucoside nf nf<br>17.8 335 179, 135 Caffeoylshikimic acid nf nf<br>18.6 367 191, 173 5-Feruloylquinic acid 5-Feruloylquinic acid 5-Feruloylquinic acid<br>19.8? 479 317 Myricetin glucoside nf nf<br>20.9 625 301 nf Quercetinlucoside-glucoside nf<br>21.3 655 493, 331 Patuletin diglucoside nf nf<br>22.0 449 287, 151 Eriodictyol glucoside nf nf<br>22.2 741 609, 475, 343, 301 Quercetin rutinoside pentoside nf nf<br>22.6 477 301, 373 Quercetin glucuronide nf nf<br>22.8 463 301, 179, 343 Quercetin galactoside Quercetin galactoside nf<br>23.2 609 301, 343, 271 Rutin Rutin Rutin, traces<br>23.3 463 301 Quercetin glucoside Quercetin glucoside Quercetin glucoside<br>24.0 493 331, 373 Patuletin glucoside nf nf<br>24.9 477 315, 433 Isorhamnetin glucoside nf nf<br>24.9 505 463, 301, 179, 151 Quercetin acetyl glucoside nf nf<br>25.4 515 353, 191 Dicaffeoylquinic acid 1 Dicaffeoylquinic acid 1 nf<br>26.4 493 331, 287 Patuletin-7-glucoside nf nf<br>26.4 373 211, 193 Pinosylvin glucoside nf nf<br>26.7 477 315, 357, 300 Isorhamnetin glucoside Isorhamnetin glucoside nf<br>$t_R$, retention time; $[M-H]^-$, ion mass; MS/MS, mass of identified fragments; nf, not found. | **Table 3**<br>Polyphenolic compounds identified in the methanolic extracts of *A. tinctoria* and *A. sylvestris* by HPLC-ion trap MS/MS. | tR, retention time; [M-H]–, ion mass; MS/MS, mass of identified fragments; nf, not found. |
| https://europepmc.org/article/MED/35406974#free-full-text | The Phytochemical Profile and Anticancer Activity of Anthemis tinctoria and Angelica sylvestris Used in Estonian Ethnomedicine. | https://europepmc.org/articles/PMC9003001/table/plants-11-00994-t004/ | **Table 4**<br>Polyphenolic compounds identified in the methanolic extract of *A. tinctoria* and *A. sylvestris* by the UPLC–triple quadrupole MS/MS (MRM) method.<br>$t_R$ (min) \| Precursor Ion (*m/z*) $[M-H]^-$ \| Product Ion (*m/z*) MS/MS \| Aerial Parts of *A. tinctoria* \| Aerial Parts of *A. sylvestris* \| Roots of *A. sylvestris*<br>0.46 191 85 Quinic acid Quinic acid Quinic acid<br>2.80 353 191 Neochlorogenic acid Neochlorogenic acid Neochlorogenic acid<br>3.75 353 191 Chlorogenic acid Chlorogenic acid Chlorogenic acid<br>3.98 179 107 Caffeic acid Caffeic acid nf<br>5.06 609 300 Rutin Rutin nf<br>5.15 593 285 Luteolin 7-rutinoside Luteolin 7-rutinoside Luteolin 7-rutinoside<br>5.18 193 134 Ferulic acid Ferulic acid Ferulic acid<br>5.39/5.55/5.76 515 353 Dicaffeoylquinic acids Dicaffeoylquinic acids Dicaffeoylquinic acids<br>5.22 463 301 Hyperoside nf nf<br>5.26 447 285 Luteolin-7-glucoside nf nf<br>5.28 463 301 Isoquercitrin Isoquercitrin Isoquercitrin<br>5.52 623 315 Isorhamnetin 3-rutinoside nf nf<br>5.70 447 300 Quercitrin Quercitrin Quercitrin<br>5.70 447 285 Luteolin 4-glucoside nf nf<br>6.79 285 133 Luteolin nf nf<br>6.86 301 151 nf Quercetin nf<br>7.22 271 151 nf Naringenin Naringenin<br>7.38 269 117 Apigenin Apigenin Apigenin<br>7.57 299 284 nf Diosmetin Diosmetin<br>7.60 315 300 nf Isorhamnetin nf<br>$t_R$, retention time; $[M-H]^-$, ion mass; MS/MS, mass of identified fragments; nf, not found. | **Table 4**<br>Polyphenolic compounds identified in the methanolic extract of *A. tinctoria* and *A. sylvestris* by the UPLC–triple quadrupole MS/MS (MRM) method. | tR, retention time; [M-H]–, ion mass; MS/MS, mass of identified fragments; nf, not found. |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35707466#free-full-text | Comparative Analysis of Chemical Composition andAntibacterial and Anti-Inflammatory Activities of theEssential Oils from Chrysanthemum morifolium ofDifferent Flowering Stages and Different Parts. | https://europepmc.org/articles/PMC9192287/table/tab1/ | **Table 1**<br>Chemical composition (%) of JM, TJ, JH, SLC, and RC from *C. morifolium*.<br>Peak \| Compound[a] \| Content (%)[b] JM TJ JH SLC RC \| RI[c] \| RI[d]<br>1 \| *cis*-1,3-Dimethylcyclohexane \| 0.32 0.15 0.04 —[e] — \| 764 \| 772<br>2 \| (*E*)-2-Hexenal \| 0.08 — 0.03 — — \| 832 \| 850<br>3 \| 1,8-Cineole \| — 0.11 — — — \| 1037 \| 1031<br>4 \| *γ*-Terpinene \| — 0.04 — — — \| 1062 \| 1064<br>5 \| *cis*-Sabinol \| — 0.07 0.04 — — \| 1133 \| 1139<br>6 \| Camphor \| 0.1 0.03 0.12 — — \| 1144 \| 1146<br>7 \| *cis*-Chrysanthenol \| 0.11 0.06 0.04 — — \| 1156 \| 1163<br>8 \| Borneol \| 1.17 1.13 0.37 0.22 — \| 1171 \| 1172<br>9 \| Terpinen-4-ol \| 0.4 0.3 0.07 0.06 — \| 1185 \| 1180<br>10 \| Fenchyl acetate \| — — 0.06 — — \| 1221 \| 1225<br>11 \| Carvacrol \| 2.33 2.09 0.4 0.47 — \| 1291 \| 1291<br>12 \| *p*-Vinyl guaiacol \| — 0.1 0.08 0.27 0.17 \| 1314 \| 1309<br>13 \| *α*-Copaene \| — — — 0.2 — \| 1361 \| 1369<br>14 \| *β*-Elemene \| — 0.56 — 0.11 — \| 1385 \| 1388<br>15 \| Italicene \| 0.14 0.13 — 0.15 0.09 \| 1401 \| 1406<br>16 \| Caryophyllene \| 1.24 1.2 0.09 8.84 0.1 \| 1427 \| 1428<br>17 \| *trans*-*α*-Bergamotene \| — — — 0.13 — \| 1437 \| 1430<br>18 \| (E)-*β*-Famesene \| 0.16 0.37 — 1.14 0.11 \| 1455 \| 1455<br>19 \| *α*-Caryophyllene \| — 0.06 — 0.52 — \| 1462 \| 1456<br>20 \| (+)-Spathulenol \| 2.15 2.97 0.15 — — \| 1468 \| 1472<br>21 \| 7-epi-1,2-Dehydrosesquicineole \| 0.05 0.06 0.15 — — \| 1484 \| 1473<br>22 \| *β*-Selinene \| 0.87 1.09 0.34 — 0.05 \| 1488 \| 1485<br>23 \| *α*-Curcumene \| 1.82 3.63 — 2.69 — \| 1492 \| 1485<br>24 \| Germacrene D \| 0.11 — — 4.33 — \| 1494 \| 1487<br>25 \| (Z,E)-*α*-Farnesene \| 1.68 0.12 0.36 0.21 — \| 1498 \| 1495<br>26 \| Zingiberene \| — 2.94 — 3.63 — \| 1502 \| 1495<br>27 \| *α*-Muurolene \| 0.66 0.11 — 0.05 — \| 1504 \| 1502<br>28 \| *α*-Farnesene \| 0.2 0.79 0.08 1.53 — \| 1510 \| 1507<br>29 \| *γ*-Cadinene \| — 0.26 — 0.4 — \| 1513 \| 1516<br>30 \| *β*-Sesquiphellandrene \| 2.21 3.75 0.58 3.49 0.08 \| 1523 \| 1525<br>31 \| *γ*-Cadinene \| 0.3 0.32 0.09 — — \| 1533 \| 1534<br>32 \| (E)-*α*-Bisabolene \| 0.6 0.21 0.07 0.07 — \| 1540 \| 1540<br>[a]Compounds are listed according to their elution order on the apolar Elite-5MS capillary column. [b]Contents (%) were calculated by electronic integration of peak areas obtained on the apolar Elite-5MS column. [c]RI: Retention indices relatives to *n*-alkanes (C7–C27) on the apolar Elite-5MS column. [d]RI: Retention indices relatives were obtained from the literature (19–23) and NIST Chemistry WebBook. [e]–: not detected. [f]/: not identified and not found. | **Table 1**<br>Chemical composition (%) of JM, TJ, JH, SLC, and RC from *C. morifolium*. | aCompounds are listed according to their elution order on the apolar Elite-5MS capillary column. bContents (%) were calculated by electronic integration of peak areas obtained on the apolar Elite-5MS column. cRI: Retention indices relatives to n-alkanes (C7–C27) on the apolar Elite-5MS column. dRI: Retention indices relatives were obtained from the literature (19–23) and NIST Chemistry WebBook. e–: not detected. f/: not identified and not found. |
| https://europepmc.org/article/MED/35625543#free-full-text | Essential Oils of Duguetia Species A. St. Hill (Annonaceae): Chemical Diversity and Pharmacological Potential. | https://europepmc.org/articles/PMC9138787/table/biomolecules-12-00615-t002/ | **Table 2**<br>Main chemical constituents of *Duguetia* species EOs.<br>Species (Popular Name in Brazil) \| Part Used \| Total Components \| Major Constituents [1] \| References<br>*D. asterotricha* (envira, envira-surucucu-da-mata, envireira) \| Flowers \| 9 (69.3%) \| Limonene (14.1%), *p*-cymene (5.5%), and α-pinene (4.2%) \| [23]<br>*D. eximia* (unknown) \| Leaves and stem \| 76 ([4]) \| α-Eudesmol (80.3%) and spathulenol (5.0%) \| [24]<br>*D. flagellaris* (ameju-preto, caniceiro-preto, pindaíva, pindaíba) \| Bark \| 76 ([4]) \| Germacrene D (16.5%), cyperene (10.6%), α-muurolol (8.6%), humulene epoxide II (5.3%), spathulenol (5.0%), caryophyllene oxide (5.0%), δ-cadinene (4.3%), α-muurolene (4.2%), and β-elemene (4.0%) \| [24]<br>Stem \| 3 ([4]) \| Germacrene D (16.5%), cyperene (10.6%) and α-muurolol (8.6%) \| [25]<br>Leaves and Stem \| 76 ([4]) \| Spathulenol (58.7%), α-muurolol (6.2%) and humulene epoxide II (4.3%) \| [24]<br>Branches \| 2 ([4]) \| Spathulenol (58.7%), and α-muurolol (6.2%) \| [25]<br>*D. furfuracea* (araticum, ata brava, pinha do campo, pindaúva do campo, marolinho-do-cerrado, pinha-de-guará) \| Stem \| 19 ([4]) \| 2,4,5-Trimethoxystyrene (29.2%), α-asarone (23.8%), bicyclogermacrene (8.6%), *epi*-globulol (6.4%), and spathulenol (4.7%) \| [26]<br>39 ([4]) \| α-Gurjunene (22.2%), 2,4,5-trimethoxystyrene (19.7%), cyperene (16.0%), α-asarone (10.1%), and *trans*-*m*-mentha-4,8-diene (6.5%) \| [27]<br>24 ([4]) \| (*E*)-Asarone (21.9%), bicyclogermacrene (16.7%), 2,4,5-trimethoxystyrene (16.1%), α-gurjunene (15.0%), cyperene (7.8%), and (*E*)-caryophyllene (4.6%) \| [28]<br>8 ([4]) \| α-Asarone (36.4%), 2,4,5-trimethoxystyrene (27.8%), bicyclogermacrene (11.1%), α-gurjunene (10.5%), and cyperene (5.8%) \| [17]<br>Leaves [2] \| 17 (99.4%) \| β-Phellandrene (42.2%), bicyclogermacrene (20.7%), myrcene (6.8%), spathulenol (5.5%), α-phellandrene (4.6%), and sabinene (4.3%) \| [29]<br>17 (99.7%) \| Sabinene (25.1%), terpinen-4-ol (16.2%), *p*-cymene (8.3%), caryophyllene oxide (7.7%), and spathulenol (5.1%)<br>18 (99.2%) \| Bicyclogermacrene (29.1%), spathulenol (18.3%), germacrene D (9.6%), *trans*-caryophyllene (9.3%), δ-cadinene (5.5%), and caryophyllene oxide (5.3%)<br>18 (100%) \| Bicyclogermacrene (24%), germacrene D (15%), trans-caryophyllene (12.9%), spathulenol (12.4%), and caryophyllene oxide (6.8%)<br>19 (100%) \| Bicyclogermacrene (21.4%), germacrene D (13.6%), spathulenol (12.2%), and caryophyllene oxide (5.2%)<br>20 (99.7%) \| Terpinen-4-ol (21.6%), spathulenol (20.9%), sabinene (17.3%), and p-cymene (5.6%)<br>31 ([4]) \| Spathulenol (17.8%), bicyclogermacrene (16.2%), germacrene D (13.0%), β-caryophyllene (11.5%), and viridiflorol (4.0%) \| [27]<br>*D. gardneriana* \| Leaves \| 33 (91.4%) \| Germacrene D (28.1%), viridiflorene (24.0%), β-pinene (12.6%), \| [30]<br>[1] Only compounds with percentage concentrations greater than or equal to 4% are displayed. [2] The chemical compositions of the leaves vary due to the intensity of the odor and the location from which they were collected. [3] The extraction times for EO obtention (2 h and 4 h) resulted in different chemical compositions. [4] Percentage not described by the authors in relation to the total composition of the essential oil. | **Table 2**<br>Main chemical constituents of *Duguetia* species EOs. | 1 Only compounds with percentage concentrations greater than or equal to 4% are displayed. 2 The chemical compositions of the leaves vary due to the intensity of the odor and the location from which they were collected. 3 The extraction times for EO obtention (2 h and 4 h) resulted in different chemical compositions. 4 Percentage not described by the authors in relation to the total composition of the essential oil. |
| https://europepmc.org/article/MED/35625543#free-full-text | Essential Oils of Duguetia Species A. St. Hill (Annonaceae): Chemical Diversity and Pharmacological Potential. | https://europepmc.org/articles/PMC9138787/table/biomolecules-12-00615-t003/ | **Table 3**<br>Pharmacological properties of *Duguetia* species EOs.<br>Pharmacological Effects \| Part Used \| Actions \| References<br>**Anti-inflammatory**<br>*D. furfuracea* \| Stem \| After 6 h, EO inhibited paw edema induced by LPS by 92.4%. \| [28]<br>Stem \| After 2 h of LPS injection, doses of 3 and 10 mg/kg of EO inhibited paw edema by 90.9% and 92.42%, respectively. After 4 h, there was a significant reduction effect, with percentages of 77.8% (3 mg/kg) and 81.5% (10 mg/kg). \| [17]<br>*D. lanceolata* \| Bark \| EO at doses of 50, 100 and 200 mg/kg significantly reduced paw edema caused by carrageenan in 20.8%, 36.5% and 49.0%, respectively. \| [43]<br>Branches \| After 4 h, EO reduced the formation of paw edema caused by carrageenan by 18.3% (50 mg/kg), 32.3% (100 mg/kg) and 44.1% (200 mg/kg). \| [35]<br>**Antinociceptive**<br>*D. furfuracea* \| Stem \| EO inhibited formalin-induced activity, and caffeine (10 mg/kg) and naloxone (5 mg/kg) administration reversed the EO's antinociceptive activity. \| [28]<br>Stem \| Inhibition of 43.4% and 44.1% of formalin-induced activity was observed during the 1st phase at doses of 10 and 30 mg/kg, respectively. In the 2nd phase, there was also reduction in licking time at doses of 10 mg/kg (30.9%) and 30 mg/kg (39.8%). \| [17]<br>*D. lanceolata* \| Bark \| Number of abdominal contractions ($ED_{50}$ = 21.8 mg/kg) and paw-licking time 1st phase ($ED_{50}$ = 5.3 mg/kg) and 2nd phase ($ED_{50}$ = 1.4 mg/kg) were reduced in the formalin test. \| [43]<br>Branches \| In the formalin test, EO caused significant and time-dependent inhibition of paw licking at doses of 50, 100 and 200 mg/kg at 1st and 2nd phases. \| [35]<br>**Antibacterial and Antifungal**<br>*D. gardneriana* \| Leaves \| EO showed weak activity against *Staphylococcus aureus* and *Candida guilliermondii*. \| [30]<br>*D. lanceolata* \| Bark \| EO inhibited the growth of *Staphylococcus pyogenes*, *Escherichia coli* and *Candida albicans* with MIC values ranging from 20 to 125 μg/mL. \| [33]<br>*D. moricandiana* \| Leaves \| EO was active against *Staphylococcus aureus* and *Candida albicans* \| [30]<br>*D. quitarensis* \| Aerial parts \| EO was active for Gram-positive microorganisms *Streptococcus mutans* and *Streptococcus pyogenes* with MIC of 37.5 μg/mL. \| [37]<br>**Trypanocidal**<br>*D. quitarensis* \| Aerial parts \| EO showed trypanocidal activity against the amastigote and trypomastigote forms of *Trypanosoma cruzi* with $IC_{50}$ values of 0.26 and 0.54 μg/mL, respectively. \| [36]<br>**Antioxidant**<br>*D. lanceolata* \| Branches \| EO presented antioxidant effect, demonstrated through the DPPH radical, potency reduction and β-carotene assays. It inhibited lipid peroxidation by 41.5% ($EC_{50}$ equal to 159.4 μg/mL). \| [35]<br>**Cytotoxic and Antitumor**<br>*D. furfuracea* \| Stem \| EO was active against *A. salina* with $LC_{50}$ values of 2.6 μg/mL. \| [26] | **Table 3**<br>Pharmacological properties of *Duguetia* species EOs. | |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35744789 | Effect of Ylang-Ylang (Cananga odorata Hook. F. & Thomson) Essential Oil on Acute Inflammatory Response In Vitro and In Vivo. | https://europepmc.org/articles/PMC9231162/table/molecules-27-03666-t001/ | **Table 1**<br>Chemical composition of *Cananga odorata* essential o<br>RT: retention time; RI: retention index on Rtx-5MS. | **Table 1**<br>Chemical composition of *Cananga odorata* essential oil. | RT: retention time; RI: retention index on Rtx-5MS. |
| | Ocimum sanctum, Zingiber officinale, and Piper nigrum extracts and their effects on gut microbiota modulations (prebiotic potential), basal inflammatory markers and lipid levels: oral supplementation study in healthy rats. | https://europepmc.org/articles/PMC8865113/table/t0002/ | **Table 2.**<br>Polyphenols in *O.sanctum, Z.officinale, P.nigrum* and combined extracts.<br>*Combination (animal dose: 1450 mg/kg.bwt) = *O. sanctum* (850 mg [58.6%]) + *Z. officinale* (500 mg [34.6%]) + *P. nigrum* (100 mg [7.0%]; BDL: Below Detectable Level. | **Table 2.**<br>Polyphenols in *O.sanctum*, *Z.officinale*, *P.nigrum* and combined extracts. | *Combination (animal dose: 1450 mg/kg.bwt) = O. sanctum (850 mg [58.6%]) + Z. officinale (500 mg [34.6%]) + P. nigrum (100 mg [7.0%]; BDL: Below Detectable Level. |
| | Ocimum sanctum, Zingiber officinale, and Piper nigrum extracts and their effects on gut microbiota modulations (prebiotic potential), basal inflammatory markers and lipid levels: oral supplementation study in healthy rats. | https://europepmc.org/articles/PMC8865113/figure/F0002a/ |  | Figure 2.<br>Chemical composition of essential oils of *Ocimum sanctum*, *Zingiber officinale* and *Piper nigrum*. (2a) GC-MS Chromatogram of *O. sanctum* essential oils. (b) GC-MS Chromatogram of *Zingiber officinale* essential oils. (c) GC-MS Chromatogram of *P. nigrum* essential oils. (d) GC-MS Chromatogram of *O. sanctum* + *Z. officinale* +*P. nigrum* essential oils. | |

| Peak | RT (min) | Compound | RI | % |
|---|---|---|---|---|
| 1 | 4.57 | prenyl acetate | 921 | 0.41 |
| 2 | 7.10 | *p*-methyl anisole | 1019 | 7.38 |
| 3 | 7.45 | 1,8-cineole | 1030 | 0.05 |
| 4 | 9.62 | methyl benzoate | 1096 | 7.64 |
| 5 | 9.79 | linalool | 1100 | 15.23 |
| 6 | 12.28 | benzyl acetate | 1164 | 18.12 |
| 7 | 15.99 | linalyl acetate | 1255 | 0.21 |
| 8 | 20.91 | α-copaene | 1374 | 0.38 |
| 9 | 21.33 | geranyl acetate | 1384 | 9.46 |
| 10 | 21.60 | β-elemene | 1390 | 0.09 |
| 11 | 22.68 | *trans*-caryophyllene | 1417 | 5.42 |
| 12 | 23.82 | cinnamyl acetate | 1445 | 6.05 |
| 13 | 24.05 | α-humulene | 1451 | 1.80 |
| 14 | 25.01 | γ-muurolene | 1475 | 0.42 |
| 15 | 25.17 | germacrene D | 1479 | 4.61 |
| 16 | 25.60 | prenyl benzoate | 1489 | 0.17 |
| 17 | 25.79 | bicyclogermacrene | 1494 | 0.13 |
| 18 | 25.98 | α-muurolene | 1499 | 0.21 |
| 19 | 26.12 | γ-bisabolene | 1502 | 0.59 |
| 20 | 26.32 | α-farnesene | 1508 | 2.02 |
| 21 | 26.49 | γ-cadinene | 1512 | 0.22 |
| 22 | 26.87 | δ-cadinene | 1522 | 0.98 |
| 23 | 29.12 | caryophyllene oxide | 1580 | 0.06 |
| 24 | 31.40 | α-cadinol | 1641 | 0.44 |
| 25 | 31.56 | α-muurolol | 1645 | 0.10 |
| 26 | 31.86 | cadin-4-en-10-ol | 1654 | 0.51 |
| 27 | 34.36 | (*z,z*)-farnesol | 1722 | 0.38 |
| 28 | 35.80 | benzyl benzoate | 1764 | 11.39 |
| 29 | 38.47 | farnesyl acetate | 1841 | 0.83 |
| 30 | 39.34 | benzyl salicylate | 1868 | 4.47 |

| Polyphenols (mg/100gm) (Mean ± SE) | *O. sanctum* | *Z. officinale* | *P. nigrum* | Combination* |
|---|---|---|---|---|
| Gallic acid | 4.02 ± 0.12 | 84.13 ± 1.2 | 6.49 ± 0.52 | 12.91 ± 0.47 |
| Protocatechuic acid | 23.53 ± 0.1.2 | 19.26 ± 0.32 | 15.38 ± 0.25 | 22.18 ± 0.62 |
| *p*-hydroxi benzoic acid | 12.61 ± 0.32 | 0.77 ± 0.11 | 3.35 ± 0.42 | 8.59 ± 0.43 |
| Catechin | 356 ± 2.02 | 9.70 ± 0.14 | 28.15 ± 0.62 | 194 ± 1.66 |
| Caffeic acid | 55.68 ± 1.21 | 1.43 ± 0.01 | 3.92 ± 0.23 | 33.32 ± 1.32 |
| Sinapic acid | 18.58 ± 0.12 | 0.95 ± 0.06 | 2.54 ± 0.31 | 12.41 ± 0.91 |
| Ferulic acid | 7.17 ± 0.23 | 2.82 ± 0.35 | 10.45 ± 0.64 | 8.17 ± 0.33 |
| *p*-Coumaric acid-4 | BDL | 3.84 ± 0.11 | 0.80 ± 0.01 | 0.82 ± 0.04 |
| Ellagic acid | 18.14 ± 0.21 | 14.64 ± 10.62 | 2.47 ± 0.26 | 11.39 ± 0.62 |
| *o*-Coumaric acid-2 | 10.19 ± 0.96 | 7.06 ± 0.14 | 1.60 ± 0.02 | 5.22 ± 0.34 |
| Luteolin-7*O*-Glucoside | 430 ± 2.3 | BDL | BDL | 266 ± 1.8 |
| Myricetin | 43.70 ± 1.2 | BDL | 2.52 ± 0.34 | 27.52 ± 0.56 |
| Resveratrol | 270 ± 2.1 | 2.72 ± 0.12 | BDL | 164 ± 2.8 |
| Daidzein | BDL | BDL | 2.26 ± 0.41 | 3.13 ± 0.24 |
| Quercetin | BDL | BDL | BDL | BDL |
| Luteolin | 64.10 ± 1.02 | BDL | 2.23 ± 0.61 | 39.09 ± 0.99 |
| Naringenin | 6.43 ± 0.31 | BDL | 24.95 ± 1.51 | 12.99 ± 0.53 |
| Apigenin | 132 ± 3.11 | BDL | BDL | 79.57 ± 1.65 |
| Kaempferol | 57.03 ± 1.21 | BDL | BDL | 36.78 ± 0.87 |
| Hesperetin | 164 ± 2.31 | BDL | 7.06 ± 0.97 | 93.66 ± 2.56 |
| Flavone | BDL | 21.89 ± 0.91 | 88.18 ± 1.8 | 33.36 ± 1.65 |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| | Ocimum sanctum, Zingiber officinale, and Piper nigrum extracts and their effects on gut microbiota modulations (prebiotic potential), basal inflammatory markers and lipid levels: oral supplementation study in healthy rats. | https://europepmc.org/articles/PMC8865113/figure/F0002b/ |  | Figure 2.<br>Chemical composition of essential oils of *Ocimum sanctum*, *Zingiber officinale* and *Piper nigrum*. (2a) GC-MS Chromatogram of *O. sanctum* essential oils. (b) GC-MS Chromatogram of *Zingiber officinale* essential oils. (c) GC-MS Chromatogram of *P. nigrum* essential oils. (d) GC-MS Chromatogram of *O. sanctum* + *Z. officinale* +*P. nigrum* essential oils. | |
| https://europepmc.org/article/MED/35883828 | Valeriana pilosa Roots Essential Oil: Chemical Composition, Antioxidant Activities, and Molecular Docking Studies on Enzymes Involved in Redox Biological Processes. | https://europepmc.org/articles/PMC9311991/table/antioxidants-11-01337-t001/ | **Table 1**<br>Percentage composition of the essential oil isolated from *Valeriana pilosa* R and P roots collected in Cajamarca, Peru.<br>[a] RI—retention index as determined on the DB-1 column using the homologous series of *n*-alkanes ($C_9$–$C_{21}$); t—trace (<0.05). [b] RI data—retention index data reported in plant essential oils on non-polar column (www.webbook.nist.gov, accessed on 21 March 2022). | **Table 1**<br>Percentage composition of the essential oil isolated from *Valeriana pilosa* R and P roots collected in Cajamarca, Peru. | a RI—retention index as determined on the DB-1 column using the homologous series of n-alkanes (C9–C21); t—trace (<0.05). b RI data—retention index data reported in plant essential oils on non-polar column (www.webbook.nist.gov, accessed on 21 March 2022). |
| https://europepmc.org/article/MED/35630757#free-full-text | Phytochemical Analysis, Antibacterial Activity and Modulating Effect of Essential Oil from Syzygium cumini (L.) Skeels. | https://europepmc.org/articles/PMC9145283/table/molecules-27-03281-t001/ | **Table 1**<br>Chemical composition of the essential oil from leaves of *Syzygium cumini* (L.) Skeels.<br>Open in a separate window<br>Source: Research data; [a] experimental retention index (based on the n-alkane $C_7$–$C_{30}$ homologous series); [b] literature retention index [19]. | **Table 1**<br>Chemical composition of the essential oil from leaves of *Syzygium cumini* (L.) Skeels. | Source: Research data; a experimental retention index (based on the n-alkane C7–C30 homologous series); b literature retention index [19]. |

| N° | Components | RI [a] | Relative Content (%) | Identification Method | RI Data [b] |
|---|---|---|---|---|---|
| 1 | Isovaleric acid | 886 | 2.6 | RI,MS | 827–888 |
| 2 | Tricyclene | 921 | t | RI,MS | 914–930 |
| 3 | α-Thujene | 924 | 0.1 | RI,MS | 905–948 |
| 4 | α-Pinene | 930 | 3.7 | RI,MS | 909–956 |
| 5 | Camphene | 938 | 1.4 | RI,MS | 929–978 |
| 6 | 3-Methyl valeric acid | 947 | 3.1 | RI,MS | 941–968 |
| 7 | Sabinene | 958 | 0.4 | RI,MS | 944–980 |
| 8 | 1-Octen-3-ol | 961 | t | RI,MS | 958–986 |
| 9 | β-Pinene | 963 | 0.6 | RI,MS | 952–986 |
| 10 | Myrcene | 975 | 0.1 | RI,MS | 962–993 |
| 11 | Limonene | 1009 | 3.2 | RI,MS | 995–1044 |
| 12 | p-Cymene | 1013 | t | RI,MS | 992–1072 |
| 13 | 1,8-Cineole | 1015 | 4.3 | RI,MS | 1007–1046 |
| 14 | Linalool | 1074 | 0.1 | RI,MS | 1078–1107 |
| 15 | Isopentyl isovalerate | 1094 | t | RI,MS | 1094–1105 |
| 16 | Camphor | 1102 | 0.2 | RI,MS | 1105–1150 |
| 17 | Menthone | 1120 | 0.8 | RI,MS | 1124–1142 |
| 18 | Isomenthone | 1126 | 0.2 | RI,MS | 1132–1159 |
| 19 | Borneol | 1134 | t | RI,MS | 1140–1188 |
| 20 | Neomenthol | 1139 | t | RI,MS | 1153–1176 |
| 21 | Menthol | 1148 | 1.2 | RI,MS | 1141–1185 |
| 22 | Carvone | 1210 | 0.1 | RI,MS | 1210–1246 |
| 23 | Menthyl acetate | 1278 | 1.4 | RI,MS | 1276–1294 |
| 24 | α-Cubebene | 1345 | 0.2 | RI,MS | 1340–1360 |
| 25 | Cyclosativene | 1363 | 0.1 | RI,MS | 1363–1368 |
| 26 | α-Copaene | 1375 | 1.0 | RI,MS | 1351–1407 |
| 27 | β-Patchoulene | 1378 | 0.4 | RI,MS | 1375–1380 |
| 28 | β-Bourbonene | 1379 | 0.4 | RI,MS | 1346–1396 |
| 29 | β-Elemene | 1388 | 0.8 | RI,MS | 1362–1410 |
| 30 | β-Caryophyllene | 1414 | 3.5 | RI,MS | 1411–1421 |
| 31 | Seychellene | 1431 | 7.6 | RI,MS | 1457–1461 |
| 32 | α-Guaiene | 1437 | 4.1 | RI,MS | 1409–1490 |
| 33 | α-Humulene | 1447 | 6.1 | RI,MS | 1428–1489 |

| Compounds | RI [a] | RI [b] | % |
|---|---|---|---|
| α-pinene | 937 | 939 | 53.21 |
| β-pinene | 979 | 981 | 3.01 |
| β-myrcene | 995 | 991 | 0.75 |
| Limonene | 1029 | 1031 | 1.25 |
| Nonalol | 1105 | 1103 | 5.62 |
| Linalool | 1099 | 1098 | 3.98 |
| α-terpineol | 1187 | 1189 | 2.09 |
| Tetradecane | 1226 | 1221 | 0.27 |
| Nerol | 1228 | 1228 | 9.38 |
| (E,Z)-2,4-decadienal | 1296 | 1295 | 0.98 |
| Geranil acetate | 1385 | 1384 | 3.43 |
| Ionone | 1387 | 1387 | 1.29 |
| Damascone | 1409 | 1411 | 1.08 |
| Caryophyllene | 1417 | 1418 | 2.81 |
| α-humulene | 1451 | 1452 | 1.57 |
| Nerolidol | 1569 | 1564 | 5.73 |
| Globulol | 1581 | 1583 | 0.06 |
| α-cadinol | 1646 | 1649 | 0.11 |
| Total identified (%) | | | 96.62 |

| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35140337#free-full-text | Chemical composition, antioxidant, and antimicrobial activity of Elsholtzia beddomei C. B. Clarke ex Hook. f. essential oil. | https://europepmc.org/articles/PMC8828703/table/Tab1/ | **Table 1**<br>Chemical constituents of *E. beddomei* essential oil.<br>No. Compound RI[a] %Peak area<br>1 2E-hexenal 854 t[b]<br>2 hexanol 872 t<br>3 benzaldehyde 962 0.31<br>4 endo-2-norborneol 991 0.07<br>5 cis-meta-mentha-2,8-diene 993 0.06<br>6 benzyl alcohol 1036 t<br>7 bergamal 1059 t<br>8 acetophenone 1070 0.81<br>9 linalool 1106 83.67<br>10 perillene 1113 1.65<br>11 exo-isocitral 1151 t<br>12 trans-α-necrodol 1155 t<br>13 β-pinene oxide 1166 0.12<br>14 lavandulol 1177 t<br>15 rosefuran epoxide 1185 t<br>16 Z-isocitral 1189 0.15<br>17 verbanol 1210 0.16<br>18 elsholtzia ketone 1214 0.24<br>19 trans-pulegol 1225 t<br>20 nerol 1239 0.33<br>21 exo-fenchyl acetate 1241 t<br>22 neral 1247 3.68<br>23 trans-chrysanthenyl acetate 1247 0.26<br>24 geraniol 1261 0.11<br>25 perillaldehyde 1282 4.68<br>26 nerylformate 1293 t<br>27 methyl geranate 1335 0.06<br>28 eugenol 1370 t<br>29 neryl acetate 1373 0.05<br>30 α-ylangene 1387 t<br>31 geranyl acetate 1393 t<br>32 p-but-E-enyl-anisole 1394 t<br>33 E-caryophyllene 1431 1.55<br>34 α-trans-bergamotene 1446 t<br>[a]Calculated retention indices obtained from a DB-5 column.<br>[b]Trace amount < 0.05. | **Table 1**<br>Chemical constituents of *E. beddomei* essential oil. | [a]Calculated retention indices obtained from a DB-5 column.<br>[b]Trace amount<0.05. |
| https://europepmc.org/article/MED/35677362 | Chemical Composition, Antioxidant, Insecticidal Activity, and Comparative Analysis of Essential Oils of Leaves and Fruits of Schinus molle and Schinus terebinthifolius. | https://europepmc.org/articles/PMC9170424/table/tab2/ | **Table 2**<br>Chemical composition of different essential oils.<br>Compound name Retention indices *Schinus molle* *Schinus terebinthifolius*<br>RI I RI a SML SMF STL STF<br>α-Pinene 936 934 — **0.70** — **3.26**<br>Camphene 950 948 **0.21** **2.22** **0.91** —<br>β-Pinene 978 974.96 **0.24** **1.21** **1.48** **0.25**<br>β-Myrcene 987 983 **10.36** **5.44** **4.22** **0.70**<br>(+)-2-Carene 996 996 **3.19** — — —<br>α-Phellandrene 998 1001 **0.22** **0.75** **0.96** —<br>α-Terpinene 1013 1014 **12.01** **8.15** **9.45** **6.70**<br>Limonene 1025 1026 **22.94** **18.49** **23.22** **6.52**<br>β-Ocimene 1027 1028 — **5.04** **13.32** —<br>γ-Terpinene 1051 1052 **12.01** **8.15** **9.45** **6.70**<br>Terpinolene 1086 1082 **0.50** **0.61** **2.09** **0.77**<br>Linalool 1086 1085.36 — **0.23** — —<br>α-Thujone 1087 1089 — **0.34** — —<br>α-Campholenal 1105 1110 — — **0.20** —<br>Sabinol 1120 1121 **2.52** **1.57** **5.07** **0.33**<br>Trans-p-menth-2-en-1-ol 1123 1127 **0.75** — **0.62** —<br>Octanoic acid methyl ester 1125 1126 **2.26** — — —<br>Verbenol 1128 1130 — — **0.28** —<br>Para-cymen-8-ol 1162 1163 **0.28** — **0.77** —<br>Terpinen-4-ol 1164 1165.28 **0.44** **0.36** **0.37** —<br>Pinocarveol 1184 1187 — — **0.40** —<br>Carvone 1214 1219 — — **0.27** —<br>Piperitone 1232 1231 — — **0.35** —<br>Carvenone 1236 1234 **0.89** **0.24** **1.26** —<br>Phellandral 1249 1250 **0.30** — **0.42** —<br>Thymol 1267 1278 **1.33** **0.82** **1.75** —<br>Bornyle acetate 1273 1279 **0.29** **0.52** — —<br>Carvacrol 1278 1286 — — — **0.37**<br>Citronellol acetate 1334 1337 — — **2.25** —<br>Geranyl acetate 1358 1357 **0.30** — **0.50** —<br>β-Elemene 1389 1389.71 **0.47** **1.39** **0.26** **8.15**<br>α-bisabolol 1665 1666 **0.61** — — —<br>Caryophyllene oxide 1578 1573 — **0.34** **0.28** **0.22** | **Table 2**<br>Chemical composition of different essential oils. | |
| https://europepmc.org/article/MED/34950215#free-full-text | Lamiaceae Essential Oils, Phytochemical Profile, Antioxidant, and Biological Activities. | https://europepmc.org/articles/PMC8692021/table/tab1/ | **Table 1**<br>Chemical composition and yield of essential oils from *Lamiaceae* species.<br>Species Plant part Essential oil yield Main compounds Reference<br>*Aeollanthus suaveolens* Mart. ex Spreng Leaves 1.6% Massoia lactone, linalool,(*E*)-B-farnesene, geraniol, 2,5-dimethoxy-p-cymene [95]<br>*Calamintha nepeta* (L.) Kuntze Leaves * Pulegone, menthone, limonene [96]<br>*Clinopodium macrostemum* (Moc. and Sessé ex Benth.) Kuntze Leaves 0.80% linalool, nerol, caryophyllene, menthone, geranyl acetate, terpineol, pulegone [97]<br>*Hyptis dilatata* Benth. Leaves * Fenchone, 3-carene, α-pinene, β-caryophyllene, limonene, β-pinene, and camphor [98]<br>*Hyptis martiusii* Benth. Leaves 0.34% 1,8-cineole, d-carene, camphor, limonene, germacrene B [99]<br>*Lavandula dentata* L. Leaves and stems * 1,8-cineole, isolimonene, thuj-3-en-10-al, trans-pinocarveol [100]<br>*Melissa officinalis* L. Leaves and flowers 0.10% Citral, caryophyllene oxide, citronellal, geraniol, geranyl acetate, β-caryophyllene [101]<br>*Mentha arvensis* L. Leaves and flowers * Citronellal and nerol [102]<br>*M. piperita* L Leaves and flowers * Menthone, menthol, pulegone and menthyl acetate [103]<br>*Minthostachys mollis* (Benth.) Griseb. Leaves 0.98% Menthone, pulegone, cis-dihydrocarvone, carvacryl acetate, linalyl acetate, and linalool [104]<br>*Ocimum basilicum* L. Leaves 1.56 ± 0.15% linalyl acetate and linalool [105]<br>*O. gratissimum* L. Leaves * 1,8-Cineole, eugenol, 4-terpineol [91]<br>*O. gratissimum* Leaves and flowers * Thymol, eugenol, 1,8-cineole, E-caryophyllene, β-selinene [106]<br>*Origanum scabrum* Boiss. and Heldr. Leaves 1.5% Carvacrol, thymol, p-cymene,γ-terpinene [107]<br>*Origanum vulgare* L. Leaves * 4-terpineol, sabinene hydrate, thymol [92]<br>*Plectranthus amboinicus* (Lour.) Spreng. Leaves 0.009% Thymol, β-pinene, γ-terpinene, caryophyllene [108]<br>*Plectranthus barbatus var. grandis* (L.H. Cramer) Lukhoba and A.J. Paton Leaves * β-caryophyllene, α-copaene, germacrene [109] | **Table 1**<br>Chemical composition and yield of essential oils from *Lamiaceae* species. | |


| URL | Title | Table Link | Table Pic (top) | Table Header | Table footer |
|---|---|---|---|---|---|
| https://europepmc.org/article/MED/35449437#free-full-text | Coriander (Coriandrum sativum L.) essential oil and oil-loaded nano-formulations as an anti-aging potentiality via TGFβ/SMAD pathway. | https://europepmc.org/articles/PMC9023561/table/Tab1/ |  | **Table 1**<br><br>Volatile oil constituents identified in the essential oils of fennel, anise, coriander, and cumin fruits were analyzed by GC–MS. | Significant values are in bold. |

**Table 1**

Volatile oil constituents identified in the essential oils of fennel, anise, coriander, and cumin fruits were analyzed by GC–MS.

| Compounds | (RI) | Rel. abundance (%) | | | | IM |
|---|---|---|---|---|---|---|
| | | Fennel | Anise | Coriander | Cumin | |
| α-Pinene | 939 | 1.58 | – | **4.58** | 0.72 | RI, MS |
| Camphene | 954 | 0.04 | – | 0.48 | – | RI, MS |
| Sabinene | 954 | 0.34 | – | 0.16 | 0.8 | RI, MS |
| β-Pinene | 979 | 0.14 | – | 0.39 | **14.43** | RI, MS |
| Myrcene | 990 | 0.2 | – | 0.13 | 0.53 | RI, MS |
| Carene | 1002 | 0.07 | – | – | 0.07 | RI, MS |
| Cymene | 1024 | 0.15 | – | 0.49 | **4.56** | RI, MS |
| Limonene | 1029 | **12.5** | – | 1.05 | – | RI, MS |
| 1,8-Cineole | 1031 | 0.18 | – | – | 0.31 | RI, MS |
| *O*-Cimene | 1037 | 0.37 | – | – | – | RI, MS |
| Terpinene | 1059 | 0.19 | – | **3.64** | **17.33** | RI, MS |
| Camphor | 1146 | 0.1 | – | **3.61** | | RI, MS |
| Methyl chavicol | 1196 | **79.88** | – | – | | RI, MS |
| Fenchyl acetate | 1220 | 0.1 | – | – | – | RI, MS |
| α-Copaene | 1376 | 0.03 | – | – | – | RI, MS |
| Germacrene | 1481 | 0.03 | – | – | – | RI, MS |
| Anethole | 1284 | – | **91.8** | – | – | RI, MS |
| Element | 1338 | – | 0.23 | – | – | RI, MS |
| α-Himachalene | 1451 | – | 0.43 | – | – | RI, MS |
| β-Himachalene | 1482 | – | **5.33** | – | – | RI, MS |
| α-Longipinene | 1352 | – | 0.04 | – | – | RI, MS |
| Cyclosativene | 1371 | – | 0.02 | – | – | RI, MS |
| α-Ylangene | 1375 | – | 0.07 | – | – | RI, MS |
| β-Elemene | 1390 | – | 0.09 | – | – | RI, MS |
| α- zingiberene | 1493 | – | 0.75 | – | – | RI, MS |
| β- Himachalene | 1500 | – | 0.31 | – | – | RI, MS |
| β-Bisabolene | 1505 | – | 0.25 | – | – | RI, MS |
| β-Sesquiphellandrene | 1522 | – | 0.06 | – | – | RI, MS |
| α-Thujene | 930 | 0.02 | – | 0.03 | 0.22 | RI, MS |
| Terpinolene | 1088 | – | – | 0.13 | – | RI, MS |
| Linalool | 1096 | – | – | **81.29** | – | RI, MS |
| Citronellal | 1153 | – | – | 0.14 | – | RI, MS |

Significant values are in bold.