

Top 1D.H Geo Prelim Exam August 2011.

- 1) (a) We want to show that X is Hausdorff. Consider the following diagram

$$\begin{array}{ccc} \mathbb{R}^2 & & \\ \downarrow q & \searrow f & \\ X = \mathbb{R}/\sim & \xrightarrow{g} & \mathbb{R} \end{array} \quad x \sim y \text{ if } f(x) = f(y)$$

Since f is constant on the fibers $q^{-1}([x])$, we know that f induces a continuous map g st. $g \circ q = f$. Now consider $[x], [y] \in X$ with $[x] \neq [y]$. Then $g([x]) \neq g([y])$ and since $g([x]), g([y]) \in \mathbb{R}$ which is Hausdorff, \exists open sets U, V in \mathbb{R} with $g([x]) \in U$, $g([y]) \in V$, and $U \cap V = \emptyset$. Well, g is continuous, so $g^{-1}(U)$ and $g^{-1}(V)$ are open, $[x] \in g^{-1}(U)$, $[y] \in g^{-1}(V)$, and $g^{-1}(U) \cap g^{-1}(V) = \emptyset$. Thus, X is Hausdorff.

- (b) Suppose that X is not connected so $X = U \sqcup V$ with U and V open. Then $\mathbb{R}^2 = q^{-1}(U) \sqcup q^{-1}(V)$, so \mathbb{R}^2 is disconnected. \square

Top/Diff Geo Prelim Exam August 2011


- 2) Suppose there exists a homeomorphism $f: D^2 \rightarrow D^2$ such that $f|_{S^1}$ is not a homeomorphism. Then, there exists some $s \in S^1$ such that $f(s) \notin S^1$. If we then consider the spaces $D^2 \setminus \{x\}$ and $D^2 \setminus \{f(x)\}$, there is an induced homeomorphism $\tilde{f}: D^2 \setminus \{x\} \rightarrow D^2 \setminus \{f(x)\}$. Consider some fixed point $y \in D^2$, $x \neq y \neq f(x)$. We know that since $x \in \text{int}(D^2)$ and $f(x) \in \text{int}(D^2)$, $D^2 \setminus \{x\}$ is homotopic to D^2 and $D^2 \setminus \{f(x)\}$ is homotopic to S^1 , so

$$\pi_1(D^2 \setminus \{x\}, y) = 1, \quad \pi_1(D^2 \setminus \{f(x)\}, y) = \mathbb{Z},$$

but then $\tilde{f}_*: 1 \rightarrow \mathbb{Z}$ cannot be an isomorphism and hence \tilde{f} is not a homeomorphism. Thus $f(x) \in S^1$ and $f|_{S^1}$ is a homeomorphism.



Top / Diff Geo Prelim Exam August 2011

3)



* (a) Q is not a topological manifold because it fails to be locally Euclidean where the corners meet. If Q were to be a topological manifold, it would need to have dimension 2. Therefore, any open neighborhood of the meeting place of the three corners would need to be homeomorphic to \mathbb{R}^2 . However, with the way that the equivalence relation is defined, any open neighborhood of the point where the corners meet would self-intersect which doesn't happen in \mathbb{R}^2 even when embedded in a higher dimensional space.

(b) Let



Then we can apply Seifert - Van Kampen (SVK) to get that

$$\pi_1(X) \cong \mathbb{Z}/3\mathbb{Z}$$

To see this, consider the diagram

$$\begin{array}{ccccc} & & \pi_1(U) & & \\ \swarrow \gamma_1^* & & \searrow \beta_1^* & & \\ \langle \alpha \rangle = \pi_1(u \cap v) & & & & \pi_1(X) \\ \searrow \gamma_2^* & & \swarrow \beta_2^* & & \\ & & \pi_1(V) & & \end{array}$$

Since $\pi_1(U) = \langle \alpha \rangle$ $\pi_1(V) = 1$ $\gamma_1^*(\alpha) = \alpha^3$ $\beta_1^*(\alpha) = 1$
 $\pi_1(u \cap v) = \langle \alpha \rangle$

$$\pi_1(X) \cong \pi_1(U) *_{\pi_1(u \cap v)} \pi_1(V) = \frac{\langle \alpha \rangle * 1}{\langle \alpha^3 = 1 \rangle} \cong \langle \alpha \mid \alpha^3 = 1 \rangle \cong \mathbb{Z}/3\mathbb{Z}$$

Top/Diff Geo Prelim Exam August 2011

4) (a) $\alpha = \cos(z) dz + \sin(z) dy$
 $d\alpha = -\sin(z) dz \wedge dx + \cos(z) dz \wedge dy$

$$\begin{aligned}\alpha \wedge d\alpha &= \cos(z)^2 dx \wedge dz \wedge dy - \sin(z)^2 dy \wedge dz \wedge dx \\ &= \cos(z)^2 + \sin(z)^2 dx \wedge dz \wedge dy \\ &= dx \wedge dz \wedge dy \neq 0 \quad \forall p \in T^3\end{aligned}$$

So α is a contact form.

(b) Suppose that there were two Reeb fields for α , A, B .
Well if $\alpha(A) = \alpha(B)$ then we have that for every point (x, y, z)

$$\cos(z) dx(A) + \sin(z) dy(B) = \cos(z) dx(B) + \sin(z) dy(A).$$

If we then pick the points $(0, 0, 0)$ and $(0, 0, \pi/2)$ we get

$$dx(A) = dx(B) \quad \text{and} \quad dy(A) = dy(B).$$

Similarly, we would have

$$dz(A, B) = -\sin(z) dz \wedge dx(A, B) + \cos(z) dz \wedge dy(A, B)$$

and by using the relation $dx(B) = dx(A)$ derived previously and evaluating at $(0, 0, \pi)$ we have

$$(dz(A) - dz(B)) dx(A) = 0.$$

We can also use $dy(B) = dy(A)$ and $(0, 0, 0)$

$$(dz(A) - dz(B)) dy(A) = 0,$$

and putting this all together, we get that

$$dz(A) = dz(B)$$

so $A \equiv B$.

Top/Diff Geo Prelim Exam August 2011

(cont.) 4) (b) For X to be the Reeb field of α we need $\alpha(X) = 1$ and $d\alpha(X) = 0$, so solving the set of equations that this gives us we have that the Reeb field of α is

$$X = \cos(z) \frac{\partial}{\partial x} + \sin(z) \frac{\partial}{\partial y} + 0 \frac{\partial}{\partial z}.$$

(c) Taking X as above, we can compute that the flow of X is given by

$$\Theta_t(x, y, z) = (x + \cos(z)t, y + \sin(z)t, z)$$

and this is not periodic because if $(x_0, y_0, z_0) \neq (x_1, y_1, z_1)$

$$(x_0 + \cos(z_0)t, y_0 + \sin(z_0)t, z_0) \neq (x_1 + \cos(z_1)t, y_1 + \sin(z_1)t, z_1).$$

since any ^{values} coordinates z_0, z_1 that could result in equality of the first two coordinates will result in inequality of the ~~the~~ third.

Top / Diff Geo Prelim Exam August 2011

- 5) Consider the canonical isomorphism $M_{2,2} \rightarrow \mathbb{R}^4: \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (a, b, c, d)$ and define the map F by
- $$F(w, x, y, z) = (w^2 + x^2 + y^2, z^2, wz - xy).$$

Then we have that

$$DF = \begin{pmatrix} 2w & 2x & 2y & 2z \\ z & -y & -x & w \end{pmatrix}$$

and this matrix has rank < 2 only when all of the following are zero:

$$\begin{aligned} &-2wy - 2xz, -2wx - 2yz, 2w^2 - 2z^2, \\ &-2x^2 + 2y^2, 2xw + 2xz, 2wy + 2xz \end{aligned} \quad \begin{aligned} &\text{(Determinants of all} \\ &\text{2x2 minors)} \end{aligned}$$

This never happens so long as $a \neq 2|b|$ so each point in the set

$$S = \{(a, b) : a \neq 2|b|\}$$

is a regular value of F so $\forall (a, b) \in S$

$$M_{ab} = F^{-1}(\{(a, b)\})$$

is a smooth submanifold of \mathbb{R}^4 by the Regular Level Set Theorem.