

# **Title of the Thesis**

Author's name

Masterarbeit in Physik  
angefertigt im Helmholtz-Institut für Strahlen- und  
Kernphysik

vorgelegt der  
Mathematisch-Naturwissenschaftlichen Fakultät  
der  
Rheinischen Friedrich-Wilhelms-Universität  
Bonn

MMM 2021

DRAFT

I hereby declare that this thesis was formulated by myself and that no sources or tools other than those cited were used.

Bonn, .....  
Date

.....  
Signature

- 1. Gutachter: Prof. Dr. John Smith
- 2. Gutachterin: Prof. Dr. Anne Jones

# Acknowledgements

---

I would like to thank ...

You should probably use `\chapter*` for acknowledgements at the beginning of a thesis and `\chapter` for the end.

DRAFT

DRAFT

# Contents

---

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Theoretical Overview</b>	<b>3</b>
2.1	Excitons . . . . .	3
2.2	Hubbard-Model . . . . .	3
2.3	HMC . . . . .	3
2.4	Correlation Functions . . . . .	3
<b>3</b>	<b>Symmetries</b>	<b>5</b>
3.1	Lattice . . . . .	5
3.2	Symmetry Operators . . . . .	7
3.2.1	Spin-flip transformation . . . . .	7
3.2.2	Charge transformation . . . . .	9
3.2.3	Particle-hole transformation . . . . .	10
3.3	Correlation functions . . . . .	13
3.3.1	One-particle correlation function . . . . .	13
3.3.2	Two-particle correlation function . . . . .	14
<b>4</b>	<b>Spectre of Excitons</b>	<b>31</b>
4.1	Data Generating . . . . .	31
4.2	Fitting Data . . . . .	31
<b>A</b>	<b>Useful information</b>	<b>33</b>
	<b>Bibliography</b>	<b>35</b>
	<b>List of Figures</b>	<b>37</b>
	<b>List of Tables</b>	<b>39</b>

DRAFT

## Introduction

---

The introduction usually gives a few pages of introduction to the whole subject, maybe even starting with the Greeks.

For more information on L<sup>A</sup>T<sub>E</sub>X and the packages that are available see for example the books of Kopka [1] and Goossens et al [2].

A lot of useful information on particle physics can be found in the “Particle Data Book” [3].

I have resisted the temptation to put a lot of definitions into the file `thesis_defs.sty`, as everyone has their own taste as to what scheme they want to use for names. However, a few examples are included to help you get started:

- cross-sections are measured in pb and integrated luminosity in pb<sup>-1</sup>;
- the  $K_S^0$  is an interesting particle;
- the missing transverse momentum,  $p_T^{\text{miss}}$ , is often called missing transverse energy, even though it is calculated using a vector sum.

Note that the examples of units assume that you are using the `siunitx` package.

It also is probably a good idea to include a few well formatted references in the thesis skeleton. More detailed suggestions on what citation types to use can be found in the “Thesis Guide” [4]:

- articles in refereed journals [3, 5];
- a book [6];
- a PhD thesis [7] and a Diplom thesis [8];
- a collection of articles [9];
- a conference note [10];
- a preprint [11] (you can also use `@online` or `@booklet` for such things);
- something that is only available online [4].

At the end of the introduction it is normal to say briefly what comes in the following chapters.

The line at the beginning of this file is used by TeXstudio etc. to specify which is the master L<sup>A</sup>T<sub>E</sub>X file, so that you can compile your thesis directly from this file. If your thesis is called something other than `mythesis`, adjust it as appropriate.





# Theoretical Overview

---

## 2.1 Excitons

## 2.2 Hubbard-Model

## 2.3 HMC

## 2.4 Correlation Functions



## Symmetries

A very important aspect of the analysis are the symmetries. They let us extract useful information from the available data. We can leverage the symmetries we have found, so that we can reduce the numerical uncertainties of our data by averaging, or we can reduce the computational cost by declaring two results are the same by symmetry. The term means any operation one can perform on an operator that leaves it invariant.

In this chapter we will see what is the symmetry of the lattice, which transformations are symmetry operators of the Hubbard model hamiltonian, and how they transform the correlation functions that we are interested in. This will help us better our data and the results we extract from it.

### 3.1 Lattice

The honeycomb lattice is a 2D lattice where the points of the primitive cell lie on the vertices of a hexagon. The lattice plane is generated by two non-orthogonal vectors

$$\vec{a}_1 = \frac{\sqrt{3}}{2} \begin{pmatrix} \sqrt{3} \\ +1 \end{pmatrix} \quad \vec{a}_2 = \frac{\sqrt{3}}{2} \begin{pmatrix} \sqrt{3} \\ -1 \end{pmatrix} \quad (3.1)$$

By repeating them  $L_1$  and  $L_2$  times respectively, we can construct the whole finite lattice. The reciprocal lattice is also hexagonal. The honeycomb lattice can be seen as made of two triangular lattices. Lattices with this property are called a bipartite. In general a bipartite lattice is any one which we can make from two overlapping lattices, where every point of one of the sublattices is only neighbouring points from the other sublattice. On Fig. 3.1 is shown the bipartite of the hexagonal lattice.

In order to investigate the symmetry operators and how they transform the correlation functions, we work with the reciprocal lattice. The primitive cell of the reciprocal is called Brillouin zone (BZ). Some of the points on the lattice which are of special interest have labels that are used in the literature and in this thesis. These are center of the first Brillouin zone called the Gamma point ( $\Gamma$ ), the two points on the vertices of the BZ called the Dirac points ( $K, K'$ ), and the three middle of the edges of the BZ points  $M, M', M''$ . They are all shown on Fig. 3.2 as well as the first neighbours. The graph shows that all symmetries are contained in the first BZ and we do not need to investigate beyond it. Therefore, we consider only the first Brillouin zone in our calculations.




Figure 3.1: Honeycomb lattice is a bipartite lattice because we can separate and label the lattice points with different colors (in this example blue and red) in such a way that every point in blue color neighbours only points with red color.




Figure 3.2: LEFT: All points of special interest shown on the lattice and how they are copied over to other lattice cells. RIGHT: Special points only in the first BZ. It contains all symmetries of the lattice




Figure 3.3: An example of the spin-flip transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the spin-flip conjugation.

The primitive cell of the reciprocal lattice of the honeycomb is a hexagon, meaning it must have all the symmetries of the dihedral group.

## 3.2 Symmetry Operators

In this section we are going to focus only on transformation that do not change the hamiltonian in general or at half filling ( $\mu = 0$ ) i.e. symmetry operations which means they commute with it

$$[H, R] = 0. \quad (3.2)$$

We can show that the expectation value of the operator does not change when we apply the symmetry to it.

$$\langle O \rangle = \frac{1}{Z} \text{tr}[O R^{-1} R e^{-\beta H}] = \frac{1}{Z} \text{tr}[O R^{-1} e^{-\beta H} R] = \frac{1}{Z} \text{tr}[R O R^{-1} e^{-\beta H}] = \langle R O R^{-1} \rangle \quad (3.3)$$

For our chosen model, there are only a few symmetry operators that satisfy this condition. We are going to see how they can be used to our advantage to increase our statistics.

### 3.2.1 Spin-flip transformation

On a bipartite lattice, we have spin-flip operation that exchanges the particles and holes but leaves the charge untouched. On Figure 3.3 there is shown an example of how the spin-flip operation transforms each operator when applied. Unfortunately, the example can show only how the spin and the charge transform but not the hamiltonian terms and amplitudes.

When applied onto the creation/annihilation operators of the particles and the holes, this spin transformation gives us

$$\begin{aligned}
 IpI^{-1} &= \Sigma h^\dagger = h^\dagger \Sigma & IhI^{-1} &= \Sigma p^\dagger = p^\dagger \Sigma \\
 Ip^\dagger I^{-1} &= \Sigma h = h \Sigma & Ih^\dagger I^{-1} &= \Sigma p = p \Sigma
 \end{aligned}$$

where  $\Sigma$  is a matrix which shows us the behaviour of the bipartite lattice

$$\Sigma_{xy} = (-1)^x \delta_{xy} \quad (3.4)$$

We can check if this symmetry commutes with the Hamiltonian by first showing that it leaves the charge invariant,

$$IqI^{-1} = Ip^\dagger I^{-1} IpI^{-1} - Ih^\dagger I^{-1} IhI^{-1} = h \Sigma \Sigma h^\dagger - p \Sigma \Sigma p^\dagger = (1 - h^\dagger h) - (1 - p^\dagger p) = p^\dagger p - h^\dagger h = q \quad (3.5)$$

It can also be shown that the third component of the spin gets flipped after we apply this operator

$$Is^3 I^{-1} = -s^3, \quad (3.6)$$

where

$$s_x^3 = \frac{1}{2} \left( \delta_{xx} - p_x^\dagger p_x - h_x^\dagger h_x \right) \quad (3.7)$$

Finally, it is important to see how the hopping term transforms under this symmetry operation

$$\begin{aligned}
 I \left( - \sum_{xy} p_x^\dagger K_{xy} p_y + \sigma_k h_x^\dagger K_{xy} h_y \right) I^{-1} &= - \sum_{xy} Ip_x^\dagger I^{-1} IK_{xy} I^{-1} Ip_y I^{-1} + \sigma_k Ih_x^\dagger I^{-1} IK_{xy} I^{-1} Ih_y I^{-1} \\
 &= - \sum_{xy} h_x (\Sigma IKI^{-1} \Sigma)_{xy} h_y^\dagger + \sigma_k p_x (\Sigma IKI^{-1} \Sigma)_{xy} p_y^\dagger \\
 &= - \sum_{xy} -(\delta_{xy} (IKI^{-1})_{xy} - h_y^\dagger (IKI^{-1})_{xy} h_x) - \\
 &\quad - \sigma_k (\delta_{xy} (IKI^{-1})_{xy} - p_y^\dagger (IKI^{-1})_{xy} p_x) \\
 &= \sigma_k \left[ - \sum_{xy} p_x^\dagger (IKI^{-1})_{xy}^\top p_y + \sigma_k h_x^\dagger (IKI^{-1})_{xy}^\top h_y \right]
 \end{aligned}$$

where  $\Sigma K \Sigma = -K$ ,  $\text{tr}[K] = 0$ , and  $\sigma_k^2 = 1$ . In general to return to our initial hopping term, we need  $I$  to be anti-unitary where  $IKI^{-1} = K^*$  and then we could replace  $K^\dagger \rightarrow K$ . Since we are working on a bipartite lattice, we can choose  $\sigma_k = 1$  which even for  $\mu \neq 0$  gives us the commutator  $[H - \vec{\mu} \cdot \vec{q}, I] = 0$ . This is true because as shown in (3.5) the charge is even in relation to the spin-flip symmetry operation.




Figure 3.4: An example of the charge transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the charge-flip conjugation.

### 3.2.2 Charge transformation

Other symmetry we can consider is the charge symmetry operation. It exchanges the charges but leaves the spin fixed. When applied onto the creation and annihilation operators, it transforms them as

$$\begin{aligned} CpC^{-1} &= h & ChC^{-1} &= p \\ Cp^{\dagger}C^{-1} &= h^{\dagger} & Ch^{\dagger}C^{-1} &= p^{\dagger} \end{aligned}$$

On Figure 3.4 is shown how the charge and spin of all creation/annihilation operators transform under this symmetry operation. If we conjugate with the charge operator, we get

$$CqC^{-1} = Cp^{\dagger}C^{-1}CpC^{-1} - Ch^{\dagger}C^{-1}ChC^{-1} = h^{\dagger}h - p^{\dagger}p = -q \quad (3.8)$$

and when conjugated with the spin operator

$$Cs^3C^{-1} = s^3, \quad (3.9)$$

so the spin is fixed for this operation. Again, we want to calculate the conjugation of the tight-binding term in the hamiltonian with the charge-flip operator

$$\begin{aligned}
 C \left( - \sum_{xy} p_x^\dagger K_{xy} p_y + \sigma_k h_x^\dagger K_{xy} h_y \right) C^{-1} &= - \sum_{xy} C p_x^\dagger C^{-1} C K_{xy} C^{-1} C p_y C^{-1} + \sigma_k C h_x^\dagger C^{-1} C K_{xy} C^{-1} C h_y C^{-1} \\
 &= - \sum_{xy} h_x^\dagger (C K C^{-1})_{xy} h_y + \sigma_k p_x^\dagger (C K C^{-1})_{xy} p_y \\
 &= - \sum_{xy} \sigma_k p_x^\dagger (C K C^{-1})_{xy} p_y + h_x^\dagger (C K C^{-1})_{xy} h_y
 \end{aligned}$$

where if we choose  $C K C^{-1} = K$ , we get to the initial term. Therefore, we are shown that on a bipartite lattice where we can choose  $\sigma_k = +1$ , we have a charge conjugation symmetry i.e. the it commutes with the tight-binding hamiltonian  $[H, C] = 0$ . This is because as shown in (3.8) the charge operator is flipped but the the interaction is even under this transformation because it is of the form  $qVq$ .

### 3.2.3 Particle-hole transformation

The particle-hole symmetry operator is a combination of two symmetry operators that do not transform the hamiltonian nicely but their combination does. It preserves the energy of the system but flips the charge and the spin. This means that we should only consider the tightbinding hamiltonian i.e. the hamiltonian without the chemical potential term.

In order to convince ourselves of the neat transformation and also the commutation of the hamiltonian with the symmetry operator, we should first see how the constituent operators transform the Hubbard model hamiltonian.

First, we look at the staggering operator  $X$ . It transforms the creation/anihilation operators like

$$\begin{aligned}
 X p^\dagger X^{-1} &= \Sigma p^\dagger = p^\dagger \Sigma & X h^\dagger X^{-1} &= \Sigma h^\dagger = h^\dagger \Sigma \\
 X p X^{-1} &= \Sigma p = p \Sigma & X h X^{-1} &= \Sigma h = h \Sigma
 \end{aligned}$$

It leaves the charge, spin, and number operators invariant but it flips the sign of the hopping term

$$\begin{aligned}
 X \left( - \sum_{xy} p_x^\dagger K_{xy} p_y + \sigma_k h_x^\dagger K_{xy} h_y \right) X^{-1} &= - \sum_{xy} X p_x^\dagger X^{-1} X K_{xy} X^{-1} X p_y X^{-1} + \sigma_k X h_x^\dagger X^{-1} X K_{xy} X^{-1} X h_y X^{-1} \\
 &= - \sum_{xy} p_x^\dagger (\Sigma X K X^{-1} \Sigma)_{xy} p_y + \sigma_k h_x^\dagger (\Sigma X K X^{-1} \Sigma)_{xy} h_y \\
 &= - \sum_{xy} p_x^\dagger (\Sigma K \Sigma)_{xy} p_y + \sigma_k h_x^\dagger (\Sigma K \Sigma)_{xy} h_y \\
 &= - \left[ - \sum_{xy} p_x^\dagger K_{xy} p_y + \sigma_k h_x^\dagger K_{xy} h_y \right]
 \end{aligned}$$

The other operator that we consider is the one that "flips" the dagger of the operators. It transforms the particle/hole operators as



$$\begin{aligned} FpF^{-1} &= p^\dagger & FhF^{-1} &= h^\dagger \\ Fp^\dagger F^{-1} &= p & Fh^\dagger F^{-1} &= h \end{aligned}$$

When applied onto the charge operator, it flips the the sign  $q \rightarrow -q$  which is shown below

$$FqF^{-1} = Fp^\dagger F^{-1} FpF^{-1} - Fh^\dagger F^{-1} FhF^{-1} = pp^\dagger - hh^\dagger = (1 - p^\dagger p) - (1 - h^\dagger h) = -q \quad (3.10)$$

This in turn means that this symmetry operation flips the chemical potential and more importantly the hopping term

$$\begin{aligned} F \left( - \sum_{xy} p_x^\dagger K_{xy} p_y + \sigma_k h_x^\dagger K_{xy} h_y \right) F^{-1} &= - \sum_{xy} Fp_x^\dagger F^{-1} FK_{xy} F^{-1} Fp_y F^{-1} + \sigma_k Fh_x^\dagger F^{-1} FK_{xy} F^{-1} Fh_y F^{-1} \\ &= - \sum_{xy} p_x (FKF^{-1})_{xy} p_y^\dagger + \sigma_k h_x (FKF^{-1})_{xy} h_y^\dagger \\ &= - \sum_{xy} (\delta_{xy} (FKF^{-1})_{xy} - p_y^\dagger (FKF^{-1})_{xy} p_x) + \sigma_k (\delta_{xy} (FKF^{-1})_{xy} - h_y^\dagger (FKF^{-1})_{xy} h_x) \\ &= - \left[ - \sum_{xy} p_y^\dagger (FKF^{-1})_{xy} p_x + \sigma_k h_y^\dagger (FKF^{-1})_{xy} h_x \right] \\ &= - \left[ - \sum_{xy} p_x^\dagger (FKF^{-1})_{xy}^\top p_y + \sigma_k h_x^\dagger (FKF^{-1})_{xy}^\top h_y \right] \end{aligned}$$

where  $\text{tr}[K] = 0$ , if we want to return to the original hopping term then we have to choose  $FKF^{-1} = K^*$  and then we can substitute  $K^\dagger \rightarrow K$ . We see on Figure 3.5 an exmaple of the dagger-flip operator switching the signs of the charge and the spin.

After we have shown how the different term of the hamiltonian transform under both symmetries separately, we can answer how the composition of the two transforms each term of the Hubbard model's hamiltonian. Starting from the hopping term, we know that each operation flips the sign of the kinetic term, meaning it is even under the composed operator. Next, we know that the charge operator is invariant under the staggering operator and it flips sign when conjugated with the dagger-flip operator. This means that the interaction term  $qVq$  must preserve its sign with the particle-hole transformation because the interaction is bilinear. Finally, the chemical potential unfortunately is linear  $\vec{\mu} \cdot \vec{q}$ , so the sign of this term is flipped. After all these considerations, we could conclude that at half-filling ( $\mu = 0$ ), the tightbinding hamiltonian commutes with the particle-hole transformation

$$[H, XF] = 0 \quad (3.11)$$

and we can use the this symmetry operation to our advantage. On Figure 3.6 it is shown again an example of the transformation of spin and charge. On this graph we can only visualize the spin and the charge of the operators but not the transformed terms of the hamiltonian, resulting in the same looking example as if only we consider  $F$  in spite of the kinetic terms having opposite signs.




Figure 3.5: An example of the dagger-flip transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the dagger-flip conjugation.




Figure 3.6: An example of the particle-hole transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the particle-hole conjugation.

It is worth noting that after working out how these symmetry operations transform the hamiltonian, we could write out  $I = XFC$  but we should still examine them all separately because  $I$  commutes with the whole hamiltonian and the other two only with the tight-binding one.

### 3.3 Correlation functions

After we have found out what the symmetries of the lattice are and how they transform the relevant operators, we can take them into consideration and apply them onto the one- and two-body correlation functions. We use them in order to average data with the same total momentum. This will increase our statistics and better our precision when performing the fitting of the final data.

#### 3.3.1 One-particle correlation function

First, we start with the one-body correlators which will give us a valuable insight on how to work with the two-body correlation functions. The full derivation is provided in EVAN'S PAPER, so only the relevant information for this thesis will be shown here.

We start with the  $\langle pp^\dagger \rangle$  correlation function which we can write as  $C_Q = +1, S = 1, S^3 = -1, \vec{k}$ . It can be represented as a  $2 \times 2$  matrix with two one-body correlators due to the internal band labels

$$C_{Q=+1, S=1, S^3=-1, \vec{k}} = \begin{bmatrix} p_+ p_+^\dagger & p_+ p_-^\dagger \\ p_- p_+^\dagger & p_- p_-^\dagger \end{bmatrix}_{\vec{k}} (\tau) \quad (3.12)$$

This matrix of correlators can be calculated and then diagonalized to

$$C_{Q=+1, S=1, S^3=-1, \vec{k}} = \frac{1}{2} (C_{p_+ p_+^\dagger} + C_{p_- p_-^\dagger}) \mathbb{1} + \frac{1}{2} \left( \sqrt{(C_{p_+ p_+^\dagger} - C_{p_- p_-^\dagger})^2 + 4C_{p_+ p_-^\dagger} C_{p_- p_+^\dagger}} \right) \sigma_3 \quad (3.13)$$

In the tight-binding case of the hamiltonian, the off-diagonal components vanish and the diagonal are equal. However, in general one must be careful because this is discretization dependent. In order to avoid this problem, we must always compute the whole matrix and then diagonalize. The whole argument can also be done for  $\langle hh^\dagger \rangle$ .

To increase the precision of our data, we apply the symmetry operators that we calculated in Section 3.2. Also, we use the cyclic property of the trace and the final results can be seen in Table 3.1. A detailed derivation of these results as mentioned previously can be found in EVAN'S PAPER. It is important to see that we can increase the precision eightfold not sixteenfold which is the number of cases in the table. This is because some of the entries are duplicates and they can be found by examining the quantum numbers in the second column. The averaging put into an equation looks like this

$$\tilde{C}_{QS^3\vec{k}}(\tau) = \frac{1}{4} \left( C_{QS^3\vec{k}}(\tau) + C_{\overline{QS^3\vec{k}}}(\beta - \tau)^\top + \sigma_1 \cdot C_{QS^3\vec{k}}(\tau) \cdot \sigma_1 + \sigma_1 \cdot C_{\overline{QS^3\vec{k}}}(\beta - \tau)^\top \cdot \sigma_1 \right) \quad (3.14)$$

after which one must diagonalize.

		$Q$	$S$	$S^3$	$\vec{k}$	$\tau$
Start with	$QSk t$	+1	$\frac{1}{2}$	$-\frac{1}{2}$	1	$\tau$
trace cycl	$\overline{QSk t}$	-1	$\frac{1}{2}$	$+\frac{1}{2}$	$\top$	$\beta - \tau$
I	$QSk t$	+1	$\frac{1}{2}$	$+\frac{1}{2}$	$\sigma_1 \cdot \sigma_1$	$\tau$
		-1	$\frac{1}{2}$	$-\frac{1}{2}$	$\sigma_1 \top \sigma_1$	$\beta - \tau$
XF	$\overline{QSk t}$	-1	$\frac{1}{2}$	$+\frac{1}{2}$	$\sigma_1 \cdot \sigma_1$	$\tau$
		+1	$\frac{1}{2}$	$-\frac{1}{2}$	$\sigma_1 \cdot \sigma_1$	$\beta - \tau$
		-1	$\frac{1}{2}$	$-\frac{1}{2}$	1	$\tau$
		+1	$\frac{1}{2}$	$+\frac{1}{2}$	$\top$	$\beta - \tau$
C	$\overline{QSk t}$	-1	$\frac{1}{2}$	$-\frac{1}{2}$	1	$\tau$
		+1	$\frac{1}{2}$	$+\frac{1}{2}$	$\top$	$\beta - \tau$
		-1	$\frac{1}{2}$	$+\frac{1}{2}$	$\sigma_1 \cdot \sigma_1$	$\tau$
		+1	$\frac{1}{2}$	$-\frac{1}{2}$	$\sigma_1 \top \sigma_1$	$\beta - \tau$
		+1	$\frac{1}{2}$	$+\frac{1}{2}$	$\sigma_1 \cdot \sigma_1$	$\tau$
		-1	$\frac{1}{2}$	$-\frac{1}{2}$	$\sigma_1 \top \sigma_1$	$\beta - \tau$
		+1	$\frac{1}{2}$	$-\frac{1}{2}$	1	$\tau$
		-1	$\frac{1}{2}$	$+\frac{1}{2}$	$\top$	$\beta - \tau$

Table 3.1: All the correlators that can be directly averaged, assuming they all take the  $2 \times 2$  matrix form. The correlators above the double line may be averaged even at finite  $\mu$  and everything below may be averaged when  $\mu = 0$ . The combination of charge and spin indicate whether  $p^\dagger$ ,  $p$ ,  $h^\dagger$ , or  $h$  is right-most. The  $\vec{k}$  column indicates what must be done in the internal  $2 \times 2$  momentum space before averaging;  $\top$  indicates transpose,  $\sigma_1 \cdot \sigma_1$  indicates conjugation by  $\sigma_1$ , 1 indicates that no operation is needed. The  $\tau$  column indicates whether the given correlator needs to be time-reversed.

### 3.3.2 Two-particle correlation function

In this thesis we are working with only two types of two-body correlators. There are correlators with particles and holes, and only particles. We start with the former correlator  $\langle p h h^\dagger p^\dagger \rangle$ . In matrix form with all labels it looks like this

$$C_{Q=0, S=1, S^3=-1; k, q, r, s} = \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k, q, r, s} \quad (\tau) \quad (3.15)$$

Now, we apply different symmetry operator to the two particle correlation function. The derivations of the conjugation with the symmetry operators are always done in two steps. First, we start with the thermal trace and then switch to matrix form, so that it is easier to see the transformations needed in order to get another two particle correlator.

### Trace cyclicity

We begin with the trace cyclicity because it will give us how the correlation function transformations when timereversed.

$$C_{Q=0,S=1,S^3=-1;k,q,r,s} = \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \frac{1}{Z} \text{tr} \left[ h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} p_k h_q e^{-H\tau} \right] \quad (3.16)$$

Now we apply time reverse

$$\beta \rightarrow \beta - \tau$$

and use the anti-commutation relations of the operators to get

$$\frac{1}{Z} \text{tr} \left[ h_r^\dagger p_s^\dagger e^{-H(\beta-(\beta-\tau))} p_k h_q e^{-H(\beta-\tau)} \right] = \frac{1}{Z} \text{tr} \left[ p_s^\dagger h_r^\dagger e^{-H\tau} h_q p_k e^{-H(\beta-\tau)} \right] \quad (3.17)$$

We switch to matrix form to see observe better how all labels transform

$$C_{Q=0,S=1,S^3=-1;k,q,r,s} = \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k,q,r,s} \quad (\tau) \quad (3.18)$$

$$= \begin{bmatrix} h_+^\dagger p_+^\dagger p_+ h_+ & h_+^\dagger p_-^\dagger p_+ h_+ & h_-^\dagger p_+^\dagger p_+ h_+ & h_-^\dagger p_-^\dagger p_+ h_+ \\ h_+^\dagger p_+^\dagger p_+ h_- & h_+^\dagger p_-^\dagger p_+ h_- & h_-^\dagger p_+^\dagger p_+ h_- & h_-^\dagger p_-^\dagger p_+ h_- \\ h_+^\dagger p_+^\dagger p_- h_+ & h_+^\dagger p_-^\dagger p_- h_+ & h_-^\dagger p_+^\dagger p_- h_+ & h_-^\dagger p_-^\dagger p_- h_+ \\ h_+^\dagger p_+^\dagger p_- h_- & h_+^\dagger p_-^\dagger p_- h_- & h_-^\dagger p_+^\dagger p_- h_- & h_-^\dagger p_-^\dagger p_- h_- \end{bmatrix}_{r,s,k,q} \quad (\beta - \tau)$$

$$= \begin{bmatrix} p_+^\dagger h_+^\dagger h_+ p_+ & p_-^\dagger h_+^\dagger h_+ p_+ & p_+^\dagger h_-^\dagger h_+ p_+ & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_- p_+ \\ p_+^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_-^\dagger h_+ p_- & p_-^\dagger h_-^\dagger h_+ p_- \\ p_+^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_- \end{bmatrix}_{s,r,q,k} \quad (\beta - \tau) \quad (3.19)$$

$$= C_{Q=0,S=1,S^3=+1;s,r,q,k}$$

We notice that order reverse is actually a composed of two transformations – particle-hole momentum switch and time reverse. After we stare at the matrix for an hour, we can figure out what the transformation between the correlation functions is

$$C_{Q=0,S=1,S^3=-1;k,q,r,s}(\tau) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} C_{Q=0,S=1,S^3=+1;s,r,q,k}^\top(\beta - \tau) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (3.20)$$

We repeat the steps again for each symmetry operator. First we start with the thermal trace and then switch to matrix for easy transformations. It is shown on the right side of the column sign if the operator transforms well the full hamiltonian or only at half-filling.

$$\underline{I : H - \vec{\mu} \cdot \vec{q}}$$

$$\begin{aligned}
 C_{Q=0, S=1, S^3=-1; k, q, r, s} &= \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ I^{-1} I p_k I^{-1} I h_q I^{-1} I e^{-H\tau} I^{-1} I h_r^\dagger I^{-1} I p_s^\dagger I^{-1} I e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ I p_k I^{-1} I h_q I^{-1} I e^{-H\tau} I^{-1} I h_r^\dagger I^{-1} I p_s^\dagger I^{-1} I e^{-H(\beta-\tau)} I^{-1} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma h)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} (\Sigma p)_r (\Sigma h)_s e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_q^\dagger (\Sigma h)_k^\dagger e^{-H\tau} (\Sigma h)_s (\Sigma p)_r e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_r (\Sigma h)_s e^{-H(\beta-\tau)} (\Sigma h)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} \right]
 \end{aligned} \tag{3.21}$$

$$\begin{aligned}
 C_{Q=0, S=1, S^3=-1; k, q, r, s} &= I^{-1} \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k, q, r, s} (\tau) I \\
 &= \begin{bmatrix} (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_- \\ (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_- \\ (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_- \\ (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_- \end{bmatrix} \\
 &= \begin{bmatrix} h_-^\dagger p_-^\dagger p_- h_- & h_-^\dagger p_-^\dagger p_- h_+ & h_-^\dagger p_-^\dagger p_+ h_- & h_-^\dagger p_-^\dagger p_+ h_+ \\ h_-^\dagger p_+^\dagger p_- h_- & h_-^\dagger p_+^\dagger p_- h_+ & h_-^\dagger p_+^\dagger p_+ h_- & h_-^\dagger p_+^\dagger p_+ h_+ \\ h_+^\dagger p_-^\dagger p_- h_- & h_+^\dagger p_-^\dagger p_- h_+ & h_+^\dagger p_-^\dagger p_+ h_- & h_+^\dagger p_-^\dagger p_+ h_+ \\ h_+^\dagger p_+^\dagger p_- h_- & h_+^\dagger p_+^\dagger p_- h_+ & h_+^\dagger p_+^\dagger p_+ h_- & h_+^\dagger p_+^\dagger p_+ h_+ \end{bmatrix}_{k, q, r, s} (\tau)
 \end{aligned} \tag{3.22}$$

$$= \begin{bmatrix} p_- h_- h_-^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger \\ p_- h_- h_+^\dagger p_-^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_+^\dagger \end{bmatrix}_{r, s, k, q} (\beta - \tau) \tag{3.23}$$

$$= \begin{bmatrix} p_-^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_+ p_- & p_-^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_-^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_+ p_- & p_+^\dagger h_-^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_+ p_+ \\ p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_+^\dagger h_+ p_+ \end{bmatrix}_{q, k, s, r} (\tau) \tag{3.24}$$

Depending on what transformation we use to get back to  $\langle phh^\dagger p^\dagger \rangle$ , we get two transformations which are time reverse to one another. We notice the same behaviour for all operators.

$$C_{Q=0,S=1,S^3=-1;k,q,r,s}(\tau) = \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0,S=1,S^3=-1;r,s,k,q}^\top(\beta-\tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} \quad (3.25)$$

$$C_{Q=0,S=1,S^3=-1;k,q,r,s}(\tau) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0,S=1,S^3=+1;q,k,s,r}(\tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (3.26)$$

### C : H

$$\begin{aligned} C_{Q=0,S=1,S^3=-1;k,q,r,s} &= \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\ &= \frac{1}{Z} \text{tr} \left[ C^{-1} C p_k C^{-1} C h_q C^{-1} C e^{-H\tau} C^{-1} C h_r^\dagger C^{-1} C p_s^\dagger C^{-1} C e^{-H(\beta-\tau)} \right] \\ &= \frac{1}{Z} \text{tr} \left[ h_k p_q e^{-H\tau} p_r^\dagger h_s^\dagger e^{-H(\beta-\tau)} \right] \\ &= \frac{1}{Z} \text{tr} \left[ p_q h_k e^{-H\tau} h_s^\dagger p_r^\dagger e^{-H(\beta-\tau)} \right] \end{aligned} \quad (3.27)$$

$$C_{Q=0,S=1,S^3=-1;k,q,r,s} = C^{-1} \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k,q,r,s} (\tau) C \quad (3.28)$$

$$\begin{aligned}
 &= \begin{bmatrix} h_+ p_+ p_+^\dagger h_+^\dagger & h_+ p_+ p_+^\dagger h_-^\dagger & h_+ p_+ p_-^\dagger h_+^\dagger & h_+ p_+ p_-^\dagger h_-^\dagger \\ h_+ p_- p_+^\dagger h_+^\dagger & h_+ p_- p_+^\dagger h_-^\dagger & h_+ p_- p_-^\dagger h_+^\dagger & h_+ p_- p_-^\dagger h_-^\dagger \\ h_- p_+ p_+^\dagger h_+^\dagger & h_- p_+ p_+^\dagger h_-^\dagger & h_- p_+ p_-^\dagger h_+^\dagger & h_- p_+ p_-^\dagger h_-^\dagger \\ h_- p_- p_+^\dagger h_+^\dagger & h_- p_- p_+^\dagger h_-^\dagger & h_- p_- p_-^\dagger h_+^\dagger & h_- p_- p_-^\dagger h_-^\dagger \end{bmatrix}_{k,q,r,s} (\tau) \\
 &= \begin{bmatrix} p_+^\dagger h_+^\dagger h_+ p_+ & p_+^\dagger h_-^\dagger h_+ p_+ & p_-^\dagger h_+^\dagger h_+ p_+ & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_-^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_-^\dagger h_+ p_- \\ p_+^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_- p_+ \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_- \end{bmatrix}_{r,s,k,q} (\beta - \tau)
 \end{aligned} \tag{3.29}$$

$$\begin{aligned}
 &= \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{q,k,s,r} (\tau)
 \end{aligned} \tag{3.30}$$

$$C_{Q=0,S=1,S^3=-1;k,q,r,s}(\tau) = C_{Q=0,S=1,S^3=+1;r,s,k,q}^\top(\beta - \tau) \tag{3.31}$$

$$C_{Q=0,S=1,S^3=-1;k,q,r,s}(\tau) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} C_{Q=0,S=1,S^3=-1;q,k,s,r}(\tau) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \tag{3.32}$$

### $XF : H$

$$\begin{aligned}
 &C_{Q=0,S=1,S^3=-1;k,q,r,s} \\
 &= \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (XF)^{-1} (XF) p_k (XF)^{-1} (XF) h_q (XF)^{-1} (XF) e^{-H\tau} (XF)^{-1} (XF) h_r^\dagger (XF)^{-1} (XF) p_s^\dagger (XF)^{-1} (XF) e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_k^\dagger (\Sigma h)_q^\dagger e^{-H\tau} (\Sigma h)_r (\Sigma p)_s e^{-H(\beta-\tau)} \right]
 \end{aligned} \tag{3.33}$$



$$\begin{aligned}
 & C_{Q=0, S=1, S^3=-1; k, q, r, s} \\
 &= (XF)^{-1} \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k, q, r, s} (\tau) (XF) \\
 &= \begin{bmatrix} (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_- \\ (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_- \\ (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_- \\ (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_- \end{bmatrix} \\
 &= \begin{bmatrix} p_-^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_+ p_- & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_+ p_+ \\ p_+^\dagger h_-^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_+ p_- & p_+^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_+^\dagger h_+ p_+ \end{bmatrix}_{k, q, r, s} (\tau) \tag{3.34}
 \end{aligned}$$

$$= \begin{bmatrix} p_- h_- h_-^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_-^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger \\ p_- h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_+^\dagger \end{bmatrix}_{s, r, k, q} (\beta - \tau) \tag{3.35}$$

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=-1; k, q, r, s}(\tau) \\
 &= \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0, S=1, S^3=+1; k, q, r, s}(\tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} \tag{3.36}
 \end{aligned}$$

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=-1; k, q, r, s}(\tau) \\
 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0, S=1, S^3=-1; s, r, q, k}^\top (\beta - \tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \tag{3.37}
 \end{aligned}$$

The table below shows the summary of all symmetries that we calculated.

For more compact form, we write

$$\sigma_5 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \tag{3.38}$$

The other correlation function that we are interested in is with two particles at the source and two at the sink. Again, we start with correlator  $\langle p p p^\dagger p^\dagger \rangle$ . In trace and matrix forms it look like this.

	$Q$	$S$	$S^3$	trafo	$kqrs$	$\tau$
Start with	0	1	-1	1	1234	$\tau$
trace cycl	0	1	+1	$\sigma_5 \top \sigma_5$	4321	$\beta - \tau$
I	0	1	-1	$\sigma_1 \top \sigma_1$	3412	$\beta - \tau$
	0	1	+1	$\sigma_5 \sigma_1 \cdot \sigma_1 \sigma_5$	2143	$\tau$
C	0	1	+1	$\top$	3412	$\beta - \tau$
	0	1	-1	$\sigma_5 \cdot \sigma_5$	2143	$\tau$
XF	0	1	+1	$\sigma_1 \cdot \sigma_1$	1234	$\tau$
	0	1	-1	$\sigma_5 \sigma_1 \top \sigma_1 \sigma_5$	4321	$\beta - \tau$

Table 3.2: All the correlators that can be directly averaged, if they all take a  $4 \times 4$  matrix form. The correlators above the double line may be averaged even at finite  $\mu$  and everything below may be averaged when  $\mu = 0$ . The combination of charge and spin indicate whether  $h^\dagger p^\dagger$  or  $hp$  is at the source (right-most). The *trafo* column indicates what must be done in the internal  $4 \times 4$  momentum space before averaging;  $\top$  indicates transpose,  $\sigma_1 \cdot \sigma_1$  indicates conjugation by  $\sigma_1$ ,  $\sigma_5 \cdot \sigma_5$  indicates conjugation by  $\sigma_5$ , and 1 indicates that no operation is needed. The next column  $kqrs$  indicates the position of the momenta at the sink and source relative to the initial correlator is. The  $\tau$  column indicates whether the given correlator needs to be time-reversed.

$$C_{Q=2, S=1, S^3=-1; k, q, r, s} = \frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \quad (3.39)$$

$$C_{Q=2, S=1, S^3=-1; k, q, r, s} = \left[ \begin{array}{cccc} p_+ p_+ p_+^\dagger p_+^\dagger & p_+ p_+ p_+^\dagger p_-^\dagger & p_+ p_+ p_-^\dagger p_+^\dagger & p_+ p_+ p_-^\dagger p_-^\dagger \\ p_+ p_- p_+^\dagger p_+^\dagger & p_+ p_- p_+^\dagger p_-^\dagger & p_+ p_- p_-^\dagger p_+^\dagger & p_+ p_- p_-^\dagger p_-^\dagger \\ p_- p_+ p_+^\dagger p_+^\dagger & p_- p_+ p_+^\dagger p_-^\dagger & p_- p_+ p_-^\dagger p_+^\dagger & p_- p_+ p_-^\dagger p_-^\dagger \\ p_- p_- p_+^\dagger p_+^\dagger & p_- p_- p_+^\dagger p_-^\dagger & p_- p_- p_-^\dagger p_+^\dagger & p_- p_- p_-^\dagger p_-^\dagger \end{array} \right]_{k, q, r, s} \quad (\tau) \quad (3.40)$$

First, we implement the anti-commutation relations which will give us the identities of this correlation function. Again, we use derive the the results through traces and then continue with the matrix form which will give us a better insight on the transformation of all labels that are included.

### Anti-commutation

$$\begin{aligned} C_{Q=2, S=1, S^3=-1; k, q, r, s} &= \frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \\ &= \frac{1}{Z} \text{tr} \left[ p_q p_k e^{-H\tau} p_s^\dagger p_r^\dagger e^{-H(\beta-\tau)} \right] \\ &= -\frac{1}{Z} \text{tr} \left[ p_q p_k e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\ &= -\frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_s^\dagger p_r^\dagger e^{-H(\beta-\tau)} \right] \end{aligned} \quad (3.41)$$

$$\begin{aligned}
 C_{Q=2,S=1,S^3=-1;k,q,r,s} &= \begin{bmatrix} p_+p_+p_+p_+^\dagger & p_+p_+p_+p_-^\dagger & p_+p_+p_-p_+^\dagger & p_+p_+p_-p_-^\dagger \\ p_+p_-p_+p_+^\dagger & p_+p_-p_+p_-^\dagger & p_+p_-p_-p_+^\dagger & p_+p_-p_-p_-^\dagger \\ p_-p_+p_+p_+^\dagger & p_-p_+p_+p_-^\dagger & p_-p_+p_-p_+^\dagger & p_-p_+p_-p_-^\dagger \\ p_-p_-p_+p_+^\dagger & p_-p_-p_+p_-^\dagger & p_-p_-p_-p_+^\dagger & p_-p_-p_-p_-^\dagger \end{bmatrix}_{k,q,r,s} \quad (\tau) \\
 &= \begin{bmatrix} p_+p_+p_+p_+^\dagger & p_+p_+p_-p_+^\dagger & p_+p_+p_-p_-^\dagger & p_+p_+p_-p_-^\dagger \\ p_-p_+p_+p_+^\dagger & p_-p_+p_-p_+^\dagger & p_-p_+p_-p_-^\dagger & p_-p_+p_-p_-^\dagger \\ p_+p_-p_+p_+^\dagger & p_+p_-p_-p_+^\dagger & p_+p_-p_-p_-^\dagger & p_+p_-p_-p_-^\dagger \\ p_-p_-p_+p_+^\dagger & p_-p_-p_-p_+^\dagger & p_-p_-p_-p_-^\dagger & p_-p_-p_-p_-^\dagger \end{bmatrix}_{q,k,s,r} \quad (\tau) \\
 &= - \begin{bmatrix} p_+p_+p_+p_+^\dagger & p_+p_+p_+p_-^\dagger & p_+p_+p_-p_+^\dagger & p_+p_+p_-p_-^\dagger \\ p_-p_+p_+p_+^\dagger & p_-p_+p_+p_-^\dagger & p_-p_+p_-p_+^\dagger & p_-p_+p_-p_-^\dagger \\ p_+p_-p_+p_+^\dagger & p_+p_-p_-p_+^\dagger & p_+p_-p_-p_-^\dagger & p_+p_-p_-p_-^\dagger \\ p_-p_-p_+p_+^\dagger & p_-p_-p_+p_-^\dagger & p_-p_-p_-p_+^\dagger & p_-p_-p_-p_-^\dagger \end{bmatrix}_{q,k,r,s} \quad (\tau) \\
 &= - \begin{bmatrix} p_+p_+p_+p_+^\dagger & p_+p_+p_-p_+^\dagger & p_+p_+p_-p_-^\dagger & p_+p_+p_-p_-^\dagger \\ p_+p_-p_+p_+^\dagger & p_+p_-p_-p_+^\dagger & p_+p_-p_-p_-^\dagger & p_+p_-p_-p_-^\dagger \\ p_-p_+p_+p_+^\dagger & p_-p_+p_-p_+^\dagger & p_-p_+p_-p_-^\dagger & p_-p_+p_-p_-^\dagger \\ p_-p_-p_+p_+^\dagger & p_-p_-p_-p_+^\dagger & p_-p_-p_-p_-^\dagger & p_-p_-p_-p_-^\dagger \end{bmatrix}_{k,q,s,r} \quad (\tau)
 \end{aligned} \tag{3.42}$$

$$\begin{aligned}
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot C_{Q=2,S=1,S^3=-1;q,k,s,r}(\tau) \cdot \sigma_5 \\
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot C_{Q=2,S=1,S^3=-1;q,k,s,r}(\tau) \\
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -C_{Q=2,S=1,S^3=-1;q,k,s,r}(\tau) \cdot \sigma_5
 \end{aligned} \tag{3.43}$$

These identities show us that we can, average our data four fold only because we are using this kind of correlator. Using the cyclicity of the trace, we find how the correlation function transforms when time is reversed. This will further increase our statistics two times.

### Trace ciclicity

$$\begin{aligned}
 C_{Q=2,S=1,S^3=-1;k,q,r,s} &= \frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \\
 &= \frac{1}{Z} \text{tr} \left[ p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} p_k p_q e^{-H\tau} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ p_s^\dagger p_r^\dagger e^{-H(\beta-\tau)} p_q p_k e^{-H\tau} \right] \\
 &= -\frac{1}{Z} \text{tr} \left[ p_s^\dagger p_r^\dagger e^{-H(\beta-\tau)} p_k p_q e^{-H\tau} \right] \\
 &= -\frac{1}{Z} \text{tr} \left[ p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} p_q p_k e^{-H\tau} \right]
 \end{aligned} \tag{3.44}$$

$$\begin{aligned}
 C_{Q=2,S=1,S^3=-1;k,q,r,s} &= \begin{bmatrix} p_+p_+p_+p_+^\dagger & p_+p_+p_+p_-^\dagger & p_+p_+p_-p_+^\dagger & p_+p_+p_-p_-^\dagger \\ p_+p_-p_+p_+^\dagger & p_+p_-p_+p_-^\dagger & p_+p_-p_-p_+^\dagger & p_+p_-p_-p_-^\dagger \\ p_-p_+p_+p_+^\dagger & p_-p_+p_+p_-^\dagger & p_-p_+p_-p_+^\dagger & p_-p_+p_-p_-^\dagger \\ p_-p_-p_+p_+^\dagger & p_-p_-p_+p_-^\dagger & p_-p_-p_-p_+^\dagger & p_-p_-p_-p_-^\dagger \end{bmatrix}_{k,q,r,s} (\tau) = \\
 &= \begin{bmatrix} p_+^\dagger p_+^\dagger p_+p_+ & p_+^\dagger p_-^\dagger p_+p_+ & p_-^\dagger p_+^\dagger p_+p_+ & p_-^\dagger p_-^\dagger p_+p_+ \\ p_+^\dagger p_+^\dagger p_+p_- & p_+^\dagger p_-^\dagger p_+p_- & p_-^\dagger p_+^\dagger p_+p_- & p_-^\dagger p_-^\dagger p_+p_- \\ p_+^\dagger p_+^\dagger p_-p_+ & p_+^\dagger p_-^\dagger p_-p_+ & p_-^\dagger p_+^\dagger p_-p_+ & p_-^\dagger p_-^\dagger p_-p_+ \\ p_+^\dagger p_+^\dagger p_-p_- & p_+^\dagger p_-^\dagger p_-p_- & p_-^\dagger p_+^\dagger p_-p_- & p_-^\dagger p_-^\dagger p_-p_- \end{bmatrix}_{r,s,k,q} (\beta - \tau) \\
 &= \begin{bmatrix} p_+^\dagger p_+^\dagger p_+p_+ & p_-^\dagger p_+^\dagger p_+p_+ & p_+^\dagger p_-^\dagger p_+p_+ & p_-^\dagger p_-^\dagger p_+p_+ \\ p_+^\dagger p_+^\dagger p_-p_+ & p_-^\dagger p_+^\dagger p_-p_+ & p_+^\dagger p_-^\dagger p_-p_+ & p_-^\dagger p_-^\dagger p_-p_+ \\ p_+^\dagger p_+^\dagger p_+p_- & p_-^\dagger p_+^\dagger p_+p_- & p_+^\dagger p_-^\dagger p_+p_- & p_-^\dagger p_-^\dagger p_+p_- \\ p_+^\dagger p_+^\dagger p_-p_- & p_-^\dagger p_+^\dagger p_-p_- & p_+^\dagger p_-^\dagger p_-p_- & p_-^\dagger p_-^\dagger p_-p_- \end{bmatrix}_{s,r,q,k} (\beta - \tau) \\
 &= - \begin{bmatrix} p_+^\dagger p_+^\dagger p_+p_+ & p_-^\dagger p_+^\dagger p_+p_+ & p_+^\dagger p_-^\dagger p_+p_+ & p_-^\dagger p_-^\dagger p_+p_+ \\ p_+^\dagger p_+^\dagger p_+p_- & p_-^\dagger p_+^\dagger p_+p_- & p_+^\dagger p_-^\dagger p_+p_- & p_-^\dagger p_-^\dagger p_+p_- \\ p_+^\dagger p_+^\dagger p_-p_+ & p_-^\dagger p_+^\dagger p_-p_+ & p_+^\dagger p_-^\dagger p_-p_+ & p_-^\dagger p_-^\dagger p_-p_+ \\ p_+^\dagger p_+^\dagger p_-p_- & p_-^\dagger p_+^\dagger p_-p_- & p_+^\dagger p_-^\dagger p_-p_- & p_-^\dagger p_-^\dagger p_-p_- \end{bmatrix}_{s,r,k,q} (\beta - \tau) \\
 &= - \begin{bmatrix} p_+^\dagger p_+^\dagger p_+p_+ & p_+^\dagger p_-^\dagger p_+p_+ & p_-^\dagger p_+^\dagger p_+p_+ & p_-^\dagger p_-^\dagger p_+p_+ \\ p_+^\dagger p_+^\dagger p_-p_+ & p_+^\dagger p_-^\dagger p_-p_+ & p_-^\dagger p_+^\dagger p_-p_+ & p_-^\dagger p_-^\dagger p_-p_+ \\ p_+^\dagger p_+^\dagger p_+p_- & p_+^\dagger p_-^\dagger p_+p_- & p_-^\dagger p_+^\dagger p_+p_- & p_-^\dagger p_-^\dagger p_+p_- \\ p_+^\dagger p_+^\dagger p_-p_- & p_+^\dagger p_-^\dagger p_-p_- & p_-^\dagger p_+^\dagger p_-p_- & p_-^\dagger p_-^\dagger p_-p_- \end{bmatrix}_{r,s,q,k} (\beta - \tau)
 \end{aligned} \tag{3.45}$$

$$\begin{aligned}
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= C_{Q=2,S=1,S^3=+1;r,s,k,q}^\top (\beta - \tau) \\
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot C_{Q=2,S=1,S^3=+1;s,r,q,k}^\top (\beta - \tau) \cdot \sigma_5 \\
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -C_{Q=2,S=1,S^3=+1;s,r,k,q}^\top (\beta - \tau) \cdot \sigma_5 \\
 C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot C_{Q=2,S=1,S^3=+1;r,s,q,k}^\top (\beta - \tau)
 \end{aligned} \tag{3.46}$$

$$I : H - \vec{\mu} \cdot \vec{q}$$

$$\begin{aligned}
C_{Q=2, S=1, S^3=-1; k, q, r, s} &= \frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ I^{-1} I p_k I^{-1} I p_q I^{-1} I e^{-H\tau} I^{-1} I p_r^\dagger I^{-1} I p_s^\dagger I^{-1} I e^{-H(\beta-\tau)} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ I p_k I^{-1} I p_q I^{-1} I e^{-H\tau} I^{-1} I p_r^\dagger I^{-1} I p_s^\dagger I^{-1} I e^{-H(\beta-\tau)} I^{-1} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma h)_k^\dagger (\Sigma h)_q^\dagger e^{-H\tau} (\Sigma h)_r (\Sigma h)_s e^{-H(\beta-\tau)} \right] \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma h)_q^\dagger (\Sigma h)_k^\dagger e^{-H\tau} (\Sigma h)_s (\Sigma h)_r e^{-H(\beta-\tau)} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma h)_q^\dagger (\Sigma h)_k^\dagger e^{-H\tau} (\Sigma h)_r (\Sigma h)_s e^{-H(\beta-\tau)} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma h)_k^\dagger (\Sigma h)_q^\dagger e^{-H\tau} (\Sigma h)_s (\Sigma h)_r e^{-H(\beta-\tau)} \right] \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma h)_r (\Sigma h)_s e^{-H(\beta-\tau)} (\Sigma h)_k^\dagger (\Sigma h)_q^\dagger e^{-H\tau} \right] \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma h)_s (\Sigma h)_r e^{-H(\beta-\tau)} (\Sigma h)_q^\dagger (\Sigma h)_k^\dagger e^{-H\tau} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma h)_s (\Sigma h)_r e^{-H(\beta-\tau)} (\Sigma h)_k^\dagger (\Sigma h)_q^\dagger e^{-H\tau} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma h)_r (\Sigma h)_s e^{-H(\beta-\tau)} (\Sigma h)_q^\dagger (\Sigma h)_k^\dagger e^{-H\tau} \right]
\end{aligned} \tag{3.47}$$

[illegible]

$$\begin{aligned}
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_1 \cdot C_{Q=-2,S=1,S^3=+1;k,q,r,s}(\tau) \cdot \sigma_1 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot \sigma_1 \cdot C_{Q=-2,S=1,S^3=+1;q,k,s,r}(\tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot \sigma_1 \cdot C_{Q=-2,S=1,S^3=+1;q,k,r,s}(\tau) \cdot \sigma_1 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_1 \cdot C_{Q=-2,S=1,S^3=+1;k,q,s,r}(\tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_1 \cdot C_{Q=-2,S=1,S^3=-1;r,s,k,q}^\top(\beta - \tau) \cdot \sigma_1 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot \sigma_1 \cdot C_{Q=-2,S=1,S^3=-1;s,r,q,k}^\top(\beta - \tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_1 \cdot C_{Q=-2,S=1,S^3=-1;s,r,k,q}^\top(\beta - \tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot \sigma_1 \cdot C_{Q=-2,S=1,S^3=-1;r,s,q,k}^\top(\beta - \tau) \cdot \sigma_1
\end{aligned} \tag{3.49}$$

**$C : H$**

$$\begin{aligned}
C_{Q=2,S=1,S^3=-1;k,q,r,s} &= \frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ C^{-1} C p_k C^{-1} C p_q C^{-1} C e^{-H\tau} C^{-1} C p_r^\dagger C^{-1} C p_s^\dagger C^{-1} C e^{-H(\beta-\tau)} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ C p_k C^{-1} C p_q C^{-1} C e^{-H\tau} C^{-1} C p_r^\dagger C^{-1} C p_s^\dagger C^{-1} C e^{-H(\beta-\tau)} C^{-1} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ h_k h_q e^{-H\tau} h_r^\dagger h_s^\dagger e^{-H(\beta-\tau)} \right] \\
&= \frac{1}{Z} \text{tr} \left[ h_q h_k e^{-H\tau} h_s^\dagger h_r^\dagger e^{-H(\beta-\tau)} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ h_q h_k e^{-H\tau} h_r^\dagger h_s^\dagger e^{-H(\beta-\tau)} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ h_k h_q e^{-H\tau} h_s^\dagger h_r^\dagger e^{-H(\beta-\tau)} \right] \\
&= \frac{1}{Z} \text{tr} \left[ h_r^\dagger h_s^\dagger e^{-H(\beta-\tau)} h_k h_q e^{-H\tau} \right] \\
&= \frac{1}{Z} \text{tr} \left[ h_s^\dagger h_r^\dagger e^{-H(\beta-\tau)} h_q h_k e^{-H\tau} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ h_s^\dagger h_r^\dagger e^{-H(\beta-\tau)} h_k h_q e^{-H\tau} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ h_r^\dagger h_s^\dagger e^{-H(\beta-\tau)} h_q h_k e^{-H\tau} \right]
\end{aligned} \tag{3.50}$$

[illegible]



$$\begin{aligned}
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= C_{Q=-2,S=1,S^3=-1;k,q,r,s}(\tau) \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot C_{Q=-2,S=1,S^3=-1;q,k,s,r}(\tau) \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot C_{Q=-2,S=1,S^3=-1;q,k,r,s}(\tau) \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -C_{Q=-2,S=1,S^3=-1;k,q,s,r}(\tau) \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= C_{Q=-2,S=1,S^3=+1;r,s,k,q}^\top(\beta - \tau) \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot C_{Q=-2,S=1,S^3=+1;s,r,q,k}^\top(\beta - \tau) \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -C_{Q=-2,S=1,S^3=+1;s,r,k,q}^\top(\beta - \tau) \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot C_{Q=-2,S=1,S^3=+1;r,s,q,k}^\top(\beta - \tau)
\end{aligned} \tag{3.52}$$

**$XF : H$**

$$\begin{aligned}
C_{Q=2,S=1,S^3=-1;k,q,r,s} &= \frac{1}{Z} \text{tr} \left[ p_k p_q e^{-H\tau} p_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ (XF)^{-1} (XF) p_k (XF)^{-1} (XF) p_q (XF)^{-1} (XF) e^{-H\tau} (XF)^{-1} (XF) p_r^\dagger (XF)^{-1} (XF) p_s^\dagger (XF)^{-1} (XF) e^{-H(\beta-\tau)} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ (XF) p_k (XF)^{-1} (XF) p_q (XF)^{-1} (XF) e^{-H\tau} (XF)^{-1} (XF) p_r^\dagger (XF)^{-1} (XF) p_s^\dagger (XF)^{-1} (XF) e^{-H(\beta-\tau)} (XF)^{-1} \right] = \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} (\Sigma p)_r (\Sigma p)_s e^{-H(\beta-\tau)} \right] \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_q^\dagger (\Sigma p)_k^\dagger e^{-H\tau} (\Sigma p)_s (\Sigma p)_r e^{-H(\beta-\tau)} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma p)_q^\dagger (\Sigma p)_k^\dagger e^{-H\tau} (\Sigma p)_r (\Sigma p)_s e^{-H(\beta-\tau)} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma p)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} (\Sigma p)_s (\Sigma p)_r e^{-H(\beta-\tau)} \right] \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_r (\Sigma p)_s e^{-H(\beta-\tau)} (\Sigma p)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} \right] \\
&= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_s (\Sigma p)_r e^{-H(\beta-\tau)} (\Sigma p)_q^\dagger (\Sigma p)_k^\dagger e^{-H\tau} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma p)_s (\Sigma p)_r e^{-H(\beta-\tau)} (\Sigma p)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} \right] \\
&= -\frac{1}{Z} \text{tr} \left[ (\Sigma p)_r (\Sigma p)_s e^{-H(\beta-\tau)} (\Sigma p)_q^\dagger (\Sigma p)_k^\dagger e^{-H\tau} \right]
\end{aligned} \tag{3.53}$$

27th July 2022 18:10

$$\begin{aligned}
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_1 \cdot C_{Q=2,S=1,S^3=+1;k,q,r,s}(\tau) \cdot \sigma_1 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot \sigma_1 \cdot C_{Q=2,S=1,S^3=+1;q,k,s,r}(\tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot \sigma_1 \cdot C_{Q=2,S=1,S^3=+1;q,k,r,s}(\tau) \cdot \sigma_1 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_1 \cdot C_{Q=2,S=1,S^3=+1;k,q,s,r}(\tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_1 \cdot C_{Q=2,S=1,S^3=-1;r,s,k,q}^\top(\beta - \tau) \cdot \sigma_1 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= \sigma_5 \cdot \sigma_1 \cdot C_{Q=2,S=1,S^3=-1;s,r,q,k}^\top(\beta - \tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_1 \cdot C_{Q=2,S=1,S^3=-1;s,r,k,q}^\top(\beta - \tau) \cdot \sigma_1 \cdot \sigma_5 \\
C_{Q=2,S=1,S^3=-1;k,q,r,s}(\tau) &= -\sigma_5 \cdot \sigma_1 \cdot C_{Q=2,S=1,S^3=-1;r,s,q,k}^\top(\beta - \tau) \cdot \sigma_1
\end{aligned} \tag{3.55}$$

Table ?? shows the summary of all symmetries that we derived. From all these symmetries, with our data we can use only the anti-commutation and trace cyclicity, and if we calculate at half-filling, we could use also the  $XF$  symmetry. In the worst case, we can make our statistics 8 times better and 16 times better if we work at  $\mu = 0$ .

	$Q$	$S$	$S^3$	trafo	$kqrs$	$\tau$
Start with	2	1	-1	1	1234	$\tau$
Anti-commutation	2	1	-1	$\sigma_5 \cdot \sigma_5$	2143	$\tau$
	2	1	-1	$-\sigma_5 \cdot$	2134	$\tau$
	2	1	-1	$- \cdot \sigma_5$	1243	$\tau$
trace cycl	2	1	+1	$\top$	3412	$\beta - \tau$
	2	1	+1	$\sigma_5 \top \sigma_5$	4321	$\beta - \tau$
	2	1	+1	$-\top \sigma_5$	4312	$\beta - \tau$
	2	1	+1	$-\sigma_5 \top$	3421	$\beta - \tau$
I	-2	1	+1	$\sigma_1 \cdot \sigma_1$	1234	$\tau$
	-2	1	+1	$\sigma_5 \sigma_1 \cdot \sigma_1 \sigma_5$	2143	$\tau$
	-2	1	+1	$-\sigma_5 \sigma_1 \cdot \sigma_1$	2134	$\tau$
	-2	1	+1	$-\sigma_1 \cdot \sigma_1 \sigma_5$	1243	$\tau$
	-2	1	-1	$\sigma_1 \top \sigma_1$	3412	$\beta - \tau$
	-2	1	-1	$\sigma_5 \sigma_1 \top \sigma_1 \sigma_5$	4321	$\beta - \tau$
	-2	1	-1	$-\sigma_1 \top \sigma_1 \sigma_5$	4312	$\beta - \tau$
	-2	1	-1	$-\sigma_5 \sigma_1 \top \sigma_1$	3421	$\beta - \tau$
C	-2	1	-1	1	1234	$\tau$
	-2	1	-1	$\sigma_5 \cdot \sigma_5$	2143	$\tau$
	-2	1	-1	$-\sigma_5 \cdot$	2134	$\tau$
	-2	1	-1	$- \cdot \sigma_5$	1243	$\tau$
	-2	1	+1	$\top$	3412	$\beta - \tau$
	-2	1	+1	$\sigma_5 \top \sigma_5$	4321	$\beta - \tau$
	-2	1	+1	$-\top \sigma_5$	4312	$\beta - \tau$
	-2	1	+1	$-\sigma_5 \top$	3421	$\beta - \tau$
XF	2	1	+1	$\sigma_1 \cdot \sigma_1$	1234	$\tau$
	2	1	+1	$\sigma_5 \sigma_1 \cdot \sigma_1 \sigma_5$	2143	$\tau$
	2	1	+1	$-\sigma_5 \sigma_1 \cdot \sigma_1$	2134	$\tau$
	2	1	+1	$-\sigma_1 \cdot \sigma_1 \sigma_5$	1243	$\tau$
	2	1	-1	$\sigma_1 \top \sigma_1$	3412	$\beta - \tau$
	2	1	-1	$\sigma_5 \sigma_1 \top \sigma_1 \sigma_5$	4321	$\beta - \tau$
	2	1	-1	$-\sigma_1 \top \sigma_1 \sigma_5$	4312	$\beta - \tau$
	2	1	-1	$-\sigma_5 \sigma_1 \top \sigma_1$	3421	$\beta - \tau$

Table 3.3: All the correlators that can be directly averaged, if they all take a  $4 \times 4$  matrix form. The correlators above the double line may be averaged even at finite  $\mu$  and everything below may be averaged when  $\mu = 0$ . The combination of charge and spin indicate whether  $p^\dagger p^\dagger$  or  $pp$  is at the source (right-most). The *trafo* column indicates what must be done in the internal  $4 \times 4$  momentum space before averaging;  $\top$  indicates transpose,  $\sigma_1 \cdot \sigma_1$  indicates conjugation by  $\sigma_1$ ,  $\sigma_5 \cdot \sigma_5$  indicates conjugation by  $\sigma_5$ , minus sign indicates that there must be added one minus in front, and 1 indicates that no operation is needed. The next column *kqrs* indicates the position of the momenta at the sink and source relative to the initial correlator is. The  $\tau$  column indicates whether the given correlator needs to be time-reversed.

## **Spectre of Excitons**

---

### **4.1 Data Generating**

### **4.2 Fitting Data**



### Useful information

---

In the appendix you usually include extra information that should be documented in your thesis, but not interrupt the flow.

The  $\text{\LaTeX}$  WikiBook [12] is a useful source of information on  $\text{\LaTeX}$ .





# Bibliography

---

- [1] H. Kopka and P. W. Daly, *Guide to L<sup>A</sup>T<sub>E</sub>X*, 4th ed., Addison-Wesley, 2004 (cit. on p. 1).
- [2] F. Mittelbach and M. Goossens, *The L<sup>A</sup>T<sub>E</sub>X Companion*, 2nd ed., Addison-Wesley, 2004 (cit. on p. 1).
- [3] Particle Data Group, K. Nakamura et al., *Review of Particle Physics*, J. Phys. G **37** (2010) 075021, URL: <http://pdg.lbl.gov> (cit. on p. 1).
- [4] I. C. Brock,  
*Users Guide to Writing a Thesis in a Physics/Astronomy Institute of the Universität Bonn*,  
URL: [http://pi.physik.uni-bonn.de/pi\\_only/thesis.php](http://pi.physik.uni-bonn.de/pi_only/thesis.php) (visited on 01/07/2021)  
(cit. on p. 1).
- [5] ATLAS Collaboration, *Measurement of the top quark-pair production cross section with ATLAS in pp collisions at  $\sqrt{s} = 7$  TeV*, Eur. Phys. J. C **71** (2011) 1577, arXiv: [1012.1792](https://arxiv.org/abs/1012.1792) (cit. on p. 1).
- [6] F. Halzen and A. D. Martin,  
*Quarks and Leptons: An Introductory Course in Modern Particle Physics*, Wiley, 1984,  
ISBN: 9780471887416 (cit. on p. 1).
- [7] T. Loddenkötter, *Implementation of a kinematic fit of single top-quark production in association with a W boson and its application in a neural-network-based analysis in ATLAS*, BONN-IR-2012-06, PhD Thesis: University of Bonn, 2012,  
URL: [http://hss.ulb.uni-bonn.de/diss\\_online](http://hss.ulb.uni-bonn.de/diss_online) (cit. on p. 1).
- [8] S. Mergelmeyer, *D<sup>\*</sup>-Photoproduktion mit assoziierten Jets und Vertices bei ZEUS*, BONN-IB-2011-01, Universität Bonn, 2011,  
URL: [http://brock.physik.uni-bonn.de/zeus\\_pub.php](http://brock.physik.uni-bonn.de/zeus_pub.php) (cit. on p. 1).
- [9] O. S. Brüning et al., eds., *LHC Design Report. 1. The LHC Main Ring*, CERN-2004-003-V-1, CERN-2004-003, Geneva, 2004,  
URL: <https://cdsweb.cern.ch/record/782076> (cit. on p. 1).
- [10] ATLAS Collaboration, *Determination of the muon reconstruction efficiency in ATLAS at the Z resonance in proton-proton collisions at  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2011-008, CERN, 2011 (cit. on p. 1).
- [11] ATLAS Collaboration,  
*Expected Performance of the ATLAS Experiment – Detector, Trigger and Physics*, 2009,  
arXiv: [0901.0512](https://arxiv.org/abs/0901.0512) (cit. on p. 1).
- [12] L<sup>A</sup>T<sub>E</sub>X, URL: <https://en.wikibooks.org/wiki/LaTeX> (visited on 01/07/2021)  
(cit. on p. 33).



## List of Figures

---

3.1	Honeycomb lattice is a bipartite lattice because we can separate and label the lattice points with different colors (in this example blue and red) in such a way that every point in blue color neighbours only points with red color. . . . .	6
3.2	LEFT: All points of special interest shown on the lattice and how they are copied over to other lattice cells. RIGHT: Special points only in the first BZ. It contains all symmetries of the lattice . . . . .	6
3.3	An example of the spin-flip transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the spin-flip conjugation. . . . .	7
3.4	An example of the charge transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the charge-flip conjugation. . . . .	9
3.5	An example of the dagger-flip transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the dagger-flip conjugation. . . . .	12
3.6	An example of the particle-hole transformation with all possibilities of creation/annihilation operators. The graphs show only the transformation of the charge and the spin of the operators. The color of the arrows shows the charge (blue = +1, red = -1) and the orientation shows the spin. Left graph is the initial configuration of the operators and the right is after the particle-hole conjugation. . . . .	12



# List of Tables

---

- 3.1 All the correlators that can be directly averaged, assuming they all take the  $2 \times 2$  matrix form. The correlators above the double line may be averaged even at finite  $\mu$  and everything below may be averaged when  $\mu = 0$ . The combination of charge and spin indicate whether  $p^\dagger$ ,  $p$ ,  $h^\dagger$ , or  $h$  is right-most. The  $\vec{k}$  column indicates what must be done in the interval  $2 \times 2$  momentum space before averaging;  $\top$  indicates transpose,  $\sigma_1 \cdot \sigma_1$  indicates conjugation by  $\sigma_1$ , 1 indicates that no operation is needed. The  $\tau$  column indicates whether the given correlator needs to be time-reversed. . . . . 14
- 3.2 All the correlators that can be directly averaged, if they all take a  $4 \times 4$  matrix form. The correlators above the double line may be averaged even at finite  $\mu$  and everything below may be averaged when  $\mu = 0$ . The combination of charge and spin indicate whether  $h^\dagger p^\dagger$  or  $hp$  is at the source (right-most). The *trafo* column indicates what must be done in the interval  $4 \times 4$  momentum space before averaging;  $\top$  indicates transpose,  $\sigma_1 \cdot \sigma_1$  indicates conjugation by  $\sigma_1$ ,  $\sigma_5 \cdot \sigma_5$  indicates conjugation by  $\sigma_5$ , and 1 indicates that no operation is needed. The next column *kqrs* indicates the position of the momenta at the sink and source relative to the initial correlator is. The  $\tau$  column indicates whether the given correlator needs to be time-reversed. . . . . 20
- 3.3 All the correlators that can be directly averaged, if they all take a  $4 \times 4$  matrix form. The correlators above the double line may be averaged even at finite  $\mu$  and everything below may be averaged when  $\mu = 0$ . The combination of charge and spin indicate whether  $p^\dagger p^\dagger$  or  $pp$  is at the source (right-most). The *trafo* column indicates what must be done in the interval  $4 \times 4$  momentum space before averaging;  $\top$  indicates transpose,  $\sigma_1 \cdot \sigma_1$  indicates conjugation by  $\sigma_1$ ,  $\sigma_5 \cdot \sigma_5$  indicates conjugation by  $\sigma_5$ , minus sign indicates that there must be added one minus in front, and 1 indicates that no operation is needed. The next column *kqrs* indicates the position of the momenta at the sink and source relative to the initial correlator is. The  $\tau$  column indicates whether the given correlator needs to be time-reversed. . . . . 30