

# **Title of the Thesis**

Author's name

Masterarbeit in Physik  
angefertigt im Helmholtz-Institut für Strahlen- und  
Kernphysik

vorgelegt der  
Mathematisch-Naturwissenschaftlichen Fakultät  
der  
Rheinischen Friedrich-Wilhelms-Universität  
Bonn

MMM 2021

DRAFT

I hereby declare that this thesis was formulated by myself and that no sources or tools other than those cited were used.

Bonn, .....  
Date

.....  
Signature

- 1. Gutachter: Prof. Dr. John Smith
- 2. Gutachterin: Prof. Dr. Anne Jones

# Acknowledgements

---

I would like to thank ...

You should probably use `\chapter*` for acknowledgements at the beginning of a thesis and `\chapter` for the end.

DRAFT

DRAFT

# Contents

---

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Symmetries</b>	<b>3</b>
2.1	Correlation functions . . . . .	3
2.1.1	One-particle correlation function . . . . .	3
<b>A</b>	<b>Useful information</b>	<b>9</b>
	<b>Bibliography</b>	<b>11</b>
	<b>List of Figures</b>	<b>13</b>
	<b>List of Tables</b>	<b>15</b>

DRAFT

## Introduction

---

The introduction usually gives a few pages of introduction to the whole subject, maybe even starting with the Greeks.

For more information on L<sup>A</sup>T<sub>E</sub>X and the packages that are available see for example the books of Kopka [1] and Goossens et al [2].

A lot of useful information on particle physics can be found in the “Particle Data Book” [3].

I have resisted the temptation to put a lot of definitions into the file `thesis_defs.sty`, as everyone has their own taste as to what scheme they want to use for names. However, a few examples are included to help you get started:

- cross-sections are measured in pb and integrated luminosity in pb<sup>-1</sup>;
- the  $K_S^0$  is an interesting particle;
- the missing transverse momentum,  $p_T^{\text{miss}}$ , is often called missing transverse energy, even though it is calculated using a vector sum.

Note that the examples of units assume that you are using the `siunitx` package.

It also is probably a good idea to include a few well formatted references in the thesis skeleton. More detailed suggestions on what citation types to use can be found in the “Thesis Guide” [4]:

- articles in refereed journals [3, 5];
- a book [6];
- a PhD thesis [7] and a Diplom thesis [8];
- a collection of articles [9];
- a conference note [10];
- a preprint [11] (you can also use `@online` or `@booklet` for such things);
- something that is only available online [4].

At the end of the introduction it is normal to say briefly what comes in the following chapters.

The line at the beginning of this file is used by TeXstudio etc. to specify which is the master L<sup>A</sup>T<sub>E</sub>X file, so that you can compile your thesis directly from this file. If your thesis is called something other than `mythesis`, adjust it as appropriate.





## Symmetries

### 2.1 Correlation functions

#### 2.1.1 One-particle correlation function

We start with correlator  $\langle phh^\dagger p^\dagger \rangle$

$$C_{Q=0, S=1, S^3=+1; k, q, r, s} = \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k, q, r, s} \quad (\tau)$$

Now, we apply different symmetry operator to the two particle correlation function. When we apply them, we first start with the thermal trace and then switch to matrix form, so that it is easier to see the transformations needed in order to get another two particle correlator.

We obviously start with the trace cyclicity.

$$C_{Q=0, S=1, S^3=+1; k, q, r, s} = \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] = \frac{1}{Z} \text{tr} \left[ h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} p_k h_q e^{-H\tau} \right]$$

Now we apply time reverse

$$\beta \rightarrow \beta - \tau$$

and use the anti-commutation relations of the operators to get

$$\frac{1}{Z} \text{tr} \left[ h_r^\dagger p_s^\dagger e^{-H(\beta-(\beta-\tau))} p_k h_q e^{-H(\beta-\tau)} \right] = \frac{1}{Z} \text{tr} \left[ p_s^\dagger h_r^\dagger e^{-H\tau} h_q p_k e^{-H(\beta-\tau)} \right] \quad (2.1)$$

Now we switch to matrix form to see better what happens.

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=+1; k, q, r, s} \\
 &= \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k, q, r, s} \quad (\tau) \\
 &= \begin{bmatrix} h_+^\dagger p_+^\dagger p_+ h_+ & h_+^\dagger p_-^\dagger p_+ h_+ & h_-^\dagger p_+^\dagger p_+ h_+ & h_-^\dagger p_-^\dagger p_+ h_+ \\ h_+^\dagger p_+^\dagger p_+ h_- & h_+^\dagger p_-^\dagger p_+ h_- & h_-^\dagger p_+^\dagger p_+ h_- & h_-^\dagger p_-^\dagger p_+ h_- \\ h_+^\dagger p_+^\dagger p_- h_+ & h_+^\dagger p_-^\dagger p_- h_+ & h_-^\dagger p_+^\dagger p_- h_+ & h_-^\dagger p_-^\dagger p_- h_+ \\ h_+^\dagger p_+^\dagger p_- h_- & h_+^\dagger p_-^\dagger p_- h_- & h_-^\dagger p_+^\dagger p_- h_- & h_-^\dagger p_-^\dagger p_- h_- \end{bmatrix}_{r, s, k, q} \quad (\beta - \tau) \\
 &= \begin{bmatrix} p_+^\dagger h_+^\dagger h_+ p_+ & p_-^\dagger h_+^\dagger h_+ p_+ & p_+^\dagger h_-^\dagger h_+ p_+ & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_- p_+ \\ p_+^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_-^\dagger h_+ p_- & p_-^\dagger h_-^\dagger h_+ p_- \\ p_+^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_- \end{bmatrix}_{s, r, q, k} \quad (\beta - \tau) = C_{Q=0, S=1, S^3=-1; s, r, q, k}
 \end{aligned}$$

We can notice that order reverse is actually a composed of two transformations – particle-hole momentum switch and time reverse. After we stare at the matrix for an hour, we can figure out what the transformation between the correlation functions is.

$$C_{Q=0, S=1, S^3=+1; k, q, r, s}(\tau) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} C_{Q=0, S=1, S^3=-1; s, r, q, k}^\top (\beta - \tau) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

We repeat the steps again for each symmetry operator. First we start with the thermal trace and then switch to matrix for easy transformations.

$$I : H - \vec{\mu} \cdot \vec{q}$$

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=+1; k, q, r, s} \\
 &= \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ I^{-1} I p_k I^{-1} I h_q I^{-1} I e^{-H\tau} I^{-1} I h_r^\dagger I^{-1} I p_s^\dagger I^{-1} I e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ I p_k I^{-1} I h_q I^{-1} I e^{-H\tau} I^{-1} I h_r^\dagger I^{-1} I p_s^\dagger I^{-1} I e^{-H(\beta-\tau)} I^{-1} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma h)_k^\dagger (\Sigma p)_q^\dagger e^{-H\tau} (\Sigma p)_r (\Sigma h)_s e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_q^\dagger (\Sigma h)_k^\dagger e^{-H\tau} (\Sigma h)_s (\Sigma p)_r e^{-H(\beta-\tau)} \right]
 \end{aligned}$$

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=+1; k, q, r, s} \\
 &= I^{-1} \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k, q, r, s} (\tau) I \\
 &= \begin{bmatrix} (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_- \\ (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_+^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_- \\ (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_+^\dagger (\Sigma p)_- (\Sigma h)_- \\ (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_+ (\Sigma h)_- & (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_+ & (\Sigma h)_-^\dagger (\Sigma p)_-^\dagger (\Sigma p)_- (\Sigma h)_- \end{bmatrix} \\
 &= \begin{bmatrix} h_-^\dagger p_-^\dagger p_- h_- & h_-^\dagger p_-^\dagger p_- h_+ & h_-^\dagger p_-^\dagger p_+ h_- & h_-^\dagger p_-^\dagger p_+ h_+ \\ h_-^\dagger p_+^\dagger p_- h_- & h_-^\dagger p_+^\dagger p_- h_+ & h_-^\dagger p_+^\dagger p_+ h_- & h_-^\dagger p_+^\dagger p_+ h_+ \\ h_+^\dagger p_-^\dagger p_- h_- & h_+^\dagger p_-^\dagger p_- h_+ & h_+^\dagger p_-^\dagger p_+ h_- & h_+^\dagger p_-^\dagger p_+ h_+ \\ h_+^\dagger p_+^\dagger p_- h_- & h_+^\dagger p_+^\dagger p_- h_+ & h_+^\dagger p_+^\dagger p_+ h_- & h_+^\dagger p_+^\dagger p_+ h_+ \end{bmatrix}_{k, q, r, s} (\tau) \\
 &= \begin{bmatrix} p_- h_- h_-^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger \\ p_- h_- h_+^\dagger p_-^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_+^\dagger \end{bmatrix}_{r, s, k, q} (\beta - \tau) \\
 &= \begin{bmatrix} p_-^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_- p_+ \\ p_+^\dagger h_-^\dagger h_- p_- & p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_+ & p_+^\dagger h_+^\dagger h_- p_+ \\ p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_+^\dagger h_+ p_+ \end{bmatrix}_{q, k, s, r} (\tau)
 \end{aligned}$$

Depending on what transformation we use to get back to  $\langle phh^\dagger p^\dagger \rangle$ , we get two transformations which are time reverse to one another. We notice the same behaviour for all operators.

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=+1; k, q, r, s}(\tau) \\
 &= \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0, S=1, S^3=+1; r, s, k, q}^\top (\beta - \tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 & C_{Q=0, S=1, S^3=+1; k, q, r, s}(\tau) \\
 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0, S=1, S^3=-1; q, k, s, r}(\tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}
 \end{aligned}$$

$C : H$

$$\begin{aligned}
 C_{Q=0,S=1,S^3=+1;k,q,r,s} &= \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ C^{-1} C p_k C^{-1} C h_q C^{-1} C e^{-H\tau} C^{-1} C h_r^\dagger C^{-1} C p_s^\dagger C^{-1} C e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ h_k p_q e^{-H\tau} p_r^\dagger h_s^\dagger e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ p_q h_k e^{-H\tau} h_s^\dagger p_r^\dagger e^{-H(\beta-\tau)} \right]
 \end{aligned}$$

$$\begin{aligned}
 C_{Q=0,S=1,S^3=+1;k,q,r,s} &= C^{-1} \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k,q,r,s} (\tau) C \\
 &= \begin{bmatrix} h_+ p_+ p_+^\dagger h_+^\dagger & h_+ p_+ p_+^\dagger h_-^\dagger & h_+ p_+ p_-^\dagger h_+^\dagger & h_+ p_+ p_-^\dagger h_-^\dagger \\ h_+ p_- p_+^\dagger h_+^\dagger & h_+ p_- p_+^\dagger h_-^\dagger & h_+ p_- p_-^\dagger h_+^\dagger & h_+ p_- p_-^\dagger h_-^\dagger \\ h_- p_+ p_+^\dagger h_+^\dagger & h_- p_+ p_+^\dagger h_-^\dagger & h_- p_+ p_-^\dagger h_+^\dagger & h_- p_+ p_-^\dagger h_-^\dagger \\ h_- p_- p_+^\dagger h_+^\dagger & h_- p_- p_+^\dagger h_-^\dagger & h_- p_- p_-^\dagger h_+^\dagger & h_- p_- p_-^\dagger h_-^\dagger \end{bmatrix}_{k,q,r,s} (\tau) \\
 &= \begin{bmatrix} p_+^\dagger h_+^\dagger h_+ p_+ & p_+^\dagger h_-^\dagger h_+ p_+ & p_-^\dagger h_+^\dagger h_+ p_+ & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_- p_+ \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_- \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_- \end{bmatrix}_{r,s,k,q} (\beta - \tau) \\
 &= \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{q,k,s,r} (\tau)
 \end{aligned}$$

$$C_{Q=0,S=1,S^3=+1;k,q,r,s}(\tau) = C_{Q=0,S=1,S^3=-1;r,s,k,q}^\top(\beta - \tau)$$

$$C_{Q=0,S=1,S^3=+1;k,q,r,s}(\tau) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} C_{Q=0,S=1,S^3=+1;q,k,s,r}(\tau) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$XF : H$$

$$\begin{aligned}
 C_{Q=0,S=1,S^3=+1;k,q,r,s} &= \frac{1}{Z} \text{tr} \left[ p_k h_q e^{-H\tau} h_r^\dagger p_s^\dagger e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (XF)^{-1} (XF) p_k (XF)^{-1} (XF) h_q (XF)^{-1} (XF) e^{-H\tau} (XF)^{-1} (XF) h_r^\dagger (XF)^{-1} (XF) p_s^\dagger (XF)^{-1} (XF) e^{-H(\beta-\tau)} \right] \\
 &= \frac{1}{Z} \text{tr} \left[ (\Sigma p)_k^\dagger (\Sigma h)_q^\dagger e^{-H\tau} (\Sigma h)_r (\Sigma p)_s e^{-H(\beta-\tau)} \right]
 \end{aligned}$$

$$\begin{aligned}
 C_{Q=0,S=1,S^3=+1;k,q,r,s} &= (XF)^{-1} \begin{bmatrix} p_+ h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_+ h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_-^\dagger \\ p_- h_+ h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_- h_- h_+^\dagger p_-^\dagger & p_- h_- h_-^\dagger p_+^\dagger & p_- h_- h_-^\dagger p_-^\dagger \end{bmatrix}_{k,q,r,s} (\tau) (XF) \\
 &= \begin{bmatrix} (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_- \\ (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_+^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_- \\ (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_+^\dagger (\Sigma h)_- (\Sigma p)_- \\ (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_+ (\Sigma p)_- & (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_+ & (\Sigma p)_-^\dagger (\Sigma h)_-^\dagger (\Sigma h)_- (\Sigma p)_- \end{bmatrix} \\
 &= \begin{bmatrix} p_-^\dagger h_-^\dagger h_- p_- & p_-^\dagger h_-^\dagger h_- p_+ & p_-^\dagger h_-^\dagger h_+ p_- & p_-^\dagger h_-^\dagger h_+ p_+ \\ p_-^\dagger h_+^\dagger h_- p_- & p_-^\dagger h_+^\dagger h_- p_+ & p_-^\dagger h_+^\dagger h_+ p_- & p_-^\dagger h_+^\dagger h_+ p_+ \\ p_+^\dagger h_-^\dagger h_- p_- & p_+^\dagger h_-^\dagger h_- p_+ & p_+^\dagger h_-^\dagger h_+ p_- & p_+^\dagger h_-^\dagger h_+ p_+ \\ p_+^\dagger h_+^\dagger h_- p_- & p_+^\dagger h_+^\dagger h_- p_+ & p_+^\dagger h_+^\dagger h_+ p_- & p_+^\dagger h_+^\dagger h_+ p_+ \end{bmatrix}_{k,q,r,s} (\tau) \\
 &= \begin{bmatrix} p_- h_- h_-^\dagger p_-^\dagger & p_+ h_- h_-^\dagger p_-^\dagger & p_- h_+ h_-^\dagger p_-^\dagger & p_+ h_+ h_-^\dagger p_-^\dagger \\ p_- h_- h_+^\dagger p_-^\dagger & p_+ h_- h_+^\dagger p_-^\dagger & p_- h_+ h_+^\dagger p_-^\dagger & p_+ h_+ h_+^\dagger p_-^\dagger \\ p_- h_- h_-^\dagger p_+^\dagger & p_+ h_- h_-^\dagger p_+^\dagger & p_- h_+ h_-^\dagger p_+^\dagger & p_+ h_+ h_-^\dagger p_+^\dagger \\ p_- h_- h_+^\dagger p_+^\dagger & p_+ h_- h_+^\dagger p_+^\dagger & p_- h_+ h_+^\dagger p_+^\dagger & p_+ h_+ h_+^\dagger p_+^\dagger \end{bmatrix}_{s,r,k,q} (\beta - \tau)
 \end{aligned}$$

$$\begin{aligned}
 C_{Q=0,S=1,S^3=+1;k,q,r,s}(\tau) &= \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0,S=1,S^3=-1;k,q,r,s}(\tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 C_{Q=0,S=1,S^3=+1;k,q,r,s}(\tau) &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} C_{Q=0,S=1,S^3=+1;s,r,q,k}^\top(\beta - \tau) \begin{bmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}
 \end{aligned}$$

The table below shows the summary of all symmetries that we calculated.

For more compact form, we write

	$Q$	$S$	$S^3$	trafo	$kqrs$	$\tau$
Start with trace cycl I	0	1	+1	1	1234	$\tau$
	0	1	-1	$\sigma_5 \top \sigma_5$	4321	$\beta - \tau$
	0	1	+1	$\sigma_1 \top \sigma_1$	3412	$\beta - \tau$
	0	1	-1	$\sigma_5 \sigma_1 \cdot \sigma_1 \sigma_5$	2143	$\tau$
C	0	1	-1	$\top$	3412	$\beta - \tau$
	0	1	+1	$\sigma_5 \cdot \sigma_5$	2143	$\tau$
XF	0	1	-1	$\sigma_1 \cdot \sigma_1$	1234	$\tau$
	0	1	+1	$\sigma_5 \sigma_1 \top \sigma_1 \sigma_5$	4321	$\beta - \tau$

$$\sigma_5 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

### Useful information

---

In the appendix you usually include extra information that should be documented in your thesis, but not interrupt the flow.

The  $\text{\LaTeX}$  WikiBook [12] is a useful source of information on  $\text{\LaTeX}$ .





# Bibliography

---

- [1] H. Kopka and P. W. Daly, *Guide to L<sup>A</sup>T<sub>E</sub>X*, 4th ed., Addison-Wesley, 2004 (cit. on p. 1).
- [2] F. Mittelbach and M. Goossens, *The L<sup>A</sup>T<sub>E</sub>X Companion*, 2nd ed., Addison-Wesley, 2004 (cit. on p. 1).
- [3] Particle Data Group, K. Nakamura et al., *Review of Particle Physics*, J. Phys. G **37** (2010) 075021, URL: <http://pdg.lbl.gov> (cit. on p. 1).
- [4] I. C. Brock,  
*Users Guide to Writing a Thesis in a Physics/Astronomy Institute of the Universität Bonn*,  
URL: [http://pi.physik.uni-bonn.de/pi\\_only/thesis.php](http://pi.physik.uni-bonn.de/pi_only/thesis.php) (visited on 01/07/2021)  
(cit. on p. 1).
- [5] ATLAS Collaboration, *Measurement of the top quark-pair production cross section with ATLAS in pp collisions at  $\sqrt{s} = 7$  TeV*, Eur. Phys. J. C **71** (2011) 1577, arXiv: [1012.1792](https://arxiv.org/abs/1012.1792) (cit. on p. 1).
- [6] F. Halzen and A. D. Martin,  
*Quarks and Leptons: An Introductory Course in Modern Particle Physics*, Wiley, 1984,  
ISBN: 9780471887416 (cit. on p. 1).
- [7] T. Loddenkötter, *Implementation of a kinematic fit of single top-quark production in association with a W boson and its application in a neural-network-based analysis in ATLAS*, BONN-IR-2012-06, PhD Thesis: University of Bonn, 2012,  
URL: [http://hss.ulb.uni-bonn.de/diss\\_online](http://hss.ulb.uni-bonn.de/diss_online) (cit. on p. 1).
- [8] S. Mergelmeyer, *D<sup>\*</sup>-Photoproduktion mit assoziierten Jets und Vertices bei ZEUS*, BONN-IB-2011-01, Universität Bonn, 2011,  
URL: [http://brock.physik.uni-bonn.de/zeus\\_pub.php](http://brock.physik.uni-bonn.de/zeus_pub.php) (cit. on p. 1).
- [9] O. S. Brüning et al., eds., *LHC Design Report. 1. The LHC Main Ring*, CERN-2004-003-V-1, CERN-2004-003, Geneva, 2004,  
URL: <https://cdsweb.cern.ch/record/782076> (cit. on p. 1).
- [10] ATLAS Collaboration, *Determination of the muon reconstruction efficiency in ATLAS at the Z resonance in proton-proton collisions at  $\sqrt{s} = 7$  TeV*, ATLAS-CONF-2011-008, CERN, 2011 (cit. on p. 1).
- [11] ATLAS Collaboration,  
*Expected Performance of the ATLAS Experiment – Detector, Trigger and Physics*, 2009,  
arXiv: [0901.0512](https://arxiv.org/abs/0901.0512) (cit. on p. 1).
- [12] L<sup>A</sup>T<sub>E</sub>X, URL: <https://en.wikibooks.org/wiki/LaTeX> (visited on 01/07/2021)  
(cit. on p. 9).



## List of Figures

---



## List of Tables

---