

Additional Problem 1

We know that the limit of the Rieman sums is equal to the integral. Thus we have

$$\left| \int_a^b F(s) ds \right| = \left| \lim_{N \rightarrow \infty} \sum_{i=0}^N F(x_i) \Delta x_N \right|$$

where $\Delta x_N = \frac{b-a}{N}$ and $x_i = a + i\Delta x_N$. Since $|\cdot|$ is continuous we can interchange the limit and the norm and apply the triangle inequality to conclude

$$\leq \lim_{N \rightarrow \infty} \sum_{i=0}^N |F(x_i) \Delta x_N| = \int_a^b |F(s)| ds$$

Additional Problem 2

(a) Notice the following

$$|\gamma_{j+1}(t) - \gamma_j(t)| = \left| \int_0^t F(\gamma_j(s)) - F(\gamma_{j-1}(s)) ds \right|$$

from problem 1

$$\leq \int_0^t |F(\gamma_j(s)) - F(\gamma_{j-1}(s))| ds$$

By Lipshitz condition

$$\leq \int_0^t L |\gamma_j(s) - \gamma_{j-1}(s)| ds$$

Now we can consisely state our inductive argument:

For the base case, notice that

$$\begin{aligned} \gamma_0(t) &= p \\ \gamma_1(t) &= p + \int_0^t F(p) ds = p + Mt \end{aligned}$$

Thus from the inequality established above

$$\begin{aligned} |\gamma_2(t) - \gamma_1(t)| &\leq L \int_0^t |\gamma_1(s) - \gamma_0(s)| ds \\ &= L \int_0^t Ms ds = \frac{MLt^2}{2} \end{aligned}$$

For the inductive step we have

$$|\gamma_{j+1}(t) - \gamma_j(t)| \leq \int_0^t L |\gamma_j(s) - \gamma_{j-1}(s)| ds$$

which by the inductive hypothesis

$$\leq \int_0^t L \frac{ML^{j-1}t^j}{j!} = \frac{ML^j t^{j+1}}{(j+1)!}$$

(b) Notice that we have a Taylor series which by Taylors Theorem converges

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{ML^j T^{j+1}}{(j+1)!} = MT e^{LT}$$

we have for $t \in [0, T]$

$$\sum_{j=1}^N |\gamma_{j+1}(t) - \gamma_j(t)| \leq \sum_{j=1}^N \frac{ML^j t^{j+1}}{(j+1)!} \leq \sum_{j=1}^N \frac{ML^j T^{j+1}}{(j+1)!}$$

by the Weierstrass M-test we thus have uniform convergence on $[0, T]$

(c) We have

$$\gamma(s) = \lim_{n \rightarrow \infty} \gamma_n(s) = \lim_{n \rightarrow \infty} \int_0^s F(\gamma_{n-1}(t)) dt$$

Since we have uniform convergence we can interchange order of limits and integration

$$= \int_0^s \lim_{n \rightarrow \infty} F(\gamma_{n-1}(t)) dt = \int_0^s F(\gamma(t)) dt$$

Additional Problem 3

We have

$$|T_A x|_{\max} = \max_i |A_{i1}x_1 + A_{i2}x_2 + \dots A_{im}x_m|$$

For each i we have

$$|A_{i1}x_1 + A_{i2}x_2 + \dots A_{im}x_m| \leq |A_{i1}x_1| + |A_{i2}x_2| + \dots |A_{im}x_m| \leq L|x|_1$$

Thus

$$|T_A x|_{\max} \leq L|x|_1$$

L is the smallest value to establish this inequality since letting $A_{kl} = L$, letting $x \in \mathbb{R}^m$ be such that $x_l = 1$ and $x_i = 0$ for $i \neq l$, we have that $|x|_1 = 1$ and

$$|A_{l1}x_1 + A_{l2}x_2 + \dots A_{lk}x_l + \dots A_{lm}x_m| = L = L|x|_1$$

and thus for this choice of x

$$|T_A x|_{\max} = L|x|_1$$

For the second part

$$\begin{aligned}
|T_A x|_E &= \sqrt{\sum_{i=1}^n \left(\sum_{j=1}^m A_{ij} x_j \right)^2} \\
&\leq \sqrt{n} |T_A x|_{\max} \leq \max A_{ij} \sqrt{nm} |x|_1 \leq \max A_{ij} \sqrt{nm} |x|_E = L |x|_E
\end{aligned}$$

For an example where we can get a stronger bound let T_A be the identity for $m = n > 1$. We have

$$|T_A x|_E = |x|_E < \sqrt{mn} |x|_E = L |x|_E$$

Thus we can choose $L = 1$

For an example where L is the best bound notice that for $m = n = 1$ $T_A x$ takes the form ax for $a \in \mathbb{R}$ and thus

$$|ax|_E = |a| |x| = L |x|$$

Additional Problem 4

(\Rightarrow) If A is invertable then assume for contradiction we can choose $x_n \neq 0$ where we can scale $|x_n|_E = 1$ so that

$$|T_A x_n|_E \leq \frac{1}{n} |x_n|_E$$

Notice that the unit ball $\{x \in \mathbb{R}^m : |x|_E = 1\}$ is compact and thus we get a convergent subsequence with a limit $x \neq 0$. We have that

$$|T_A x|_E < \frac{1}{n} \forall n \Rightarrow T_A x = 0$$

which contradicts T_A have trivial nullspace

(\Leftarrow) If A is not invertable then the kernel is nontrivial so there is some $x \neq 0 \in \mathbb{R}^m$ such that

$$|T_A x|_E = 0 \not\geq c |x|_E$$

for all c (proved by contrapositive)

If (*) is true then we have

$$c |T_{A^{-1}} x|_E \leq |T_A T_{A^{-1}} x|_E = |x|_E$$

Thus

$$|T_{A^{-1}} x|_E \leq c^{-1} |x|_E$$