

**Exercise 19**

(a) Consider any limit point  $x$  of  $D_k$ . We will show  $\text{osc}(x) \geq 1/k \Rightarrow x \in D_k$  so  $D_k$  is closed. (We will use the diameter definition of  $\text{osc}$ , where  $\text{osc}(x) = \lim_{r \rightarrow 0} \text{diam } f(B_r(x))$ ) For any  $r > 0$ , since  $x$  is a limit point of  $D_k$ , there exists  $y \in B_r(x) \cap D_k$ . Since  $B_r(x)$  is open there exists  $B_s(y) \subset B_r(x)$ . Since  $y \in D_k$  we know that  $\text{diam } f(B_s(y)) \geq 1/k$ . Since  $B_s(y) \subset B_r(x)$  we have

$$\text{diam } f(B_r(x)) \geq \text{diam } f(B_s(y)) \geq 1/k$$

Thus for every  $r > 0$   $\text{diam } f(B_r(x)) \geq 1/k$  so we know

$$\text{osc}(x) = \lim_{r \rightarrow 0} \text{diam } f(B_r(x)) \geq 1/k$$

(b) Since every discontinuity point has  $\text{osc} > 0$ , the discontinuity set can be written as a countable union of  $D_k$  where each  $D_k$  is closed as proven in part a.

$$D = \bigcup_{k=1}^{\infty} D_k$$

(c) Since the continuity set is the complement of the discontinuity set, we have

$$C = [a, b] \setminus \left( \bigcup_{k=1}^{\infty} D_k \right)$$

From Demorgans law

$$C = \bigcap_{k=1}^{\infty} ([a, b] \setminus D_k)$$

Which is a countable intersection of open sets (since the complement of closed sets are open so  $[a, b] \setminus D_k$  is open)

**Exercise 27**

(b) Consider the indicator function on the rationals  $\chi_{\mathbb{Q}} : [0, 1] \rightarrow \mathbb{R}$ . We have that for any  $n \in \mathbb{N}$ ,

$$x_k^* = \frac{2a + (2k - 1)(b - a)}{2n} \in \mathbb{Q}$$

Thus  $\chi_{\mathbb{Q}}(x_k^*) = 1$  for all  $k, n$  so our calc limit yields 1, while  $\chi_{\mathbb{Q}}$  is not Riemann integrable since it is continuous nowhere

**Exercise 28**

$(i \Rightarrow ii)$  : This follows directly from the definition of a zero set. If  $Z$  is a zero set, then for each  $\epsilon > 0$  there is a countable covering of  $Z$  by open intervals  $(a_i, b_i)$  with total length  $\sum b_i - a_i < \epsilon$ . We can replace  $(a_i, b_i)$  with  $[a_i, b_i]$  and since  $(a_i, b_i) \subset [a_i, b_i]$  this is still a covering of  $Z$ , and the lengths are the same.

$(ii \Rightarrow i)$  : Given  $\epsilon > 0$  from  $ii$  there exists a countable covering  $C_i = [a_i, b_i]$  with total length  $\sum b_i - a_i < \epsilon/2$ . Since the covering is countable we can replace each  $C_i = [a_i, b_i]$  with  $U_i = (a_i - \frac{\epsilon}{2} \frac{1}{4^i}, b_i + \frac{\epsilon}{2} \frac{1}{4^i})$ . The  $U_i$  make up a covering since each  $C_i \subset U_i$  so  $Z \subset \bigcup C_i \subset \bigcup U_i$ . The total length is

$$\sum_{i=1}^{\infty} b_i + \frac{\epsilon}{2} \frac{1}{4^i} - (a_i - \frac{\epsilon}{2} \frac{1}{4^i}) = \sum b_i - a_i + \frac{\epsilon}{2} \sum_{i=1}^{\infty} \frac{1}{2^i} = \frac{\epsilon}{2} + \sum b_i - a_i < \epsilon$$

### Exercise Additional Problem 1

For any  $\epsilon > 0$  and  $x \in [0, 1] \setminus \mathbb{Q}$  we have that there exists  $k \in \mathbb{N}$  such that  $\frac{1}{k} < \epsilon$ . We have that there are only finitely many  $\frac{p}{q} \in \mathbb{Q}$  where  $q < k$  (we can bound the number of these rational points from above by the sum of denominators less than  $k$  which is finite). Thus there exists a minimum distance  $d > 0$  from these numbers and  $x$  (this is because there are a finite number of distances, each of which are  $> 0$  since  $x \notin \mathbb{Q}$  so not equal to any of these numbers). We have that  $\forall y \in B_d(x)$ ,  $f(y) < \epsilon$  and thus  $f$  is continuous at  $x$ . The reason for this is as follows, if  $y \notin \mathbb{Q}$  then  $f(y) = 0$  and we are done. If  $y \in \mathbb{Q}$  then we have  $|y - x| < d$  thus  $y$  cannot have denominator  $< k$  since otherwise we would contradict the minimality of  $d$ . Thus  $f(y) \leq \frac{1}{k} < \epsilon$  so  $f(y) < \epsilon$ .

### Exercise Additional Problem 2

If  $x \in \partial S$ , then for all  $r > 0$ , we have  $B_r(x) \cap S, B_r(x) \cap S^c \neq \emptyset$ . Thus there exists  $y, z \in B_r(x)$  where  $\chi_S(y) = 0$  and  $\chi_S(z) = 1$ . Thus  $\text{diam } \chi_S(B_r(x)) \geq 1$  since  $\chi_S$  cannot take any values besides 1 and 0, we have equality:  $\text{diam } \chi_S(B_r(x)) = 1$  for all  $r > 0$

$$\text{osc}(x) = \lim_{r \rightarrow 0} \text{diam } \chi_S(B_r(x)) = 1$$

If  $x \notin \partial S$  then there exists  $r > 0$  such that  $B_r(x) \subset S$  or  $S^c$ . We thus have  $\chi_S(B_r(x)) = \{1\}$  or  $\{0\}$  which are sets with diameter 0. Thus  $\text{osc}(x) = 0$