

**2-3, 1**

$A$  is continuous since its components are continuous functions. Given any point  $p \in S^2$  we can choose parametrizations  $x_1 : U_1 \subset \mathbb{R}^2 \rightarrow S^2$ ,  $x_2 : U_2 \subset \mathbb{R}^2 \rightarrow S^2$  such that  $p \in x_1(U_1)$ ,  $A(p) \in x_2(U_2)$ .

From class we established there are the following parametrizations we can choose from:

$$x_1, x_2 \in \{(\cos \theta \sin \varphi, \sin \theta \sin \varphi, \cos \varphi), (\cos \theta \sin \varphi + \pi, \sin \theta \sin \varphi + \pi, \cos \varphi + \pi), \\ (\cos \theta + \pi \sin \varphi, \sin \theta + \pi \sin \varphi, \cos \varphi) \dots \}$$

We must show that

$$x_2^{-1} \circ A \circ x_1$$

is differentiable at  $p$ . Notice that  $A = A^{-1}$  and thus by showing  $A$  is differentiable we have shown its inverse to be differentiable. Hence concluding  $A$  is a diffeomorphism.

It is clear this composition is differentiable since each component

$$x_2^{-1} \circ A \circ x_1(p) = (x_{2,1}^{-1}(-x_{1,1}(p_1)), x_{2,2}^{-1}(-x_{1,2}(p_2)), x_{2,3}^{-1}(-x_{1,3}(p_3)))$$

Is a composition of differentiable functions and thus differentiable regardless of choice of  $x_1, x_2$ .

**2-3 3**

We have the diffeomorphism  $f : \mathbb{R}^2 \rightarrow P$  with  $f(x, y) = (x, y, z^2)$ . To establish  $f$  is a diffeomorphism, notice that  $f$  is also a parametrization. Thus the conditions of being a diffeomorphism rely on asking whether over any open set if

$$f^{-1} \circ f \circ \text{id}$$

is differentiable. This mapping is the identity mapping so clearly differentiable.

**2-3 4**

The diffeomorphism  $f : S^2 \rightarrow E$  is defined by

$$f(x, y, z) = (ax, by, cz)$$

We have that this is a diffeomorphism since notice that for any parametrization  $x_1 : U \subset \mathbb{R}^2 \rightarrow S^2$  around a point  $p \in S^2$  we have a parametrization  $x_2 = f \circ x_1$  of  $E$  around  $f(p)$ . Thus we have

$$x_1^{-1} \circ f \circ x_2 = x_1^{-1} \circ f^{-1} \circ f \circ x_1 = \text{id}$$

is differentiable

**2-3 6**

For diffeomorphism  $f : S_1 \rightarrow S_2$ , suppose we have the parameterizations around a point  $p \in S_1$  and  $f(p) \in S_2$

$$x_1 : U \subset \mathbb{R}^2 \rightarrow S_1, x_2 : V \subset \mathbb{R}^2 \rightarrow S_2$$

$$y_1 : U \subset \mathbb{R}^2 \rightarrow S_1, y_2 : V \subset \mathbb{R}^2 \rightarrow S_2$$

We need to show that if the parameterization with  $x_1, x_2$  establishes  $f$  to be differentiable then so does  $y_1, y_2$

**2-3 7**