

Supporting Information: Load-dependent sacroiliac joint motion in 278 pelvic finite-element models

Petr Henyš, Niels Hammer

1 Supplementary methods and reproducibility

All supplementary statistics use the corrected unloaded-subject reference $\mathbf{X} + \phi_i$ and loaded coordinates $\mathbf{X} + \phi_i + \mathbf{u}_{i\ell}$, as specified in the mathematical appendix of the main article. The earlier template-to-subject extraction is superseded. The three simulation variants share subject indices; resampling preserves their pairing.

Component differences compare bilateral absolute magnitudes, not signed motion vectors. Their bootstrap intervals describe medians; primary tests are exact binomial sign tests. Signed-rank results are retained only as machine-readable sensitivity diagnostics. The six fully adjusted sex comparisons use HC3 covariance and a single six-test FDR family. Variance ratios are non-additive and are not causal contributions.

The scaling model is

$$\begin{aligned} \log Y = & \beta_0 + \beta_v \log V + \beta_s \text{Female} + \beta_{vs} \log V \text{Female} \\ & + \beta_a \text{Age}_z + \beta_{AP} \text{AP}_z + \beta_b \text{Biischadic}_z + \beta_p \text{Subpubic}_z + \epsilon. \end{aligned} \quad (1)$$

Male and female slopes are β_v and $\beta_v + \beta_{vs}$. Allometry slope tests are corrected jointly across 20 sex-specific slopes; the ten interaction tests form a separate family. Table S4 is an exact reparameterization with $\log s = (\log V - \log V_{\text{template}})/3$, so scale exponents equal three times the volume exponents. It is not an independent validation or sensitivity analysis.

The accompanying reproduction guide specifies the complete command sequence for corrected extraction, extraction sensitivity, statistics, table formatting, numerical manuscript passages and publication figures. Original FE displacement archives are unchanged. Corrected checkpoints are stored separately from statistical outputs. Existing checkpoints must be removed or regenerated if the source fields or extraction definitions change.

Table S1: Directional re-routing metrics

Load	Component	Median difference [95% CI]	Raw p	BH q
SP1leg	AP (mm)	0.024 [0.014, 0.039]	< 0.001	< 0.001
SP1leg	ML (mm)	0.144 [0.124, 0.160]	< 0.001	< 0.001
SP1leg	CC (mm)	0.439 [0.416, 0.471]	< 0.001	< 0.001
LAB1	AP (mm)	-0.303 [-0.322, -0.270]	< 0.001	< 0.001
LAB1	ML (mm)	-0.007 [-0.018, 0.002]	0.254	0.254
LAB1	CC (mm)	-0.854 [-0.886, -0.817]	< 0.001	< 0.001
LAB2	AP (mm)	-0.181 [-0.207, -0.157]	< 0.001	< 0.001
LAB2	ML (mm)	0.052 [0.047, 0.061]	< 0.001	< 0.001
LAB2	CC (mm)	-0.877 [-0.900, -0.841]	< 0.001	< 0.001
LAB3	AP (mm)	0.032 [0.018, 0.043]	< 0.001	< 0.001
LAB3	ML (mm)	0.011 [0.003, 0.017]	0.007	0.007
LAB3	CC (mm)	-0.556 [-0.586, -0.537]	< 0.001	< 0.001

Table S2: Variance ratios and paired agreement

Load	Endpoint	Shape/full % [95% CI]	Material/full % [95% CI]
SP2leg	Rotation (deg)	103.89 [100.48, 107.29]	0.823 [0.589, 1.145]
SP2leg	Translation (mm)	100.37 [98.00, 102.93]	0.984 [0.745, 1.297]
SP2leg	AP (mm)	99.94 [98.11, 102.22]	0.233 [0.163, 0.333]
SP2leg	ML (mm)	104.66 [101.49, 107.06]	0.426 [0.217, 1.096]
SP2leg	CC (mm)	99.85 [97.58, 102.21]	1.069 [0.793, 1.464]
SP1leg	Rotation (deg)	101.77 [100.44, 103.13]	0.111 [0.074, 0.169]
SP1leg	Translation (mm)	100.29 [99.16, 101.54]	0.152 [0.110, 0.209]
SP1leg	AP (mm)	100.43 [97.90, 103.31]	1.047 [0.808, 1.361]
SP1leg	ML (mm)	101.13 [99.60, 102.64]	0.014 [0.008, 0.024]
SP1leg	CC (mm)	100.43 [99.20, 101.77]	0.244 [0.181, 0.333]
LAB1	Rotation (deg)	120.61 [110.69, 130.00]	5.789 [4.458, 7.466]
LAB1	Translation (mm)	107.43 [101.54, 113.20]	2.113 [1.641, 2.698]
LAB1	AP (mm)	100.68 [95.89, 105.48]	2.704 [2.061, 3.527]
LAB1	ML (mm)	109.31 [99.08, 118.28]	1.405 [0.965, 1.981]
LAB1	CC (mm)	107.81 [100.42, 115.61]	5.737 [4.496, 7.151]
LAB2	Rotation (deg)	102.97 [100.20, 105.68]	0.145 [0.089, 0.226]
LAB2	Translation (mm)	100.16 [97.41, 102.96]	0.126 [0.098, 0.164]
LAB2	AP (mm)	101.21 [99.17, 103.15]	0.089 [0.069, 0.115]
LAB2	ML (mm)	96.43 [92.57, 100.45]	1.098 [0.742, 1.606]
LAB2	CC (mm)	104.39 [98.60, 111.08]	0.568 [0.347, 0.873]
LAB3	Rotation (deg)	102.20 [100.80, 103.58]	0.030 [0.016, 0.052]
LAB3	Translation (mm)	99.44 [98.62, 100.24]	0.051 [0.039, 0.069]
LAB3	AP (mm)	100.15 [99.12, 101.13]	0.066 [0.047, 0.093]
LAB3	ML (mm)	98.32 [95.49, 101.45]	0.031 [0.017, 0.060]
LAB3	CC (mm)	98.97 [97.73, 100.14]	0.080 [0.061, 0.103]

Table S2 continued: shape/full agreement

Load	Endpoint	Identity R^2	RMSE	MAE
SP2leg	Rotation (deg)	0.9794	0.0318	0.0216
SP2leg	Translation (mm)	0.9849	0.0280	0.0198
SP2leg	AP (mm)	0.9926	0.0142	0.0082
SP2leg	ML (mm)	0.9956	0.0070	0.0041
SP2leg	CC (mm)	0.9861	0.0254	0.0187
SP1leg	Rotation (deg)	0.9974	0.0223	0.0150
SP1leg	Translation (mm)	0.9975	0.0185	0.0128
SP1leg	AP (mm)	0.9877	0.0135	0.0084
SP1leg	ML (mm)	0.9984	0.0060	0.0039
SP1leg	CC (mm)	0.9972	0.0193	0.0141
LAB1	Rotation (deg)	0.8777	0.0200	0.0149
LAB1	Translation (mm)	0.9457	0.0111	0.0080
LAB1	AP (mm)	0.9795	0.0040	0.0025
LAB1	ML (mm)	0.9472	0.0052	0.0035
LAB1	CC (mm)	0.9102	0.0134	0.0099
LAB2	Rotation (deg)	0.9835	0.0147	0.0093
LAB2	Translation (mm)	0.9842	0.0113	0.0070
LAB2	AP (mm)	0.9966	0.0066	0.0048
LAB2	ML (mm)	0.9547	0.0070	0.0046
LAB2	CC (mm)	0.9510	0.0129	0.0079
LAB3	Rotation (deg)	0.9973	0.0128	0.0077
LAB3	Translation (mm)	0.9987	0.0064	0.0044
LAB3	AP (mm)	0.9984	0.0070	0.0047
LAB3	ML (mm)	0.9985	0.0020	0.0013
LAB3	CC (mm)	0.9966	0.0059	0.0041

Table S3: Fully adjusted LAB sex models

Load	Outcome	Female minus male [95% CI]	Raw p	BH q
LAB1	Rotation (deg)	-0.009 [-0.026, 0.008]	0.283	0.566
LAB1	Translation (mm)	0.013 [-0.005, 0.031]	0.166	0.566
LAB2	Rotation (deg)	-0.017 [-0.070, 0.037]	0.543	0.736
LAB2	Translation (mm)	-0.031 [-0.078, 0.016]	0.192	0.566
LAB3	Rotation (deg)	-0.021 [-0.105, 0.062]	0.613	0.736
LAB3	Translation (mm)	0.002 [-0.060, 0.064]	0.950	0.950

Table S4: Exact scaling reparameterization using log(scale)

Load	Outcome	Male scale slope [95% CI]	Female scale slope [95% CI]
SP2leg	Rotation (deg)	-2.038 [-2.670, -1.406]	-1.816 [-2.293, -1.339]
SP2leg	Translation (mm)	-1.097 [-1.592, -0.601]	-1.205 [-1.566, -0.845]
SP1leg	Rotation (deg)	-1.866 [-2.685, -1.047]	-1.649 [-2.281, -1.018]
SP1leg	Translation (mm)	-1.379 [-2.132, -0.625]	-1.191 [-1.774, -0.609]
LAB1	Rotation (deg)	-2.747 [-3.432, -2.062]	-3.186 [-3.698, -2.674]
LAB1	Translation (mm)	-1.703 [-2.776, -0.629]	-3.010 [-3.811, -2.210]
LAB2	Rotation (deg)	-1.622 [-2.142, -1.101]	-2.021 [-2.412, -1.631]
LAB2	Translation (mm)	-2.453 [-3.538, -1.368]	-2.840 [-3.861, -1.818]
LAB3	Rotation (deg)	-2.549 [-3.144, -1.954]	-3.136 [-3.720, -2.552]
LAB3	Translation (mm)	-2.100 [-2.766, -1.434]	-2.711 [-3.477, -1.946]

Table S5: Pooled models with subject-clustered covariance

Coefficients with interaction terms are conditional: the volume main effect refers to male SP2leg. Outcome-SD coefficients divide by the pooled outcome SD; binary predictors retain their coding. Log-volume and the other continuous covariates are standardized. No robust-Wald statistic is interpreted as partial variance explained.


Outcome	Predictor	β [95% CI]	Outcome-SD β	q_{FDR}
Rotation magnitude ($ \theta $, deg)	SP1leg (vs SP2leg)	1.087 [1.048, 1.126]	1.667	< 0.001
	LAB1 (vs SP2leg)	-0.746 [-0.764, -0.728]	-1.145	< 0.001
	LAB2 (vs SP2leg)	-0.241 [-0.260, -0.223]	-0.370	< 0.001
	LAB3 (vs SP2leg)	-0.104 [-0.119, -0.089]	-0.160	< 0.001
	Female sex	0.011 [-0.046, 0.068]	0.017	0.705
	Age (z)	0.018 [0.004, 0.032]	0.027	0.017
	Log pelvic volume (z)	-0.118 [-0.142, -0.094]	-0.181	< 0.001
	SP1leg $\times$ Log volume (z)	-0.115 [-0.163, -0.067]	-0.176	< 0.001
	LAB1 $\times$ Log volume (z)	0.109 [0.090, 0.128]	0.167	< 0.001
	LAB2 $\times$ Log volume (z)	0.086 [0.067, 0.104]	0.131	< 0.001
	LAB3 $\times$ Log volume (z)	-0.015 [-0.031, 0.000]	-0.024	0.069
	Inlet AP diameter (z)	-0.009 [-0.024, 0.007]	-0.013	0.286
	Biischadic width (z)	0.048 [0.031, 0.065]	0.074	< 0.001
	Subpubic angle (z)	0.016 [-0.010, 0.043]	0.025	0.254
	Female $\times$ Log volume (z)	-0.036 [-0.068, -0.005]	-0.056	0.033
Translation magnitude ($ \mathbf{d} $, mm)	SP1leg (vs SP2leg)	0.605 [0.579, 0.630]	1.014	< 0.001
	LAB1 (vs SP2leg)	-0.907 [-0.927, -0.887]	-1.520	< 0.001
	LAB2 (vs SP2leg)	-0.792 [-0.812, -0.772]	-1.328	< 0.001
	LAB3 (vs SP2leg)	-0.478 [-0.494, -0.461]	-0.801	< 0.001
	Female sex	0.046 [0.003, 0.089]	0.077	0.045
	Age (z)	0.013 [0.002, 0.024]	0.021	0.034
	Log pelvic volume (z)	-0.103 [-0.124, -0.083]	-0.173	< 0.001
	SP1leg $\times$ Log volume (z)	-0.066 [-0.096, -0.036]	-0.111	< 0.001
	LAB1 $\times$ Log volume (z)	0.118 [0.099, 0.136]	0.197	< 0.001
	LAB2 $\times$ Log volume (z)	0.109 [0.092, 0.126]	0.183	< 0.001
	LAB3 $\times$ Log volume (z)	0.041 [0.024, 0.057]	0.068	< 0.001
	Inlet AP diameter (z)	-0.004 [-0.016, 0.008]	-0.007	0.507
	Biischadic width (z)	0.067 [0.052, 0.083]	0.113	< 0.001
	Subpubic angle (z)	0.012 [-0.008, 0.033]	0.021	0.245
	Female $\times$ Log volume (z)	-0.022 [-0.045, 0.001]	-0.036	0.077

Table S6: Extraction sensitivity

Absolute differences between rigid-only and affine sacral correction in six subjects at archive indices 0, 55, 110, 165, 220 and 277. Each contributes five loads. These are subset maxima,

	Load	Number	Max $ \Delta \text{ rotation} $ (deg)	Max $ \Delta \text{ translation} $ (mm)
not cohort-wide bounds.	SP2leg	6	0.0225	0.0049
	SP1leg	6	0.0047	0.0055
	LAB1	6	0.0013	0.0007
	LAB2	6	0.0030	0.0006
	LAB3	6	0.0229	0.0016

Supplementary Figure S1: Raw data and adjusted curves



Points are raw observations. Lines and bands are conditional geometric means and pointwise HC3 95% confidence intervals from the adjusted model, with age and the morphometric triad held at their cohort means. Thus the curves need not pass through the raw group means. Lines are restricted to each sex's observed volume range. Displayed exponents come from the same model as the curves and Table 4.