

Whole-pelvis 3D canonical deformation patterns


■ Axial ■ Bending ■ Shear ■ Torsion ■ Residual

Displacement amplitude $|u|$
0 max

Each panel: Anterior view (left) · Cranial view (right) · Bars: Shapley nonrigid shape-norm partition

Mode 1


Fitted: 42.7%



18% 23% 57%
Residual: 57.3% · Nonrigid norm: 3.3%

Mode 2


Fitted: 68.6%



21% 24% 24% 31%
Residual: 31.4% · Nonrigid norm: 5.5%

Mode 3


Fitted: 76.7%



42% 28% 23%
Residual: 23.3% · Nonrigid norm: 29.0%

Mode 4


Fitted: 74.4%



14% 44% 16% 26%
Residual: 25.6% · Nonrigid norm: 24.5%

Mode 5


Fitted: 86.3%



47% 24% 14%
Residual: 13.7% · Nonrigid norm: 73.0%

Mode 6


Fitted: 82.3%



27% 25% 27% 18%
Residual: 17.7% · Nonrigid norm: 71.9%

Mode 7


Fitted: 72.1%



17% 44% 28%
Residual: 27.9% · Nonrigid norm: 69.5%

Mode 8


Fitted: 72.1%



22% 20% 29% 28%
Residual: 27.9% · Nonrigid norm: 57.0%

Mode 9


Fitted: 91.8%



61% 18% 12%
Residual: 8.2% · Nonrigid norm: 92.1%

Mode 10


Fitted: 74.9%



35% 34% 25%
Residual: 25.1% · Nonrigid norm: 95.7%

Mode 11


Fitted: 43.1%



21% 13% 57%
Residual: 56.9% · Nonrigid norm: 71.4%

Mode 12


Fitted: 62.3%



22% 27% 38%
Residual: 37.7% · Nonrigid norm: 94.2%

Mode 13


Fitted: 48.4%



20% 19% 52%
Residual: 51.6% · Nonrigid norm: 98.3%

Mode 14


Fitted: 53.4%



16% 33% 47%
Residual: 46.6% · Nonrigid norm: 92.0%

Mode 15

Fitted: 55.6%



18% 23% 14% 44%
Residual: 44.4% · Nonrigid norm: 98.7%